



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 07:55 am GMT

PDB ID : 5N16  
Title : First Bromodomain (BD1) from *Candida albicans* Bdf1 bound to a dibenzothiazepinone (compound 1)  
Authors : Mietton, F.; Ferri, E.; Champleboux, M.; Zala, N.; Maubon, D.; Zhou, Y.; Harbut, M.; Spittler, D.; Garnaud, C.; Courcon, M.; Chauvel, M.; d'Enfert, C.; Kashemirov, B.A.; Hull, M.; Cornet, M.; McKenna, C.E.; Govin, J.; Petosa, C.  
Deposited on : 2017-02-05  
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	89% 7%
1	B	138	88% 7%
1	C	138	88% 6% 7%

*Continued on next page...*

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	D	138	 A horizontal bar chart showing the quality of chain. The bar is divided into segments: a small red segment at the start, a large green segment labeled '85%', a yellow segment labeled '8%', and a small grey segment at the end labeled '7%'. A '%' symbol is positioned above the bar on the left side.

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bromodomain-containing factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	1025	658	177	184	6	0	0	0
1	B	128	1019	655	173	185	6	0	0	0
1	C	129	1033	664	177	186	6	0	0	0
1	D	129	1033	664	177	186	6	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	ALA	-	expression tag	UNP Q5A4W8
A	191	ALA	-	expression tag	UNP Q5A4W8
A	192	GLY	-	expression tag	UNP Q5A4W8
B	190	ALA	-	expression tag	UNP Q5A4W8
B	191	ALA	-	expression tag	UNP Q5A4W8
B	192	GLY	-	expression tag	UNP Q5A4W8
C	190	ALA	-	expression tag	UNP Q5A4W8
C	191	ALA	-	expression tag	UNP Q5A4W8
C	192	GLY	-	expression tag	UNP Q5A4W8
D	190	ALA	-	expression tag	UNP Q5A4W8
D	191	ALA	-	expression tag	UNP Q5A4W8
D	192	GLY	-	expression tag	UNP Q5A4W8

- Molecule 2 is 5-cyclopropyl-2-(5-pyrazin-2-yl-1,2,4-oxadiazol-3-yl)benzo[b][1,4]benzothiazepine-6-one (three-letter code: 8FN) (formula: C<sub>22</sub>H<sub>15</sub>N<sub>5</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total	C	N	O	S	0	0
			30	22	5	2	1		
2	A	1	Total	C	N	O	S	0	0
			30	22	5	2	1		
2	C	1	Total	C	N	O	S	0	0
			30	22	5	2	1		
2	D	1	Total	C	N	O	S	0	0
			30	22	5	2	1		

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ni		
3	A	2	Total	Ni	0	0
			2	2		
3	B	1	Total	Ni	0	0
			1	1		
3	C	1	Total	Ni	0	0
			1	1		
3	D	1	Total	Ni	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	165	Total O 165 165	0	0
5	B	176	Total O 176 176	0	0
5	C	132	Total O 132 132	0	0
5	D	175	Total O 175 175	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bromodomain-containing factor 1




- Molecule 1: Bromodomain-containing factor 1



- Molecule 1: Bromodomain-containing factor 1



- Molecule 1: Bromodomain-containing factor 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.44Å 77.29Å 102.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.74 – 1.76 42.74 – 1.68	Depositor EDS
% Data completeness (in resolution range)	99.5 (42.74-1.76) 98.3 (42.74-1.68)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 1.68Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.178 , 0.215 0.179 , 0.215	Depositor DCC
$R_{free}$ test set	3419 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtrriage
Anisotropy	0.281	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.021 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4889	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NI, 8FN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1050	0.47	0/1424
1	B	0.36	0/1044	0.49	0/1417
1	C	0.32	0/1058	0.46	0/1435
1	D	0.37	0/1058	0.49	0/1435
All	All	0.34	0/4210	0.48	0/5711

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1025	0	1044	6	0
1	B	1019	0	1036	5	0
1	C	1033	0	1058	6	0
1	D	1033	0	1058	10	0
2	A	60	0	0	0	0
2	C	30	0	0	0	0
2	D	30	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	C	6	0	7	0	0
5	A	165	0	0	1	1
5	B	176	0	0	2	5
5	C	132	0	0	1	0
5	D	175	0	0	5	5
All	All	4889	0	4203	24	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:300:MET:SD	5:C:609:HOH:O	2.26	0.92
1:B:220:ASN:OD1	5:B:501:HOH:O	2.05	0.74
1:D:212:GLN:NE2	5:D:502:HOH:O	2.25	0.70
1:D:318:LYS:NZ	5:D:501:HOH:O	2.21	0.67
1:A:289:LYS:NZ	5:A:502:HOH:O	2.32	0.62
1:B:319:GLU:OE2	1:B:319:GLU:HA	1.98	0.61
1:B:212:GLN:HG2	1:B:216:LYS:HE3	1.85	0.58
1:D:317:PRO:HD2	1:D:320:LEU:HG	1.88	0.54
1:A:204:ASN:ND2	1:D:263:ARG:HG3	2.27	0.50
1:D:206:PRO:CB	1:D:271:GLU:HG2	2.43	0.49
1:D:285:LYS:NZ	5:D:505:HOH:O	2.37	0.48
1:D:198:GLN:HG3	5:D:642:HOH:O	2.14	0.47
1:A:201:ASP:OD2	1:D:267:LEU:HD21	2.14	0.47
1:C:230:GLU:O	1:C:300:MET:HE3	2.14	0.47
1:C:222:ILE:HG22	1:C:226:LYS:HD2	1.97	0.47
1:B:299:LYS:HZ3	1:B:303:ASN:HD21	1.63	0.47
1:D:303:ASN:HB3	5:D:534:HOH:O	2.15	0.46
1:A:203:ASN:O	1:D:266:ASN:HB3	2.17	0.45
1:C:218:VAL:HG23	1:C:265:ILE:HD13	1.97	0.45
1:A:318:LYS:HB3	1:A:318:LYS:HE2	1.83	0.44
1:C:300:MET:HE3	1:C:300:MET:HB3	1.81	0.41
1:C:272:ASP:OD1	1:C:273:VAL:N	2.54	0.41
1:B:216:LYS:NZ	5:B:503:HOH:O	2.32	0.40
1:A:226:LYS:HB3	1:A:226:LYS:HE2	1.97	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:637:HOH:O	5:D:653:HOH:O[3_555]	2.13	0.07
5:A:542:HOH:O	5:A:660:HOH:O[4_555]	2.14	0.06
5:B:566:HOH:O	5:D:640:HOH:O[1_565]	2.15	0.05
5:B:533:HOH:O	5:D:618:HOH:O[1_565]	2.16	0.04
5:B:625:HOH:O	5:D:606:HOH:O[4_455]	2.17	0.03
5:B:644:HOH:O	5:D:501:HOH:O[4_455]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/138 (92%)	127 (100%)	0	0	100	100
1	B	126/138 (91%)	126 (100%)	0	0	100	100
1	C	127/138 (92%)	126 (99%)	1 (1%)	0	100	100
1	D	127/138 (92%)	127 (100%)	0	0	100	100
All	All	507/552 (92%)	506 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/121 (95%)	115 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	115/121 (95%)	112 (97%)	3 (3%)	46	23
1	C	117/121 (97%)	117 (100%)	0	100	100
1	D	117/121 (97%)	116 (99%)	1 (1%)	78	67
All	All	464/484 (96%)	460 (99%)	4 (1%)	78	67

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	299	LYS
1	B	310	LYS
1	B	319	GLU
1	D	320	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	204	ASN
1	D	204	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	8FN	A	404	-	32,35,35	2.05	5 (15%)	42,51,51	2.26	12 (28%)
2	8FN	C	401	-	32,35,35	2.43	6 (18%)	42,51,51	2.45	10 (23%)
2	8FN	A	401	-	32,35,35	2.45	6 (18%)	42,51,51	2.66	9 (21%)
4	GOL	C	402	2	5,5,5	0.75	0	5,5,5	2.24	2 (40%)
2	8FN	D	401	4	32,35,35	2.12	5 (15%)	42,51,51	2.53	16 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8FN	A	404	-	-	2/10/30/30	0/5/6/6
2	8FN	C	401	-	-	2/10/30/30	0/5/6/6
2	8FN	A	401	-	-	2/10/30/30	0/5/6/6
4	GOL	C	402	2	-	2/4/4/4	-
2	8FN	D	401	4	-	2/10/30/30	0/5/6/6

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	8FN	C4-S1	-9.60	1.69	1.78
2	C	401	8FN	C4-S1	-8.94	1.69	1.78
2	A	401	8FN	C13-S1	-8.58	1.70	1.78
2	C	401	8FN	C13-S1	-8.55	1.70	1.78
2	D	401	8FN	C4-S1	-8.54	1.70	1.78
2	A	404	8FN	C4-S1	-7.64	1.71	1.78
2	A	404	8FN	C13-S1	-6.97	1.71	1.78
2	D	401	8FN	C13-S1	-6.24	1.72	1.78
2	C	401	8FN	C7-N1	-3.69	1.32	1.39
2	D	401	8FN	C7-N1	-3.23	1.33	1.39
2	A	404	8FN	C7-N1	-2.91	1.34	1.39
2	A	404	8FN	C22-C20	2.65	1.54	1.48
2	A	401	8FN	C7-N1	-2.62	1.34	1.39
2	A	404	8FN	C21-C20	2.42	1.54	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	8FN	C21-C20	2.40	1.54	1.48
2	C	401	8FN	C22-C20	2.35	1.54	1.48
2	D	401	8FN	C21-C20	2.31	1.53	1.48
2	D	401	8FN	C22-C20	2.31	1.53	1.48
2	C	401	8FN	C8-N1	-2.26	1.40	1.43
2	A	401	8FN	C8-N1	-2.21	1.40	1.43
2	A	401	8FN	C22-C20	2.19	1.53	1.48
2	A	401	8FN	C21-C20	2.14	1.53	1.48

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	8FN	C22-C20-N1	-9.26	108.94	118.29
2	C	401	8FN	C22-C20-N1	-8.73	109.48	118.29
2	A	401	8FN	C14-N2-C15	8.10	107.69	101.13
2	D	401	8FN	C21-C20-N1	-7.94	110.27	118.29
2	C	401	8FN	C14-N2-C15	7.31	107.04	101.13
2	D	401	8FN	C22-C20-N1	-7.26	110.96	118.29
2	A	401	8FN	C21-C20-N1	-7.13	111.09	118.29
2	A	404	8FN	C14-N2-C15	7.01	106.80	101.13
2	A	404	8FN	C22-C20-N1	-7.00	111.22	118.29
2	C	401	8FN	C21-C20-N1	-6.69	111.54	118.29
2	D	401	8FN	C14-N2-C15	5.89	105.89	101.13
2	A	404	8FN	C21-C20-N1	-5.47	112.77	118.29
2	A	401	8FN	C11-C14-N3	4.99	125.61	119.12
4	C	402	GOL	C3-C2-C1	-3.90	96.52	111.70
2	A	401	8FN	C5-C7-N1	3.59	124.87	118.39
2	A	404	8FN	C9-C8-N1	-3.59	116.52	120.48
2	D	401	8FN	C9-C8-N1	-3.51	116.61	120.48
2	A	404	8FN	C13-C8-N1	3.27	123.92	120.54
2	A	404	8FN	C5-C7-N1	3.19	124.15	118.39
2	C	401	8FN	C9-C8-N1	-3.12	117.03	120.48
2	D	401	8FN	C17-C16-C15	-3.05	115.32	121.22
2	C	401	8FN	C11-C14-N3	3.03	123.06	119.12
2	D	401	8FN	O1-C7-C5	-2.94	115.06	120.01
2	D	401	8FN	C5-C7-N1	2.94	123.70	118.39
2	A	401	8FN	O1-C7-C5	-2.80	115.30	120.01
2	A	401	8FN	C9-C8-N1	-2.77	117.42	120.48
2	D	401	8FN	C11-C14-N2	2.76	127.16	123.71
2	D	401	8FN	C17-C16-N5	2.76	125.08	120.46
2	D	401	8FN	C13-C8-N1	2.67	123.30	120.54
2	D	401	8FN	C9-C10-C11	-2.65	117.31	121.13

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	401	8FN	C12-C13-C8	-2.65	118.34	120.77
2	D	401	8FN	C11-C14-N3	2.58	122.48	119.12
2	C	401	8FN	C10-C11-C12	2.55	121.77	118.16
2	A	404	8FN	O1-C7-C5	-2.51	115.79	120.01
2	C	401	8FN	C13-C8-N1	2.50	123.13	120.54
2	D	401	8FN	C15-C16-N5	2.48	119.58	116.75
2	C	401	8FN	C18-C19-N5	-2.44	118.65	122.17
2	D	401	8FN	C10-C11-C12	2.43	121.61	118.16
2	A	404	8FN	C11-C14-N2	2.41	126.73	123.71
2	A	401	8FN	C10-C11-C12	2.41	121.57	118.16
4	C	402	GOL	O3-C3-C2	2.37	121.55	110.20
2	A	401	8FN	C13-C8-N1	2.34	122.96	120.54
2	D	401	8FN	C16-C17-N4	-2.33	117.64	121.85
2	A	404	8FN	C10-C11-C12	2.32	121.45	118.16
2	A	404	8FN	C11-C14-N3	2.21	122.00	119.12
2	C	401	8FN	C18-N4-C17	2.14	120.56	116.85
2	C	401	8FN	C11-C14-N2	2.12	126.36	123.71
2	A	404	8FN	C9-C10-C11	-2.11	118.10	121.13
2	A	404	8FN	C12-C13-C8	-2.10	118.85	120.77

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	8FN	C22-C20-N1-C8
2	A	401	8FN	C21-C20-N1-C8
2	A	404	8FN	C22-C20-N1-C8
2	A	404	8FN	C21-C20-N1-C8
2	C	401	8FN	C22-C20-N1-C8
2	C	401	8FN	C21-C20-N1-C8
2	D	401	8FN	C22-C20-N1-C8
2	D	401	8FN	C21-C20-N1-C8
4	C	402	GOL	C1-C2-C3-O3
4	C	402	GOL	O2-C2-C3-O3

There are no ring outliers.



No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	129/138 (93%)	-0.21	1 (0%) 86 90	16, 24, 39, 65	0
1	B	128/138 (92%)	-0.32	0 100 100	15, 22, 38, 53	0
1	C	129/138 (93%)	0.16	7 (5%) 25 32	15, 29, 46, 67	0
1	D	129/138 (93%)	-0.09	2 (1%) 72 79	14, 21, 39, 64	0
All	All	515/552 (93%)	-0.11	10 (1%) 66 74	14, 24, 43, 67	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	320	LEU	6.6
1	C	319	GLU	4.4
1	A	320	LEU	3.3
1	C	203	ASN	3.1
1	D	319	GLU	2.6
1	C	204	ASN	2.5
1	C	313	VAL	2.5
1	C	316	PRO	2.4
1	C	212	GLN	2.2
1	C	210	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	C	402	6/6	0.81	0.23	19,22,27,57	0
3	NI	A	403	1/1	0.89	0.17	78,78,78,78	0
2	8FN	D	401	30/30	0.94	0.09	12,17,27,32	0
2	8FN	A	401	30/30	0.95	0.09	14,20,25,30	0
2	8FN	A	404	30/30	0.96	0.08	14,17,25,29	0
2	8FN	C	401	30/30	0.97	0.08	14,18,22,24	0
3	NI	B	401	1/1	0.99	0.07	31,31,31,31	0
3	NI	C	403	1/1	0.99	0.09	30,30,30,30	0
3	NI	D	402	1/1	0.99	0.10	29,29,29,29	0
3	NI	A	402	1/1	0.99	0.06	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 8FN A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 8FN A 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.