



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 11:12 PM EDT

PDB ID : 5NJ2
Title : Crystal structure of BlaC from Mycobacterium tuberculosis bound to phosphate
Authors : Tassoni, R.; Pannu, N.S.; Ubbink, M.
Deposited on : 2017-03-27
Resolution : 1.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
1	B	274	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	401	-	X	-	-
2	PO4	B	401	-	X	-	-
3	ACT	A	404	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	272	Total	C	N	O	S	0	3	0
			2076	1300	376	394	6			
1	B	265	Total	C	N	O	S	0	1	0
			1992	1247	352	387	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	MET	-	initiating methionine	UNP A0A0T9EA39
A	295	LEU	-	expression tag	UNP A0A0T9EA39
A	296	GLU	-	expression tag	UNP A0A0T9EA39
A	297	HIS	-	expression tag	UNP A0A0T9EA39
A	298	HIS	-	expression tag	UNP A0A0T9EA39
A	299	HIS	-	expression tag	UNP A0A0T9EA39
A	300	HIS	-	expression tag	UNP A0A0T9EA39
A	301	HIS	-	expression tag	UNP A0A0T9EA39
A	302	HIS	-	expression tag	UNP A0A0T9EA39
B	28	MET	-	initiating methionine	UNP A0A0T9EA39
B	295	LEU	-	expression tag	UNP A0A0T9EA39
B	296	GLU	-	expression tag	UNP A0A0T9EA39
B	297	HIS	-	expression tag	UNP A0A0T9EA39
B	298	HIS	-	expression tag	UNP A0A0T9EA39
B	299	HIS	-	expression tag	UNP A0A0T9EA39
B	300	HIS	-	expression tag	UNP A0A0T9EA39
B	301	HIS	-	expression tag	UNP A0A0T9EA39
B	302	HIS	-	expression tag	UNP A0A0T9EA39

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 2-{2-[2-2-(METHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHANOL (three-letter code: ETE) (formula: C₉H₂₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			14	9	5		

- Molecule 5 is 3,6,9,12,15-PENTAOXAHEPTADECAN-1-OL (three-letter code: AE4) (formula: C₁₂H₂₆O₆).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	C O	0	0
			15	10 5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	147	Total	O	0	0
			147	147		
6	B	141	Total	O	0	0
			141	141		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	39.59Å 41.68Å 76.84Å 101.28° 90.11° 90.43°	Depositor
Resolution (Å)	75.36 – 1.19 40.88 – 1.19	Depositor EDS
% Data completeness (in resolution range)	62.7 (75.36-1.19) 62.7 (40.88-1.19)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.19Å)	Xtrriage
Refinement program	REFMAC 7.0.033	Depositor
R, R_{free}	0.167 , 0.190 0.176 , 0.195	Depositor DCC
R_{free} test set	4773 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	10.1	Xtrriage
Anisotropy	0.018	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4416	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, AE4, ACT, ETE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.09	3/2131 (0.1%)	1.21	22/2908 (0.8%)
1	B	1.00	1/2034 (0.0%)	1.20	17/2777 (0.6%)
All	All	1.04	4/4165 (0.1%)	1.21	39/5685 (0.7%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	48	TYR	CE2-CZ	-7.34	1.29	1.38
1	A	126	ARG	CZ-NH2	-5.74	1.25	1.33
1	B	253	SER	C-N	5.09	1.44	1.34
1	A	167	GLU	CD-OE2	5.06	1.31	1.25

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	ARG	NE-CZ-NH1	14.62	127.61	120.30
1	B	65	ARG	NE-CZ-NH1	13.61	127.10	120.30
1	A	126	ARG	NE-CZ-NH2	-11.98	114.31	120.30
1	B	65	ARG	NE-CZ-NH2	-10.57	115.01	120.30
1	B	126	ARG	NE-CZ-NH2	9.35	124.97	120.30
1	B	278	ARG	NE-CZ-NH2	8.90	124.75	120.30
1	A	101	ARG	NE-CZ-NH2	-8.74	115.93	120.30
1	B	126	ARG	NE-CZ-NH1	-8.63	115.98	120.30
1	A	40	ARG	NE-CZ-NH1	8.31	124.46	120.30
1	A	65	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	A	267	ASP	CB-CG-OD1	7.61	125.15	118.30
1	A	101	ARG	NE-CZ-NH1	7.61	124.10	120.30
1	B	267	ASP	CB-CG-OD1	7.53	125.08	118.30
1	B	101	ARG	NE-CZ-NH1	7.31	123.96	120.30
1	A	65	ARG	NE-CZ-NH2	-7.30	116.65	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40	ARG	NE-CZ-NH1	-7.29	116.66	120.30
1	A	204	ASP	CB-CG-OD1	7.03	124.63	118.30
1	B	204	ASP	CB-CG-OD1	6.41	124.07	118.30
1	A	244	ARG	NE-CZ-NH2	-6.41	117.10	120.30
1	B	37	LEU	CB-CG-CD2	6.26	121.64	111.00
1	A	282	LEU	CB-CG-CD2	6.14	121.43	111.00
1	A	178	ASP	CB-CG-OD2	-6.00	112.89	118.30
1	A	186[A]	HIS	N-CA-CB	-5.91	99.95	110.60
1	A	186[B]	HIS	N-CA-CB	-5.91	99.95	110.60
1	A	165	ASP	CB-CG-OD1	5.89	123.60	118.30
1	A	40	ARG	NE-CZ-NH2	-5.69	117.45	120.30
1	B	235	ASP	CB-CG-OD2	5.59	123.33	118.30
1	A	126	ARG	CD-NE-CZ	5.50	131.30	123.60
1	B	159	ASP	CB-CG-OD2	-5.44	113.40	118.30
1	A	215	ARG	NE-CZ-NH2	5.38	122.99	120.30
1	B	98	ASP	CB-CG-OD2	-5.37	113.47	118.30
1	A	90	ASP	CB-CG-OD2	-5.33	113.51	118.30
1	B	178	ASP	CB-CG-OD1	5.27	123.05	118.30
1	A	86	LEU	CB-CG-CD1	5.25	119.92	111.00
1	A	173	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	B	222	ARG	NE-CZ-NH2	5.21	122.91	120.30
1	A	230	ASP	CB-CG-OD1	5.21	122.99	118.30
1	B	101	ARG	NE-CZ-NH2	-5.19	117.70	120.30
1	B	178	ASP	CB-CG-OD2	-5.10	113.71	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2076	0	2040	18	0
1	B	1992	0	1965	15	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	12	0	9	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	4	0	3	0	0
4	B	14	0	20	1	0
5	B	15	0	16	8	0
6	A	147	0	0	1	0
6	B	141	0	0	0	0
All	All	4416	0	4053	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:ALA:HB2	5:B:405:AE4:H121	1.45	0.98
1:A:264[B]:VAL:HG12	1:A:282:LEU:HD22	1.46	0.96
1:B:264[A]:VAL:HG12	1:B:282:LEU:HD22	1.56	0.86
1:A:116:THR:H	1:A:119:GLN:HE21	1.21	0.83
1:B:116:THR:H	1:B:119:GLN:HE21	1.28	0.81
1:A:48:TYR:CE1	1:A:186[B]:HIS:CD2	2.69	0.80
1:A:48:TYR:CE1	1:A:186[B]:HIS:HD2	2.07	0.73
1:A:88:HIS:HE1	1:A:139:ASP:OD2	1.80	0.64
1:B:88:HIS:HE1	1:B:139:ASP:OD2	1.81	0.64
1:B:61:ARG:H	5:B:405:AE4:C8	2.12	0.60
1:B:31:ALA:HB2	5:B:405:AE4:C12	2.25	0.60
1:A:264[B]:VAL:CG1	1:A:282:LEU:HD22	2.27	0.60
1:B:61:ARG:H	5:B:405:AE4:H8C2	1.68	0.58
1:A:48:TYR:CZ	1:A:186[B]:HIS:HD2	2.22	0.57
1:A:48:TYR:CD1	1:A:186[B]:HIS:CD2	2.97	0.53
1:A:173:ARG:HE	3:A:404:ACT:H3	1.74	0.53
1:A:173:ARG:HE	3:A:404:ACT:CH3	2.23	0.52
1:B:60:TYR:HD1	5:B:405:AE4:H122	1.77	0.49
1:B:210:THR:HG21	4:B:404:ETE:H141	1.93	0.49
1:A:241:ASP:OD2	3:A:404:ACT:CH3	2.61	0.48
1:A:48:TYR:CD1	1:A:59:GLU:HG2	2.48	0.48
1:B:31:ALA:CB	5:B:405:AE4:C12	2.94	0.45
1:B:31:ALA:CB	5:B:405:AE4:H121	2.31	0.45
1:B:116:THR:H	1:B:119:GLN:NE2	2.06	0.44
1:B:264[A]:VAL:HG12	1:B:282:LEU:CD2	2.39	0.44
1:B:264[A]:VAL:CG1	1:B:282:LEU:HD22	2.36	0.43
1:A:241:ASP:OD2	3:A:404:ACT:H3	2.20	0.42
1:A:116:THR:H	1:A:119:GLN:NE2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:59:GLU:H	5:B:405:AE4:C17	2.33	0.41
1:A:110:HIS:HD2	6:A:502:HOH:O	2.03	0.41
1:A:264[B]:VAL:HG12	1:A:282:LEU:CD2	2.32	0.41
1:A:241:ASP:OD2	3:A:404:ACT:H2	2.21	0.41
1:A:48:TYR:HD1	1:A:59:GLU:HG2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/274 (100%)	268 (98%)	5 (2%)	0	100	100
1	B	264/274 (96%)	261 (99%)	3 (1%)	0	100	100
All	All	537/548 (98%)	529 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/211 (100%)	209 (99%)	3 (1%)	67	32
1	B	203/211 (96%)	199 (98%)	4 (2%)	55	17
All	All	415/422 (98%)	408 (98%)	7 (2%)	60	24

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LEU
1	A	98	ASP
1	A	235	ASP
1	B	86	LEU
1	B	113	THR
1	B	128	SER
1	B	235	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	HIS
1	A	88	HIS
1	A	109	GLN
1	A	110	HIS
1	A	119	GLN
1	A	199	ASN
1	B	88	HIS
1	B	108	GLN
1	B	119	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	A	403	-	3,3,3	0.83	0	3,3,3	0.66	0
3	ACT	A	404	-	3,3,3	0.88	0	3,3,3	0.18	0
2	PO4	B	402	-	4,4,4	0.79	0	6,6,6	0.94	0
2	PO4	B	401	-	4,4,4	1.47	1 (25%)	6,6,6	2.38	3 (50%)
3	ACT	A	402	-	3,3,3	1.09	0	3,3,3	1.06	0
3	ACT	B	403	-	3,3,3	0.57	0	3,3,3	0.76	0
4	ETE	B	404	-	13,13,13	0.77	0	12,12,12	1.20	1 (8%)
2	PO4	A	401	-	4,4,4	0.82	0	6,6,6	4.41	4 (66%)
5	AE4	B	405	-	14,14,17	0.87	0	13,13,16	1.58	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ETE	B	404	-	-	4/11/11/11	-
5	AE4	B	405	-	-	7/12/12/15	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	PO4	P-O2	-2.22	1.47	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PO4	O3-P-O2	8.19	134.25	107.97
2	A	401	PO4	O3-P-O1	-4.71	93.66	110.89
2	A	401	PO4	O2-P-O1	-4.20	95.52	110.89
2	B	401	PO4	O3-P-O2	3.63	119.64	107.97
2	B	401	PO4	O3-P-O1	-2.94	100.14	110.89
2	A	401	PO4	O4-P-O1	-2.77	100.74	110.89
4	B	404	ETE	C24-OH4-C13	2.70	124.97	113.29
5	B	405	AE4	O10-C11-C12	2.61	122.17	110.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	405	AE4	O13-C12-C11	2.48	121.56	110.39
2	B	401	PO4	O4-P-O1	-2.39	102.17	110.89
5	B	405	AE4	O7-C6-C5	2.15	120.08	110.39
5	B	405	AE4	O7-C8-C9	2.12	119.95	110.39

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	404	ETE	OH2-C12-C22-OH3
5	B	405	AE4	O10-C11-C12-O13
4	B	404	ETE	OH5-C14-C24-OH4
5	B	405	AE4	O4-C5-C6-O7
5	B	405	AE4	C12-C11-O10-C9
5	B	405	AE4	C9-C8-O7-C6
5	B	405	AE4	C14-C15-O16-C17
5	B	405	AE4	C11-C12-O13-C14
4	B	404	ETE	OH6-C15-C25-OH5
4	B	404	ETE	C15-C25-OH5-C14
5	B	405	AE4	C15-C14-O13-C12

There are no ring outliers.

3 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	ACT	5	0
4	B	404	ETE	1	0
5	B	405	AE4	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/274 (99%)	0.22	6 (2%) 62 61	5, 9, 22, 39	0
1	B	265/274 (96%)	0.10	5 (1%) 66 67	6, 11, 23, 36	0
All	All	537/548 (97%)	0.16	11 (2%) 65 66	5, 10, 23, 39	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	301	HIS	3.9
1	B	113	THR	3.9
1	A	53	GLY	3.4
1	B	53	GLY	3.1
1	A	143	PRO	2.9
1	A	103	ILE	2.9
1	B	103	ILE	2.4
1	A	299	HIS	2.4
1	B	101	ARG	2.4
1	A	54	THR	2.3
1	B	109	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	A	404	4/4	0.69	0.16	27,29,32,37	0
5	AE4	B	405	15/18	0.75	0.19	22,31,42,46	0
4	ETE	B	404	14/14	0.79	0.19	29,33,39,43	0
3	ACT	A	403	4/4	0.82	0.10	27,29,31,41	0
3	ACT	A	402	4/4	0.93	0.10	17,20,21,22	0
3	ACT	B	403	4/4	0.96	0.14	18,19,21,21	0
2	PO4	A	401	5/5	0.97	0.08	13,16,19,21	0
2	PO4	B	401	5/5	0.97	0.09	17,18,21,27	0
2	PO4	B	402	5/5	0.97	0.17	25,28,32,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.