



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 08:42 AM EDT

PDB ID : 5NS7  
Title : Crystal structure of beta-glucosidase BglM-G1 mutant H75R from marine metagenome  
Authors : Mhaindarkar, D.C.; Gasper, R.; Lupilova, N.; Leichert, L.I.; Hofmann, E.  
Deposited on : 2017-04-25  
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	 7% 93%
1	B	454	 2% 93%
1	C	454	 3% 92%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 22024 atoms, of which 10399 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-glucosidase M - G1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	439	Total 6996	C 2306	H 3424	N 593	O 659	S 14	2	8	0
1	B	440	Total 7003	C 2307	H 3428	N 595	O 659	S 14	2	7	0
1	C	440	Total 7010	C 2310	H 3430	N 595	O 661	S 14	2	8	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	12	3	6	3	0	0
3	A	1	13	3	7	3	0	0
3	A	1	12	3	6	3	0	0
3	A	1	12	3	6	3	0	0
3	A	1	13	3	7	3	0	0
3	B	1	13	3	7	3	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			13	3	7	3		
3	B	1	Total	C	H	O	0	0
			12	3	6	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			12	3	6	3		
3	C	1	Total	C	H	O	0	0
			12	3	6	3		
3	C	1	Total	C	H	O	0	0
			13	3	7	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			12	3	6	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	185	Total	O	0	0
			185	185		
4	B	317	Total	O	0	0
			317	317		
4	C	254	Total	O	0	0
			254	254		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: beta-glucosidase M - G1



- Molecule 1: beta-glucosidase M - G1



- Molecule 1: beta-glucosidase M - G1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.92Å 137.92Å 98.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.76 – 1.54 48.76 – 1.54	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.76-1.54) 99.9 (48.76-1.54)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.21 (at 1.54Å)	Xtrriage
Refinement program	PHENIX dev_2621	Depositor
R, $R_{free}$	0.159 , 0.172 0.159 , 0.172	Depositor DCC
$R_{free}$ test set	13515 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.1	Xtrriage
Anisotropy	0.005	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	22024	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3711	0.59	0/5034
1	B	0.48	0/3712	0.66	1/5037 (0.0%)
1	C	0.46	0/3720	0.64	1/5048 (0.0%)
All	All	0.45	0/11143	0.63	2/15119 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	82	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	B	435	ARG	NE-CZ-NH2	-5.46	117.57	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3572	3424	3430	12	0
1	B	3575	3428	3433	14	0
1	C	3580	3430	3437	15	0
2	A	10	0	0	0	0
2	B	15	0	0	0	0
2	C	15	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	30	32	39	1	0
3	B	42	52	56	3	0
3	C	30	33	40	3	0
4	A	185	0	0	3	0
4	B	317	0	0	4	0
4	C	254	0	0	4	0
All	All	11625	10399	10435	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:339:GLU:OE1	4:C:601:HOH:O	1.84	0.95
1:B:129:LYS:NZ	4:B:601:HOH:O	2.12	0.82
1:C:339:GLU:OE2	4:C:602:HOH:O	1.97	0.82
1:C:210:SER:OG	1:C:212[A]:GLU:OE2	1.99	0.80
1:C:207[B]:ARG:NH2	1:C:282:PRO:O	2.18	0.73
1:A:210:SER:OG	1:A:212[A]:GLU:OE2	2.14	0.65
1:B:210:SER:OG	1:B:212[A]:GLU:OE2	2.14	0.65
1:C:274:GLY:H	3:C:506:GOL:H11	1.64	0.62
1:C:276:MET:HB2	3:C:506:GOL:H2	1.81	0.62
1:C:171:LEU:HB3	1:C:178[B]:MET:HG2	1.84	0.57
1:A:207[B]:ARG:NH2	1:A:282:PRO:O	2.33	0.55
1:C:335:ARG:NH2	4:C:601:HOH:O	2.38	0.55
1:A:357:GLU:HG3	1:A:359:VAL:HG23	1.88	0.54
1:C:160:ILE:HD12	4:C:630:HOH:O	2.09	0.53
3:C:504:GOL:H31	3:C:507:GOL:H2	1.90	0.52
1:B:4:TRP:CE2	1:C:383:LYS:HE3	2.44	0.52
1:A:120:TRP:CZ3	1:A:178[B]:MET:HE1	2.45	0.52
1:B:178[B]:MET:CE	4:B:744:HOH:O	2.58	0.51
1:B:160:ILE:HD12	4:B:647:HOH:O	2.09	0.51
1:A:373:GLU:OE1	3:A:506:GOL:H12	2.11	0.50
1:A:120:TRP:CE3	1:A:178[B]:MET:HE1	2.47	0.50
1:A:383:LYS:HE3	1:C:4:TRP:CE2	2.46	0.50
1:B:120:TRP:CE3	1:B:178[B]:MET:HE1	2.47	0.49
1:B:120:TRP:CZ3	1:B:178[B]:MET:HE1	2.50	0.47
1:A:178[B]:MET:CE	4:A:749:HOH:O	2.63	0.46
1:B:178[B]:MET:HE2	4:B:744:HOH:O	2.16	0.46
1:B:207[B]:ARG:NH2	1:B:282:PRO:O	2.45	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:256:LYS:O	1:C:256:LYS:HD2	2.15	0.46
1:B:291:TYR:HB2	3:B:509:GOL:H32	1.98	0.45
1:A:4:TRP:CE2	1:B:383:LYS:HE3	2.52	0.45
1:A:160:ILE:HD12	4:A:641:HOH:O	2.18	0.44
1:C:259:GLU:HG2	1:C:260:ASP:N	2.33	0.43
1:A:357:GLU:HG2	1:A:357:GLU:O	2.18	0.43
1:B:137:CYS:HA	1:B:140:TRP:CE3	2.54	0.42
1:C:137:CYS:HA	1:C:140:TRP:CE3	2.53	0.42
1:B:222:THR:OG1	3:B:509:GOL:H2	2.19	0.42
1:A:178[B]:MET:HE2	4:A:749:HOH:O	2.18	0.41
1:B:222:THR:OG1	3:B:509:GOL:C2	2.69	0.41
1:C:0:HIS:O	1:C:1:MET:HB2	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/454 (98%)	435 (98%)	8 (2%)	0	100	100
1	B	445/454 (98%)	437 (98%)	8 (2%)	0	100	100
1	C	446/454 (98%)	438 (98%)	8 (2%)	0	100	100
All	All	1334/1362 (98%)	1310 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/379 (98%)	370 (99%)	3 (1%)	81	64
1	B	373/379 (98%)	371 (100%)	2 (0%)	88	77
1	C	374/379 (99%)	370 (99%)	4 (1%)	73	51
All	All	1120/1137 (98%)	1111 (99%)	9 (1%)	81	64

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	LYS
1	A	338	LYS
1	A	435	ARG
1	B	256	LYS
1	B	435	ARG
1	C	0	HIS
1	C	256	LYS
1	C	259	GLU
1	C	435	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	505	-	5,5,5	1.38	1 (20%)	5,5,5	1.05	0
3	GOL	A	504	-	5,5,5	1.21	0	5,5,5	1.04	0
3	GOL	C	508	-	5,5,5	1.20	0	5,5,5	0.83	0
2	SO4	A	501	-	4,4,4	0.13	0	6,6,6	0.11	0
2	SO4	C	501	-	4,4,4	0.13	0	6,6,6	0.09	0
2	SO4	C	503	-	4,4,4	0.14	0	6,6,6	0.37	0
3	GOL	B	507	-	5,5,5	1.01	0	5,5,5	0.94	0
3	GOL	B	509	-	5,5,5	1.14	0	5,5,5	0.76	0
3	GOL	C	507	-	5,5,5	0.80	0	5,5,5	1.03	0
2	SO4	A	502	-	4,4,4	0.17	0	6,6,6	0.07	0
3	GOL	C	506	-	5,5,5	1.18	0	5,5,5	0.98	0
3	GOL	A	505	-	5,5,5	0.92	0	5,5,5	0.89	0
3	GOL	C	504	-	5,5,5	1.21	0	5,5,5	1.13	0
3	GOL	B	510	-	5,5,5	1.13	0	5,5,5	0.88	0
2	SO4	B	501	-	4,4,4	0.23	0	6,6,6	0.68	0
2	SO4	C	502	-	4,4,4	0.14	0	6,6,6	0.18	0
3	GOL	B	506	-	5,5,5	1.18	0	5,5,5	0.93	0
3	GOL	B	504	-	5,5,5	1.39	1 (20%)	5,5,5	1.07	1 (20%)
3	GOL	B	508	-	5,5,5	1.03	0	5,5,5	1.05	0
2	SO4	B	502	-	4,4,4	0.15	0	6,6,6	0.16	0
3	GOL	A	507	-	5,5,5	1.08	0	5,5,5	1.03	0
2	SO4	B	503	-	4,4,4	0.15	0	6,6,6	0.17	0
3	GOL	C	505	-	5,5,5	1.02	0	5,5,5	0.93	0
3	GOL	A	506	-	5,5,5	0.92	0	5,5,5	0.96	0
3	GOL	A	503	-	5,5,5	1.27	0	5,5,5	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	506	-	-	4/4/4/4	-
3	GOL	C	507	-	-	2/4/4/4	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	505	-	-	0/4/4/4	-
3	GOL	B	505	-	-	1/4/4/4	-
3	GOL	C	504	-	-	2/4/4/4	-
3	GOL	B	510	-	-	0/4/4/4	-
3	GOL	A	507	-	-	3/4/4/4	-
3	GOL	B	507	-	-	2/4/4/4	-
3	GOL	B	506	-	-	0/4/4/4	-
3	GOL	B	504	-	-	0/4/4/4	-
3	GOL	A	504	-	-	0/4/4/4	-
3	GOL	B	508	-	-	4/4/4/4	-
3	GOL	B	509	-	-	3/4/4/4	-
3	GOL	C	505	-	-	0/4/4/4	-
3	GOL	C	508	-	-	2/4/4/4	-
3	GOL	A	506	-	-	2/4/4/4	-
3	GOL	A	503	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	505	GOL	C1-C2	2.49	1.62	1.51
3	B	504	GOL	C1-C2	2.22	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	504	GOL	C3-C2-C1	-2.07	103.65	111.70

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	507	GOL	O1-C1-C2-C3
3	B	508	GOL	C1-C2-C3-O3
3	C	504	GOL	C1-C2-C3-O3
3	C	506	GOL	O1-C1-C2-O2
3	C	506	GOL	O1-C1-C2-C3
3	C	506	GOL	O2-C2-C3-O3
3	C	507	GOL	O1-C1-C2-C3
3	A	506	GOL	C1-C2-C3-O3
3	A	507	GOL	C1-C2-C3-O3

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	507	GOL	C1-C2-C3-O3
3	B	508	GOL	O1-C1-C2-C3
3	B	509	GOL	O1-C1-C2-C3
3	B	509	GOL	C1-C2-C3-O3
3	C	506	GOL	C1-C2-C3-O3
3	C	508	GOL	C1-C2-C3-O3
3	B	507	GOL	O2-C2-C3-O3
3	C	504	GOL	O2-C2-C3-O3
3	A	506	GOL	O2-C2-C3-O3
3	A	507	GOL	O2-C2-C3-O3
3	B	509	GOL	O2-C2-C3-O3
3	C	507	GOL	O1-C1-C2-O2
3	C	508	GOL	O2-C2-C3-O3
3	B	508	GOL	O2-C2-C3-O3
3	B	508	GOL	O1-C1-C2-O2
3	B	505	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	509	GOL	3	0
3	C	507	GOL	1	0
3	C	506	GOL	2	0
3	C	504	GOL	1	0
3	A	506	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	439/454 (96%)	0.16	31 (7%) 16 18	19, 29, 52, 88	0
1	B	440/454 (96%)	-0.19	7 (1%) 72 77	15, 20, 38, 102	0
1	C	440/454 (96%)	-0.16	12 (2%) 54 61	16, 22, 42, 82	0
All	All	1319/1362 (96%)	-0.07	50 (3%) 40 46	15, 23, 45, 102	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	8.8
1	C	1	MET	8.4
1	A	1	MET	7.5
1	C	0	HIS	6.8
1	A	0	HIS	5.7
1	A	88	TYR	5.7
1	A	313	PRO	5.4
1	A	310	VAL	5.4
1	A	227	LEU	5.1
1	A	314	ASN	4.9
1	A	40	PRO	4.8
1	B	0	HIS	4.7
1	A	229	GLU	4.6
1	C	229	GLU	4.3
1	A	268	GLU	4.2
1	A	312	ALA	4.2
1	A	301	THR	4.1
1	B	88	TYR	3.7
1	A	231	SER	3.7
1	A	228	GLY	3.6
1	A	311	THR	3.5
1	A	300	ALA	3.4
1	A	38	LYS	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	39	ILE	3.3
1	A	41	GLY	3.2
1	A	298	TYR	3.2
1	B	229	GLU	3.0
1	C	339	GLU	2.7
1	C	228	GLY	2.7
1	A	230	SER	2.6
1	C	234	LYS	2.6
1	C	235	LEU	2.5
1	B	89	GLY	2.5
1	A	302	ALA	2.4
1	A	303	LYS	2.4
1	C	268	GLU	2.4
1	B	301	THR	2.4
1	A	235	LEU	2.4
1	C	259	GLU	2.3
1	C	338	LYS	2.3
1	A	177	VAL	2.3
1	A	90	ALA	2.3
1	C	419	GLN	2.2
1	C	231	SER	2.2
1	A	125	SER	2.2
1	A	338	LYS	2.1
1	A	128	ASP	2.1
1	A	234	LYS	2.1
1	A	315	VAL	2.0
1	B	40	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	C	505	6/6	0.23	0.36	88,93,110,110	0
3	GOL	B	506	6/6	0.34	0.31	60,72,83,86	0
3	GOL	C	508	6/6	0.45	0.30	46,67,81,81	0
3	GOL	C	507	6/6	0.49	0.20	73,88,88,88	0
3	GOL	A	503	6/6	0.57	0.15	45,54,64,65	0
3	GOL	A	505	6/6	0.57	0.32	59,72,84,84	0
3	GOL	B	507	6/6	0.58	0.21	73,78,91,94	0
3	GOL	B	508	6/6	0.59	0.28	49,73,90,90	0
3	GOL	B	505	6/6	0.59	0.23	38,55,67,71	0
3	GOL	A	507	6/6	0.64	0.21	55,65,76,78	0
3	GOL	A	506	6/6	0.67	0.16	72,81,96,96	0
2	SO4	C	502	5/5	0.69	0.26	89,94,96,102	0
2	SO4	B	502	5/5	0.70	0.22	89,93,98,98	0
2	SO4	A	502	5/5	0.75	0.23	89,89,93,95	0
3	GOL	C	504	6/6	0.80	0.19	32,38,46,47	0
3	GOL	A	504	6/6	0.82	0.18	47,55,66,66	0
3	GOL	C	506	6/6	0.82	0.21	55,62,69,75	0
3	GOL	B	509	6/6	0.85	0.29	29,71,86,87	0
2	SO4	C	501	5/5	0.88	0.20	73,76,81,81	0
2	SO4	A	501	5/5	0.89	0.19	77,79,81,86	0
2	SO4	B	503	5/5	0.92	0.20	65,66,68,69	0
3	GOL	B	510	6/6	0.94	0.09	22,27,31,31	0
3	GOL	B	504	6/6	0.96	0.10	21,28,34,34	0
2	SO4	C	503	5/5	0.98	0.09	39,40,41,45	0
2	SO4	B	501	5/5	0.99	0.08	24,28,31,32	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.