



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 12:34 am BST

PDB ID : 5O69
Title : The structure of the thermobifida fusca guanidine III riboswitch with agmatine.
Authors : Huang, L.; Wang, J.; Lilley, D.M.J.
Deposited on : 2017-06-06
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)
RNA backbone	3102	1031 (2.70-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	41	
2	B	41	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AG2	A	101	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2569 atoms, of which 860 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called RNA (41-MER).

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	Br	C	H	N	O	P			
1	A	41	1335	2	393	450	168	282	40	0	0	0

- Molecule 2 is a RNA chain called RNA (37-MER).

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	Br	C	H	N	O	P			
2	B	37	1192	2	355	396	153	251	35	0	0	0

- Molecule 3 is AGMATINE (three-letter code: AG2) (formula: C₅H₁₄N₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	N		
3	A	1	23	5	14	4	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	10	Total O 10 10	0	0
5	B	8	Total O 8 8	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (41-MER)

Chain A: 



- Molecule 2: RNA (37-MER)

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.89Å 83.89Å 67.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.32 – 2.32 49.34 – 2.32	Depositor EDS
% Data completeness (in resolution range)	98.9 (36.32-2.32) 99.2 (49.34-2.32)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 2.32Å)	Xtrriage
Refinement program	PHENIX (dev_2219: ???)	Depositor
R, R_{free}	0.243 , 0.270 0.251 , 0.276	Depositor DCC
R_{free} test set	618 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	48.3	Xtrriage
Anisotropy	0.303	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 32.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.048 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2569	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CBV, AG2, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/940	0.83	0/1462
2	B	0.30	0/838	0.85	0/1297
All	All	0.37	0/1778	0.84	0/2759

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	885	450	448	6	0
2	B	796	396	407	10	0
3	A	9	14	14	2	0
4	A	1	0	0	0	0
5	A	10	0	0	0	0
5	B	8	0	0	1	0
All	All	1709	860	869	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:G:O2'	1:A:30:A:N7	2.33	0.59
2:B:24:G:O3'	5:B:101:HOH:O	2.18	0.56
2:B:33:A:O2'	2:B:34:G:O5'	2.26	0.51
1:A:6:C:O2'	3:A:101:AG2:HD2	2.11	0.50
1:A:17:G:O6	3:A:101:AG2:NH1	2.38	0.50
1:A:2:CBV:O5'	1:A:2:CBV:H6	2.13	0.49
1:A:26:C:H3'	1:A:27:A:H5'	1.93	0.49
2:B:39:CBV:H3'	2:B:40:G:OP1	2.14	0.48
2:B:23:G:H1'	2:B:27:A:O2'	2.14	0.47
2:B:33:A:C2'	2:B:34:G:O5'	2.63	0.47
2:B:33:A:H2'	2:B:34:G:O4'	2.15	0.47
2:B:33:A:O2'	2:B:34:G:P	2.74	0.46
2:B:39:CBV:H2'	2:B:40:G:OP2	2.17	0.44
1:A:2:CBV:O2'	1:A:27:A:N1	2.45	0.43
2:B:5:A:C2	2:B:29:G:C2	3.07	0.42
2:B:2:CBV:O5'	2:B:2:CBV:H6	2.20	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	38/41 (92%)	7 (18%)	0
2	B	30/41 (73%)	9 (30%)	2 (6%)
All	All	68/82 (82%)	16 (23%)	2 (2%)

All (16) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	12	G
1	A	16	C
1	A	23	G
1	A	27	A
1	A	29	G
1	A	35	A
1	A	36	C
2	B	5	A
2	B	6	C
2	B	7	G
2	B	12	G
2	B	16	C
2	B	23	G
2	B	34	G
2	B	35	A
2	B	36	C

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	33	A
2	B	35	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CBV	A	39	1	23,23,23	2.96	9 (39%)	34,35,35	2.22	6 (17%)
1	CBV	A	2	1	23,23,23	3.04	9 (39%)	34,35,35	0.80	1 (2%)
2	CBV	B	2	2	23,23,23	3.07	9 (39%)	34,35,35	0.91	2 (5%)
2	CBV	B	39	2	23,23,23	3.01	9 (39%)	34,35,35	3.51	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CBV	A	39	1	-	2/10/26/26	0/2/2/2
1	CBV	A	2	1	-	3/10/26/26	0/2/2/2
2	CBV	B	2	2	-	4/10/26/26	0/2/2/2
2	CBV	B	39	2	-	3/10/26/26	0/2/2/2

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	CBV	C6-C5	7.86	1.49	1.34
2	B	2	CBV	C6-C5	7.83	1.49	1.34
1	A	39	CBV	C6-C5	7.54	1.49	1.34
2	B	39	CBV	C6-C5	7.46	1.49	1.34
2	B	39	CBV	C4-N3	6.47	1.45	1.34
1	A	39	CBV	C4-N3	6.29	1.44	1.34
2	B	2	CBV	C4-N3	6.21	1.44	1.34
1	A	2	CBV	C4-N3	6.09	1.44	1.34
2	B	2	CBV	C2-N3	6.06	1.48	1.36
2	B	39	CBV	C2-N3	6.00	1.48	1.36
1	A	2	CBV	C2-N3	5.92	1.48	1.36
1	A	39	CBV	C2-N3	5.40	1.47	1.36
1	A	39	CBV	C4-N4	4.57	1.46	1.34
2	B	2	CBV	C6-N1	4.54	1.45	1.38
1	A	2	CBV	C6-N1	4.45	1.45	1.38
2	B	39	CBV	C4-N4	4.37	1.45	1.34
1	A	2	CBV	C4-N4	4.36	1.45	1.34
2	B	2	CBV	C4-N4	4.33	1.45	1.34
2	B	2	CBV	C2-N1	4.11	1.48	1.40
2	B	39	CBV	C6-N1	4.07	1.45	1.38
1	A	39	CBV	C6-N1	4.04	1.44	1.38
2	B	39	CBV	C2-N1	4.02	1.48	1.40
1	A	2	CBV	C2-N1	3.74	1.48	1.40
1	A	39	CBV	C2-N1	3.60	1.47	1.40
1	A	39	CBV	C4-C5	3.03	1.49	1.42
1	A	2	CBV	O2-C2	-3.02	1.18	1.23
2	B	2	CBV	C4-C5	2.95	1.49	1.42
2	B	39	CBV	O2-C2	-2.84	1.18	1.23
2	B	39	CBV	C4-C5	2.81	1.49	1.42
1	A	2	CBV	C4-C5	2.73	1.49	1.42
2	B	2	CBV	O2-C2	-2.62	1.18	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	CBV	O2-C2	-2.58	1.18	1.23
2	B	2	CBV	BR-C5	2.51	1.94	1.88
1	A	39	CBV	BR-C5	2.39	1.94	1.88
1	A	2	CBV	BR-C5	2.35	1.94	1.88
2	B	39	CBV	BR-C5	2.01	1.93	1.88

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	39	CBV	O3P-P-O5'	-13.98	69.53	106.73
2	B	39	CBV	O1P-P-O5'	11.82	138.20	106.73
2	B	39	CBV	O1P-P-O3P	-8.14	76.53	107.64
1	A	39	CBV	O3P-P-O5'	-6.63	89.09	106.73
1	A	39	CBV	O3P-P-O2P	-6.60	84.84	110.68
1	A	39	CBV	O1P-P-O3P	-5.18	87.84	107.64
1	A	39	CBV	O1P-P-O2P	5.08	130.56	110.68
1	A	39	CBV	O5'-P-O2P	3.08	115.12	106.47
2	B	2	CBV	O3P-P-O2P	-2.67	100.22	110.68
1	A	39	CBV	O1P-P-O5'	2.57	113.57	106.73
2	B	2	CBV	O3P-P-O5'	2.51	113.42	106.73
1	A	2	CBV	O3P-P-O5'	2.13	112.39	106.73

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	2	CBV	C5'-O5'-P-O3P
1	A	2	CBV	C5'-O5'-P-O1P
1	A	39	CBV	C5'-O5'-P-O3P
1	A	39	CBV	C5'-O5'-P-O1P
2	B	2	CBV	C5'-O5'-P-O3P
2	B	2	CBV	C5'-O5'-P-O1P
2	B	2	CBV	C5'-O5'-P-O2P
2	B	39	CBV	C5'-O5'-P-O3P
2	B	39	CBV	O4'-C4'-C5'-O5'
2	B	39	CBV	C3'-C4'-C5'-O5'
1	A	2	CBV	C5'-O5'-P-O2P
2	B	2	CBV	O4'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	2	CBV	2	0
2	B	2	CBV	1	0
2	B	39	CBV	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AG2	A	101	-	8,8,8	2.69	3 (37%)	7,8,8	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AG2	A	101	-	-	2/6/6/6	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	101	AG2	CZ-NE	6.44	1.46	1.33
3	A	101	AG2	CZ-NH1	3.38	1.45	1.32
3	A	101	AG2	CZ-NH2	-2.15	1.26	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	101	AG2	N-CA-CB-CG
3	A	101	AG2	CG-CD-NE-CZ

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	101	AG2	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	39:CBV	O3'	40:G	P	4.16

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	39/41 (95%)	1.31	3 (7%) 13 18	43, 53, 66, 87	0
2	B	35/41 (85%)	1.68	12 (34%) 0 0	51, 94, 142, 153	0
All	All	74/82 (90%)	1.49	15 (20%) 1 1	43, 63, 124, 153	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	27	A	4.4
2	B	32	A	3.4
2	B	6	C	3.3
2	B	31	C	3.3
2	B	20	C	2.8
2	B	38	G	2.6
2	B	21	C	2.5
1	A	26	C	2.4
2	B	15	C	2.3
2	B	3	G	2.2
2	B	17	G	2.2
2	B	5	A	2.0
2	B	16	C	2.0
1	A	30	A	2.0
1	A	16	C	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CBV	B	2	22/22	0.84	0.17	88,108,140,141	0
1	CBV	A	2	22/22	0.87	0.19	47,80,107,125	0
2	CBV	B	39	22/22	0.89	0.28	43,81,165,198	0
1	CBV	A	39	22/22	0.93	0.19	29,46,74,80	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	AG2	A	101	9/9	0.53	0.50	43,154,209,223	0
4	MG	A	102	1/1	0.85	1.17	275,275,275,275	0

6.5 Other polymers [i](#)

There are no such residues in this entry.