



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 01:33 PM EDT

PDB ID : 5OMA  
Title : CH3 chimera of human 14-3-3 sigma with the StARD1 peptide including Ser57  
Authors : Sluchanko, N.N.; Tugaeva, K.V.; Greive, S.J.; Antson, A.A.  
Deposited on : 2017-07-28  
Resolution : 3.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1002 (4.14-3.66)
Clashscore	141614	1004 (4.12-3.68)
Ramachandran outliers	138981	1021 (4.14-3.66)
Sidechain outliers	138945	1014 (4.14-3.66)
RSRZ outliers	127900	1275 (4.20-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	 83% 10% • 6%
1	B	248	 82% 12% 5%
1	C	248	 17% 79% 12% • 8%
1	D	248	 2% 81% 9% • 9%
2	H	4	 100%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7285 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein sigma, Steroidogenic acute regulatory protein, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	232	1819	1131	308	369	1	10	0	0	0
1	B	236	1859	1155	320	373	1	10	0	0	0
1	C	227	1791	1119	303	360		9	0	0	0
1	D	225	1777	1111	301	356		9	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P31947
A	-1	PRO	-	expression tag	UNP P31947
A	0	HIS	-	expression tag	UNP P31947
A	75	ALA	GLU	engineered mutation	UNP P31947
A	76	ALA	GLU	engineered mutation	UNP P31947
A	77	ALA	LYS	engineered mutation	UNP P31947
A	232	GLY	-	linker	UNP P31947
A	233	SER	-	linker	UNP P31947
A	234	GLY	-	linker	UNP P31947
A	235	SEP	-	linker	UNP P31947
A	236	LEU	-	linker	UNP P31947
B	-2	GLY	-	expression tag	UNP P31947
B	-1	PRO	-	expression tag	UNP P31947
B	0	HIS	-	expression tag	UNP P31947
B	75	ALA	GLU	engineered mutation	UNP P31947
B	76	ALA	GLU	engineered mutation	UNP P31947
B	77	ALA	LYS	engineered mutation	UNP P31947
B	232	GLY	-	linker	UNP P31947
B	233	SER	-	linker	UNP P31947
B	234	GLY	-	linker	UNP P31947

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	235	SEP	-	linker	UNP P31947
B	236	LEU	-	linker	UNP P31947
C	-2	GLY	-	expression tag	UNP P31947
C	-1	PRO	-	expression tag	UNP P31947
C	0	HIS	-	expression tag	UNP P31947
C	75	ALA	GLU	engineered mutation	UNP P31947
C	76	ALA	GLU	engineered mutation	UNP P31947
C	77	ALA	LYS	engineered mutation	UNP P31947
C	232	GLY	-	linker	UNP P31947
C	233	SER	-	linker	UNP P31947
C	234	GLY	-	linker	UNP P31947
C	235	SEP	-	linker	UNP P31947
C	236	LEU	-	linker	UNP P31947
D	-2	GLY	-	expression tag	UNP P31947
D	-1	PRO	-	expression tag	UNP P31947
D	0	HIS	-	expression tag	UNP P31947
D	75	ALA	GLU	engineered mutation	UNP P31947
D	76	ALA	GLU	engineered mutation	UNP P31947
D	77	ALA	LYS	engineered mutation	UNP P31947
D	232	GLY	-	linker	UNP P31947
D	233	SER	-	linker	UNP P31947
D	234	GLY	-	linker	UNP P31947
D	235	SEP	-	linker	UNP P31947
D	236	LEU	-	linker	UNP P31947

- Molecule 2 is a protein called Undetermined peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	H	4	20	12	4	4	0	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	B	1	8	4	1	3	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	O			P
4	C	1	5	4	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O	0	0
			1	1		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3 protein sigma, Steroidogenic acute regulatory protein, mitochondrial

Chain A: 





- Molecule 1: 14-3-3 protein sigma, Steroidogenic acute regulatory protein, mitochondrial

Chain B: 




- Molecule 1: 14-3-3 protein sigma, Steroidogenic acute regulatory protein, mitochondrial


Chain C: 



- Molecule 1: 14-3-3 protein sigma, Steroidogenic acute regulatory protein, mitochondrial

Chain D: 





- Molecule 2: Undetermined peptide

Chain H:  100%

There are no outlier residues recorded for this chain.



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.25Å 123.25Å 162.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.00 – 3.90 49.09 – 3.79	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.00-3.90) 99.8 (49.09-3.79)	Depositor EDS
$R_{merge}$	0.27	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.11 (at 3.77Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.209 , 0.248 0.239 , 0.271	Depositor DCC
$R_{free}$ test set	1049 reflections (8.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	139.3	Xtrriage
Anisotropy	0.156	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 159.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.38$ , $\langle L^2 \rangle = 0.20$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7285	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	198.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, PO4, TRS, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/1834	0.65	0/2467
1	B	0.53	0/1874	0.69	0/2518
1	C	0.50	0/1816	0.66	0/2445
1	D	0.50	0/1802	0.67	0/2426
All	All	0.51	0/7326	0.66	0/9856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1819	0	1781	10	0
1	B	1859	0	1825	8	0
1	C	1791	0	1761	16	0
1	D	1777	0	1747	7	0
2	H	20	0	6	0	0
3	B	8	0	12	0	0
4	C	5	0	0	0	0
5	D	5	0	0	0	0
6	A	1	0	0	0	0
All	All	7285	0	7132	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:70:ASN:O	1:C:72:GLU:HA	1.51	1.10
1:C:70:ASN:O	1:C:72:GLU:OE2	1.84	0.95
1:C:70:ASN:O	1:C:72:GLU:CD	2.06	0.94
1:C:70:ASN:C	1:C:72:GLU:HA	1.90	0.91
1:C:71:GLU:HA	1:C:72:GLU:HB2	1.55	0.88
1:A:221:GLN:HA	1:A:224:ARG:HE	1.42	0.82
1:A:221:GLN:HA	1:A:224:ARG:NE	1.95	0.82
1:C:174:LEU:HD13	1:C:223:LEU:HG	1.62	0.80
1:C:70:ASN:O	1:C:72:GLU:CA	2.32	0.78
1:B:121:LEU:HB3	1:B:154:ALA:HB2	1.78	0.66
1:B:156:ASP:O	1:B:160:LYS:HD2	2.01	0.60
1:A:49:LYS:HG2	1:B:238:ARG:HH22	1.68	0.58
1:C:45:SER:O	1:C:49:LYS:HB2	2.07	0.55
1:C:70:ASN:O	1:C:72:GLU:CG	2.54	0.55
1:C:70:ASN:O	1:C:72:GLU:CB	2.55	0.55
1:D:155:MET:HG3	1:D:176:PHE:HD2	1.74	0.53
1:D:198:PHE:HB2	1:D:223:LEU:HD13	1.93	0.50
1:D:155:MET:HA	1:D:172:LEU:HD21	1.96	0.47
1:A:221:GLN:HB2	1:A:224:ARG:HH21	1.80	0.46
1:A:27:LYS:HG3	1:A:102:LEU:HD11	1.97	0.46
1:D:3:ARG:HE	1:D:34:GLU:HB2	1.81	0.46
1:B:176:PHE:O	1:B:179:PHE:HB3	2.17	0.45
1:C:158:SER:HB2	1:C:169:ARG:HG3	1.98	0.45
1:A:221:GLN:O	1:A:224:ARG:HG2	2.17	0.45
1:C:71:GLU:HA	1:C:72:GLU:CB	2.31	0.45
1:B:3:ARG:CZ	1:B:34:GLU:HB2	2.46	0.44
1:A:3:ARG:NE	1:A:34:GLU:HB2	2.32	0.44
1:A:128:TYR:HB3	1:A:147:ALA:HB2	1.99	0.44
1:D:27:LYS:HG3	1:D:102:LEU:HD11	1.99	0.44
1:C:3:ARG:NE	1:C:34:GLU:HB2	2.34	0.43
1:D:3:ARG:NE	1:D:34:GLU:HB2	2.33	0.43
1:A:3:ARG:HE	1:A:34:GLU:HB2	1.83	0.43
1:C:3:ARG:HE	1:C:34:GLU:HB2	1.83	0.42
1:C:59:TRP:CE2	1:C:134:VAL:HG12	2.55	0.42
1:C:106:HIS:HB3	1:C:109:LYS:HG3	2.02	0.42
1:A:205:LEU:HD13	1:A:213:TYR:HE1	1.85	0.41
1:B:166:ASN:HA	1:B:167:PRO:HD3	1.97	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:LYS:HG3	1:B:102:LEU:HD11	2.02	0.41
1:D:45:SER:O	1:D:49:LYS:HB2	2.20	0.40
1:B:3:ARG:NE	1:B:34:GLU:HB2	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/248 (92%)	219 (96%)	8 (4%)	1 (0%)	34	71
1	B	231/248 (93%)	217 (94%)	13 (6%)	1 (0%)	34	71
1	C	223/248 (90%)	212 (95%)	9 (4%)	2 (1%)	17	54
1	D	221/248 (89%)	215 (97%)	5 (2%)	1 (0%)	29	67
All	All	903/992 (91%)	863 (96%)	35 (4%)	5 (1%)	25	63

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	233	SER
1	C	77	ALA
1	B	2	GLU
1	C	137	GLY
1	D	107	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/204 (94%)	177 (92%)	15 (8%)	12	41
1	B	196/204 (96%)	178 (91%)	18 (9%)	9	34
1	C	190/204 (93%)	172 (90%)	18 (10%)	8	33
1	D	189/204 (93%)	174 (92%)	15 (8%)	12	41
All	All	767/816 (94%)	701 (91%)	66 (9%)	10	38

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	49	LYS
1	A	52	VAL
1	A	64	SER
1	A	123	MET
1	A	139	ASP
1	A	145	ASP
1	A	149	SER
1	A	160	LYS
1	A	185	ASN
1	A	192	SER
1	A	204	ASP
1	A	210	GLU
1	A	211	ASP
1	A	224	ARG
1	B	0	HIS
1	B	2	GLU
1	B	45	SER
1	B	49	LYS
1	B	52	VAL
1	B	64	SER
1	B	108	ILE
1	B	113	ASP
1	B	123	MET
1	B	138	ASP
1	B	145	ASP
1	B	185	ASN
1	B	192	SER
1	B	204	ASP
1	B	211	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	213	TYR
1	B	228	THR
1	B	236	LEU
1	C	45	SER
1	C	49	LYS
1	C	52	VAL
1	C	64	SER
1	C	108	ILE
1	C	121	LEU
1	C	123	MET
1	C	138	ASP
1	C	139	ASP
1	C	141	LYS
1	C	145	ASP
1	C	185	ASN
1	C	192	SER
1	C	204	ASP
1	C	211	ASP
1	C	213	TYR
1	C	222	LEU
1	C	223	LEU
1	D	49	LYS
1	D	52	VAL
1	D	64	SER
1	D	105	SER
1	D	108	ILE
1	D	123	MET
1	D	138	ASP
1	D	145	ASP
1	D	185	ASN
1	D	192	SER
1	D	204	ASP
1	D	211	ASP
1	D	213	TYR
1	D	222	LEU
1	D	223	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	8	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	SEP	B	235	1	8,9,10	1.18	1 (12%)	8,12,14	3.25	6 (75%)
1	SEP	A	235	1	8,9,10	1.23	1 (12%)	8,12,14	5.35	4 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	235	1	-	4/5/8/10	-
1	SEP	A	235	1	-	3/5/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	235	SEP	P-OG	-2.64	1.51	1.60
1	A	235	SEP	CB-CA	2.06	1.58	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	SEP	OG-CB-CA	12.86	120.66	108.14
1	A	235	SEP	O2P-P-OG	6.70	124.57	106.73
1	B	235	SEP	O3P-P-OG	-4.74	94.11	106.73
1	B	235	SEP	O2P-P-OG	4.48	118.66	106.73

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	235	SEP	OG-CB-CA	4.04	112.08	108.14
1	B	235	SEP	P-OG-CB	-3.00	110.04	118.30
1	A	235	SEP	O3P-P-OG	-2.95	98.90	106.73
1	B	235	SEP	OG-P-O1P	2.57	113.69	106.47
1	B	235	SEP	O3P-P-O2P	2.48	117.10	107.64
1	A	235	SEP	P-OG-CB	-2.44	111.57	118.30

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	235	SEP	CB-OG-P-O2P
1	A	235	SEP	CB-OG-P-O3P
1	B	235	SEP	N-CA-CB-OG
1	B	235	SEP	CB-OG-P-O1P
1	B	235	SEP	CB-OG-P-O2P
1	B	235	SEP	CB-OG-P-O3P
1	A	235	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	SO4	D	301	-	4,4,4	0.38	0	6,6,6	0.47	0



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TRS	B	301	-	7,7,7	0.27	0	9,9,9	0.30	0
4	PO4	C	301	-	4,4,4	2.49	2 (50%)	6,6,6	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	B	301	-	-	3/9/9/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	301	PO4	P-O1	4.11	1.60	1.50
4	C	301	PO4	P-O4	2.01	1.60	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	TRS	C1-C-C3-O3
3	B	301	TRS	C2-C-C3-O3
3	B	301	TRS	N-C-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	231/248 (93%)	-0.36	1 (0%) 92 87	107, 176, 235, 262	0
1	B	235/248 (94%)	-0.23	1 (0%) 92 87	113, 168, 234, 267	0
1	C	227/248 (91%)	0.62	43 (18%) 1 1	178, 266, 297, 298	0
1	D	225/248 (90%)	-0.20	5 (2%) 62 51	116, 188, 240, 279	0
2	H	0/4	-	-	-	-
All	All	918/996 (92%)	-0.05	50 (5%) 25 21	107, 191, 291, 298	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	220	MET	7.1
1	D	210	GLU	6.3
1	C	29	ALA	5.9
1	C	26	MET	5.7
1	C	110	GLU	5.5
1	C	223	LEU	5.4
1	C	197	THR	5.1
1	C	43	LEU	4.5
1	C	222	LEU	4.4
1	C	217	THR	4.4
1	C	199	ASP	4.3
1	C	224	ARG	4.1
1	D	213	TYR	4.1
1	D	211	ASP	4.0
1	C	216	SER	3.9
1	C	3	ARG	3.8
1	C	219	ILE	3.6
1	C	111	ALA	3.5
1	C	201	ALA	3.5
1	C	84	TYR	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	55	GLN	3.2
1	C	30	VAL	3.1
1	C	44	LEU	3.1
1	C	170	LEU	3.1
1	C	194	ALA	3.0
1	C	195	LYS	3.0
1	C	227	LEU	2.9
1	C	52	VAL	2.9
1	C	196	THR	2.9
1	C	198	PHE	2.9
1	C	221	GLN	2.8
1	B	213	TYR	2.8
1	C	28	GLY	2.5
1	C	178	VAL	2.5
1	C	173	ALA	2.5
1	A	111	ALA	2.4
1	C	56	ARG	2.4
1	D	59	TRP	2.4
1	C	48	TYR	2.3
1	C	91	GLU	2.3
1	C	127	TYR	2.2
1	C	7	ILE	2.2
1	C	34	GLU	2.2
1	C	35	GLU	2.2
1	C	40	GLU	2.2
1	C	92	LEU	2.2
1	D	209	SER	2.1
1	C	33	GLY	2.1
1	C	88	VAL	2.1
1	C	218	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	SEP	A	235	10/11	0.77	0.39	173,208,211,222	0
1	SEP	B	235	10/11	0.87	0.24	161,164,170,173	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	D	301	5/5	0.70	0.19	173,174,174,174	0
4	PO4	C	301	5/5	0.89	0.20	280,281,281,281	0
3	TRS	B	301	8/8	0.89	0.24	143,154,159,161	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.