



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 02:00 AM EDT

PDB ID : 5Q7J  
Title : PanDDA analysis group deposition – Crystal Structure of DCLRE1A after initial refinement with no ligand modelled (structure 192)  
Authors : Newman, J.A.; Aitkenhead, H.; Lee, S.Y.; Kupinska, K.; Burgess-Brown, N.; Tallon, R.; Krojer, T.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.; Bountra, C.; Gileadi, O.  
Deposited on : 2017-05-25  
Resolution : 1.54 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DCLRE1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2726	1765	450	491	20	0	4	0

- Molecule 2 is MALONATE ION (three-letter code: MLI) (formula:  $C_3H_2O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	7	3	4	0	0

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ni	0	0
			1	1		


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	310	Total 310	O 310	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DCLRE1A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.79Å 57.87Å 114.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.78 – 1.54 51.65 – 1.54	Depositor EDS
% Data completeness (in resolution range)	97.5 (51.78-1.54) 97.6 (51.65-1.54)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.54Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.290 , 0.339 0.300 , 0.351	Depositor DCC
$R_{free}$ test set	2442 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.2	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3044	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/2802	0.84	4/3809 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1032	ARG	NE-CZ-NH1	-5.89	117.35	120.30
1	A	1032	ARG	NE-CZ-NH2	5.40	123.00	120.30
1	A	960	ARG	NE-CZ-NH1	5.25	122.93	120.30
1	A	829	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2726	0	2676	26	0
2	A	7	0	2	0	0
3	A	1	0	0	0	0
4	A	310	0	0	15	1
All	All	3044	0	2678	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:909:GLN:NE2	4:A:1203:HOH:O	2.16	0.75
1:A:955:GLN:OE1	4:A:1201:HOH:O	2.15	0.65
1:A:871:HIS:CE1	4:A:1300:HOH:O	2.51	0.63
1:A:815:ASP:OD1	4:A:1202:HOH:O	2.16	0.62
1:A:906:LYS:NZ	4:A:1210:HOH:O	2.41	0.53
1:A:980:THR:HG23	4:A:1236:HOH:O	2.11	0.51
1:A:743:LYS:HG3	1:A:744:HIS:CE1	2.47	0.50
1:A:717:PHE:CE1	1:A:728:TYR:HB3	2.47	0.49
1:A:742:SER:O	1:A:765:VAL:HG22	2.12	0.49
1:A:784:VAL:HG21	4:A:1481:HOH:O	2.13	0.48
1:A:875:VAL:O	1:A:958:ALA:HA	2.16	0.46
1:A:860:ASN:O	1:A:861:THR:C	2.55	0.46
1:A:1032:ARG:NE	4:A:1219:HOH:O	2.49	0.45
1:A:973:ILE:N	4:A:1216:HOH:O	2.49	0.45
1:A:859:ILE:HD11	1:A:889:ALA:HB1	1.99	0.45
1:A:702:PRO:HG2	1:A:705:LYS:HG3	2.00	0.43
1:A:973:ILE:HG22	4:A:1216:HOH:O	2.17	0.43
1:A:973:ILE:HD13	1:A:990:PRO:HD2	1.99	0.43
1:A:972:ARG:HB3	4:A:1216:HOH:O	2.19	0.43
1:A:1032:ARG:CZ	4:A:1219:HOH:O	2.68	0.41
1:A:864:GLU:HB2	4:A:1367:HOH:O	2.19	0.41
1:A:983:ASN:ND2	4:A:1213:HOH:O	2.45	0.41
1:A:978:PRO:HD3	4:A:1440:HOH:O	2.20	0.41
1:A:760:LYS:O	1:A:764[A]:HIS:HA	2.21	0.41
1:A:856:ARG:HE	1:A:856:ARG:HA	1.85	0.41
1:A:718:GLN:HB3	1:A:739:ALA:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1391:HOH:O	4:A:1402:HOH:O[3_545]	2.15	0.05



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	343/343 (100%)	324 (94%)	18 (5%)	1 (0%)	41 19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	978	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	299/305 (98%)	292 (98%)	7 (2%)	50 20

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	816	PHE
1	A	830	GLN
1	A	848	PHE
1	A	879	TYR
1	A	893	VAL
1	A	928	LEU
1	A	957	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	851	GLN
1	A	869	ASN
1	A	909	GLN
1	A	955	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MLI	A	1101	3	6,6,6	1.30	0	7,7,7	1.42	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	A	1101	3	-	4/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	MLI	O7-C2-C1	2.55	122.69	114.54
2	A	1101	MLI	O7-C2-O6	-2.02	118.27	123.30

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	MLI	C3-C1-C2-O6
2	A	1101	MLI	C2-C1-C3-O9
2	A	1101	MLI	C3-C1-C2-O7
2	A	1101	MLI	C2-C1-C3-O8

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	341/343 (99%)	1.58	84 (24%) <b>0</b> <b>0</b>	16, 32, 95, 155	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	970	PHE	27.5
1	A	966	HIS	19.7
1	A	744	HIS	14.3
1	A	768	GLN	12.2
1	A	939	PHE	10.6
1	A	769	TYR	10.5
1	A	972	ARG	10.5
1	A	941	GLY	10.4
1	A	973	ILE	10.3
1	A	745	PHE	9.4
1	A	980	THR	9.3
1	A	977	ILE	9.0
1	A	984	ILE	8.6
1	A	942	LEU	8.6
1	A	765	VAL	8.2
1	A	937	ILE	8.0
1	A	967	SER	8.0
1	A	968	ASN	7.9
1	A	979	GLN	7.7
1	A	986	ILE	7.5
1	A	971	THR	7.2
1	A	950	GLY	6.9
1	A	943	GLN	6.6
1	A	747	PHE	6.3
1	A	743	LYS	6.2
1	A	940	LYS	6.0
1	A	938	ASN	5.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	987	TYR	5.8
1	A	746	THR	5.7
1	A	974	ALA	5.4
1	A	982	GLY	5.0
1	A	780	ILE	4.9
1	A	978	PRO	4.7
1	A	969	LYS	4.6
1	A	951	GLY	4.4
1	A	976	VAL	4.4
1	A	928	LEU	4.3
1	A	750	TYR	4.3
1	A	879	TYR	4.3
1	A	767	GLU	4.2
1	A	981	LYS	4.0
1	A	944	SER	4.0
1	A	770	ILE	4.0
1	A	946	LEU	3.9
1	A	766	GLN	3.9
1	A	781	VAL	3.7
1	A	742	SER	3.7
1	A	741	LEU	3.6
1	A	764[A]	HIS	3.6
1	A	988	GLY	3.6
1	A	985	SER	3.6
1	A	711	GLY	3.5
1	A	856	ARG	3.5
1	A	724	GLY	3.4
1	A	721	VAL	3.3
1	A	983	ASN	3.3
1	A	963	GLY	3.2
1	A	964	TRP	3.0
1	A	935	MET	2.9
1	A	748	PRO	2.9
1	A	947	LYS	2.9
1	A	936	GLN	2.8
1	A	722	VAL	2.8
1	A	861	THR	2.7
1	A	957	LEU	2.6
1	A	954	ASN	2.6
1	A	826	LEU	2.6
1	A	700	THR	2.6
1	A	956	ILE	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	932	LEU	2.5
1	A	726	THR	2.5
1	A	862	ALA	2.5
1	A	925	CYS	2.4
1	A	894	LEU	2.4
1	A	989	ILE	2.4
1	A	717	PHE	2.3
1	A	945	HIS	2.3
1	A	965	THR	2.3
1	A	871	HIS	2.2
1	A	848	PHE	2.2
1	A	828	ALA	2.2
1	A	749	VAL	2.2
1	A	771	HIS	2.1
1	A	868	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MLI	A	1101	7/7	0.92	0.13	27,34,35,36	0
3	NI	A	1102	1/1	0.99	0.02	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.