



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2022 – 04:01 PM EDT

PDB ID : 5QHJ
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of human FAM83B in complex with FMOPL000709a
Authors : Pinkas, D.M.; Bufton, J.C.; Fox, A.E.; Talon, R.; Krojer, T.; Douangamath, A.; Collins, P.; Zhang, R.; von Delft, F.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Bullock, A.N.
Deposited on : 2018-05-18
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)

ENTRY-COMPOSITION INFOmissingINFO

SEQUENCE-PLOTS INFOmissingINFO

2 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.63Å 74.27Å 63.49Å 90.00° 96.45° 90.00°	Depositor
Resolution (Å)	63.09 – 1.68	Depositor
% Data completeness (in resolution range)	99.2 (63.09-1.68)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.68Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.171 , 0.213	Depositor
Wilson B-factor (Å ²)	19.5	Xtrriage
Anisotropy	0.146	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3124	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

3 Model quality [i](#)

3.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOV, IOD, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.07	1/1418 (0.1%)	1.13	8/1908 (0.4%)
1	B	1.09	2/1422 (0.1%)	1.03	5/1911 (0.3%)
All	All	1.08	3/2840 (0.1%)	1.08	13/3819 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	172	ARG	CZ-NH1	6.16	1.41	1.33
1	B	230	LYS	CB-CG	-5.56	1.37	1.52
1	A	172	ARG	CZ-NH1	5.38	1.40	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	172	ARG	NE-CZ-NH2	-14.51	113.04	120.30
1	A	172	ARG	NE-CZ-NH1	11.21	125.91	120.30
1	B	140	ARG	NE-CZ-NH2	-8.75	115.93	120.30
1	A	140	ARG	NE-CZ-NH2	-8.13	116.23	120.30
1	B	172	ARG	NE-CZ-NH2	-7.33	116.63	120.30
1	A	147	ARG	NE-CZ-NH1	6.59	123.60	120.30
1	B	140	ARG	NE-CZ-NH1	6.29	123.44	120.30
1	A	155	ASP	CB-CG-OD1	5.89	123.60	118.30
1	A	147	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	B	155	ASP	CB-CG-OD1	5.68	123.42	118.30
1	A	172	ARG	CD-NE-CZ	5.42	131.18	123.60
1	B	172	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	140	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

3.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1389	0	1397	4	0
1	B	1395	0	1413	6	0
2	A	4	0	0	0	0
3	A	16	0	24	0	0
3	B	12	0	18	0	0
4	A	15	0	0	1	0
4	B	15	0	0	0	0
5	A	138	0	0	3	0
5	B	140	0	0	4	0
All	All	3124	0	2852	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:LYS:NZ	5:B:401:HOH:O	1.60	1.25
4:A:309:GOV:O	5:A:401:HOH:O	1.84	0.93
1:A:144:LYS:NZ	5:A:402:HOH:O	2.08	0.85
1:B:279:ARG:N	5:B:404:HOH:O	2.20	0.73
1:B:145:GLU:OE1	5:B:402:HOH:O	2.08	0.71
1:B:278:ALA:C	5:B:404:HOH:O	2.33	0.65
1:A:236[B]:CYS:SG	1:A:236[B]:CYS:O	2.70	0.50
1:A:168:GLU:HG2	5:A:401:HOH:O	2.15	0.46
1:A:272:GLU:OE1	1:B:129:ARG:NH1	2.48	0.42
1:B:126:HIS:HA	1:B:127:PRO:C	2.41	0.41

There are no symmetry-related clashes.

3.3 Torsion angles [i](#)

3.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	6/180 (3%)	6 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

3.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

3.3.3 RNA [i](#)

There are no RNA molecules in this entry.

3.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

3.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

3.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic and 2 could not be matched to an existing wwPDB Chemical Component Dictionary definition at this stage - leaving 7 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

3.7 Other polymers [i](#)

There are no such residues in this entry.

3.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

4 Fit of model and data

4.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

4.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

4.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

4.4 Ligands

EDS failed to run properly - this section is therefore empty.

4.5 Other polymers

EDS failed to run properly - this section is therefore empty.