



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 07:54 PM EDT

PDB ID : 5R1C  
Title : PanDDA analysis group deposition – Auto-refined data of Aar2/RNaseH for ground state model 27, DMSO-free  
Authors : Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.; Weiss, M.S.  
Deposited on : 2020-02-12  
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	 9% 78% 12% 8%
2	B	308	 11% 85% 12% 8%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4655 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	2002	1283	335	372	12	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2568	1645	420	483	20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	170	SER	LEU	conflict	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ASP	deletion	UNP P32357


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	64	Total O 64 64	0	0
3	B	21	Total O 21 21	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.43Å 82.56Å 92.84Å 90.00° 108.09° 90.00°	Depositor
Resolution (Å)	23.55 – 1.91 44.89 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.7 (23.55-1.91) 99.7 (44.89-1.91)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.232 , 0.281 0.244 , 0.298	Depositor DCC
$R_{free}$ test set	2101 reflections (4.21%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.6	Xtrriage
Anisotropy	0.327	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 52.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4655	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/2049	0.97	8/2775 (0.3%)
2	B	0.72	2/2638 (0.1%)	0.82	2/3563 (0.1%)
All	All	0.75	2/4687 (0.0%)	0.89	10/6338 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	121	VAL	CB-CG2	-6.48	1.39	1.52
2	B	58	TYR	CD2-CE2	5.11	1.47	1.39

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1908	LEU	CB-CG-CD1	-8.21	97.03	111.00
1	A	1905	LEU	CB-CG-CD1	6.37	121.83	111.00
2	B	18	ASP	CB-CG-OD2	6.25	123.92	118.30
1	A	1942	ASP	CB-CG-OD2	6.00	123.70	118.30
2	B	265	ASP	CB-CG-OD2	5.89	123.60	118.30
1	A	1854	ASP	CB-CG-OD2	5.58	123.32	118.30
1	A	1957	ARG	NE-CZ-NH1	5.45	123.03	120.30
1	A	2012	LEU	CB-CG-CD2	5.38	120.14	111.00
1	A	1908	LEU	CB-CG-CD2	5.36	120.11	111.00
1	A	2011	LEU	CB-CG-CD1	-5.02	102.47	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	2029	20	0
2	B	2568	0	2441	22	0
3	A	64	0	0	2	0
3	B	21	0	0	2	0
All	All	4655	0	4470	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1:MET:N	3:B:401:HOH:O	2.14	0.80
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.85	0.77
2:B:70:GLN:HB3	2:B:81:MET:HE2	1.67	0.76
2:B:70:GLN:HB3	2:B:81:MET:CE	2.29	0.63
2:B:53:SER:HA	3:B:405:HOH:O	1.99	0.61
1:A:1980:LYS:NZ	3:A:2102:HOH:O	2.33	0.58
1:A:1962:ARG:O	1:A:2013:ARG:NH1	2.31	0.58
2:B:251:CYS:O	2:B:296:SER:HB2	2.07	0.54
1:A:2050:THR:HG22	1:A:2054:GLN:HE21	1.74	0.51
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.92	0.51
2:B:6:PHE:HZ	2:B:44[A]:ILE:HD11	1.77	0.50
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ1	1.59	0.50
1:A:1942:ASP:HB2	1:A:1943:PRO:HD3	1.95	0.49
1:A:1976:ASP:HB3	3:A:2102:HOH:O	2.13	0.49
2:B:237:TYR:CE2	2:B:241:LEU:HD11	2.49	0.48
1:A:1913:THR:HB	1:A:1940:MET:HE1	1.94	0.47
1:A:1910:LYS:HG2	1:A:1940:MET:SD	2.55	0.46
1:A:2056[B]:ARG:O	1:A:2060:LEU:HG	2.15	0.46
2:B:70:GLN:CB	2:B:81:MET:CE	2.94	0.46
2:B:214:PHE:O	2:B:215:LYS:HB2	2.15	0.46
2:B:41[B]:VAL:HG12	2:B:123:MET:HG2	1.98	0.45
2:B:188:ALA:HA	2:B:204:TYR:CD1	2.51	0.45
2:B:190:ARG:HG3	2:B:203[B]:TYR:CE2	2.51	0.45
1:A:1987:VAL:HG12	1:A:1989:PHE:CE1	2.51	0.45
2:B:17:ILE:O	2:B:17:ILE:HG23	2.17	0.45
1:A:1859:ARG:NH1	1:A:1979[A]:MET:SD	2.87	0.44
1:A:1914:ALA:O	1:A:1918[A]:SER:OG	2.33	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2062:GLU:HB3	1:A:2066:LYS:HE3	2.00	0.44
2:B:17:ILE:HD12	2:B:44[B]:ILE:CG1	2.48	0.44
1:A:1940:MET:HE2	1:A:1940:MET:HB3	1.76	0.43
1:A:1941:LEU:O	1:A:1945:GLU:HB2	2.19	0.43
1:A:2050:THR:HG22	1:A:2054:GLN:NE2	2.34	0.42
2:B:58:TYR:CE1	2:B:139:TYR:HD2	2.36	0.42
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.54	0.42
2:B:229:LEU:HD23	2:B:232:MET:HE2	2.02	0.42
2:B:281:ASP:OD1	2:B:281:ASP:N	2.49	0.42
2:B:193:HIS:ND1	2:B:196:GLU:OE1	2.45	0.42
1:A:2029:ASP:OD1	1:A:2031:THR:HG23	2.20	0.41
2:B:1:MET:HB3	2:B:35:ASP:HA	2.01	0.41
2:B:69:ILE:HD13	2:B:80:MET:HA	2.02	0.41
2:B:6:PHE:CZ	2:B:44[A]:ILE:HD11	2.55	0.41
1:A:1848:ILE:H	1:A:1931[A]:LYS:NZ	2.19	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/258 (96%)	239 (96%)	9 (4%)	0	100	100
2	B	305/308 (99%)	294 (96%)	10 (3%)	1 (0%)	41	31
All	All	553/566 (98%)	533 (96%)	19 (3%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	106	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/233 (97%)	219 (97%)	7 (3%)	40	30
2	B	286/284 (101%)	282 (99%)	4 (1%)	67	63
All	All	512/517 (99%)	501 (98%)	11 (2%)	62	46

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1918[A]	SER
1	A	1918[B]	SER
1	A	1962	ARG
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
2	B	17	ILE
2	B	77	LEU
2	B	301	SER
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1907	GLN
1	A	2038	HIS
1	A	2054	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	237/258 (91%)	0.80	23 (9%) <b>7</b> <b>9</b>	39, 63, 101, 123	0
2	B	300/308 (97%)	0.95	34 (11%) <b>5</b> <b>6</b>	45, 72, 121, 175	0
All	All	537/566 (94%)	0.88	57 (10%) <b>6</b> <b>7</b>	39, 69, 111, 175	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	53	SER	10.5
2	B	52	SER	7.3
2	B	1	MET	6.8
2	B	110	GLU	5.6
2	B	174	HIS	5.3
1	A	2060	LEU	4.7
2	B	175	SER	4.5
2	B	54[A]	MET	4.0
2	B	109	ASP	3.9
2	B	101	MET	3.8
2	B	277	GLU	3.6
1	A	1833	GLY	3.5
1	A	1838	SER	3.5
2	B	316	LEU	3.5
1	A	2027	LEU	3.5
2	B	279	TYR	3.3
2	B	281	ASP	3.1
1	A	1940	MET	3.1
2	B	92	ILE	3.1
2	B	96	PHE	3.1
1	A	2063	TYR	3.0
2	B	181	ILE	2.9
1	A	1979[A]	MET	2.9
2	B	86	GLY	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	108	ILE	2.7
2	B	189	ILE	2.7
2	B	89	PHE	2.7
2	B	193	HIS	2.6
1	A	1988	LEU	2.6
1	A	1969	MET	2.5
2	B	173	ALA	2.5
2	B	171	ASP	2.5
2	B	280	SER	2.4
2	B	191	PRO	2.4
1	A	2034	ILE	2.4
1	A	1889	LEU	2.3
1	A	1866	PHE	2.3
2	B	100	GLN	2.1
1	A	1861	THR	2.1
2	B	150	ASN	2.1
1	A	1850	LEU	2.1
1	A	1891	LEU	2.1
2	B	47	GLN	2.1
1	A	1865	THR	2.1
2	B	77	LEU	2.1
2	B	237	TYR	2.1
2	B	146	THR	2.1
2	B	224	LEU	2.1
1	A	1839	ASN	2.1
2	B	29	PRO	2.1
2	B	73	PRO	2.1
1	A	1954	ILE	2.1
1	A	2057[A]	ASP	2.0
1	A	1878	CYS	2.0
1	A	1840	TYR	2.0
1	A	2015	LEU	2.0
1	A	2017[A]	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.