

Full wwPDB X-ray Structure Validation Report (i)

Jun 17, 2024 – 08:39 PM EDT

PDB ID	:	5T58
Title	:	Structure of the MIND Complex Shows a Regulatory Focus of Yeast Kineto-
		chore Assembly
Authors	:	Dimitrova, Y.; Jenni, S.; Valverde, R.; Khin, Y.; Harrison, S.C.
Deposited on	:	2016-08-30
Resolution	:	3.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	1335 (3.24-3.20)
Clashscore	141614	1460 (3.24-3.20)
Ramachandran outliers	138981	1437 (3.24-3.20)
Sidechain outliers	138945	1436 (3.24-3.20)
RSRZ outliers	127900	1291 (3.24-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	233	<mark>6%</mark> 76%		21% •
2	В	205	9% 71%	21%	• 7%
3	D	250	.% 71%	9% •	20%
4	Ν	216	6%	24%	• 9%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 13336 atoms, of which 6743 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called KLLA0F02343p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	А	233	Total 3885	C 1210	Н 1969	N 329	O 370	S 7	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1	ALA	-	expression tag	UNP Q6CLK3

• Molecule 2 is a protein called KLLA0E05809p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	В	190	Total 3104	C 961	H 1551	N 274	0 312	${ m S}{ m 6}$	0	0	0

• Molecule 3 is a protein called KLLA0D15741p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	D	201	Total 3132	C 955	Н 1610	N 278	0 284	${f S}{5}$	0	0	0

• Molecule 4 is a protein called KLLA0C15939p.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
4	Ν	196	Total 3215	C 1004	Н 1613	N 267	O 327	${S \over 4}$	0	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: KLLA0F02343p







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	58.75Å 172.57Å 59.48Å	Demeriter
a, b, c, α , β , γ	90.00° 110.46° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	86.28 - 3.21	Depositor
Resolution (A)	86.28 - 3.21	EDS
% Data completeness	74.5 (86.28-3.21)	Depositor
(in resolution range)	$74.6\ (86.28-3.21)$	EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.14 (at 3.19 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
P. P.	0.320 , 0.362	Depositor
n, n_{free}	0.325 , 0.344	DCC
R_{free} test set	678 reflections $(5.03%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	80.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36,47.3	EDS
L-test for twinning ²	$< L > = 0.47, < L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.046 for l,-k,h	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	13336	wwPDB-VP
Average B, all atoms $(Å^2)$	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.26	0/1940	0.43	0/2621	
2	В	0.24	0/1568	0.41	0/2104	
3	D	0.24	0/1351	0.42	0/1803	
4	N	0.27	0/1619	0.47	0/2178	
All	All	0.25	0/6478	0.43	0/8706	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	174	ASN	Sidechain
1	А	209	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1916	1969	1972	38	0
2	В	1553	1551	1551	32	2
3	D	1522	1610	1475	18	0
4	N	1602	1613	1613	43	2
All	All	6593	6743	6611	110	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:103:ARG:NH2	2:B:103:ILE:O	2.04	0.90
3:D:359:LYS:NZ	4:N:88:TYR:O	2.09	0.83
1:A:91:ILE:O	2:B:16:ARG:NH1	2.17	0.77
1:A:2:THR:O	1:A:3:THR:OG1	2.05	0.72
1:A:25:ASP:OD2	4:N:104:TYR:OH	2.07	0.71
1:A:114:LEU:O	1:A:116:ASP:N	2.27	0.67
1:A:14:GLU:O	4:N:119:ARG:NH2	2.29	0.65
1:A:229:THR:O	1:A:230:SER:OG	2.13	0.64
2:B:143:LEU:O	2:B:146:ILE:N	2.30	0.64
1:A:141:ASN:ND2	4:N:147:MET:SD	2.70	0.64
4:N:138:GLU:O	4:N:142:SER:OG	2.13	0.61
1:A:119:THR:O	1:A:123:THR:OG1	2.13	0.61
2:B:113:ARG:O	2:B:119:ARG:NH2	2.34	0.60
1:A:50:ASN:ND2	1:A:58:SER:OG	2.32	0.59
2:B:143:LEU:O	2:B:146:ILE:HG22	2.03	0.59
1:A:103:ARG:NH2	2:B:100:ALA:O	2.36	0.59
1:A:173:ASP:OD2	1:A:177:THR:N	2.35	0.58
2:B:176:ASP:OD1	3:D:434:ARG:NH1	2.34	0.58
4:N:91:PRO:O	4:N:93:ASP:N	2.37	0.57
1:A:176:THR:OG1	4:N:173:ASN:OD1	2.23	0.57
1:A:9:THR:O	1:A:13:THR:OG1	2.19	0.56
2:B:71:ASN:OD1	2:B:74:ARG:NH2	2.39	0.56
2:B:62:GLN:O	2:B:66:ILE:HD12	2.09	0.52
4:N:121:ALA:HB3	4:N:122:PRO:HD3	1.92	0.52
4:N:29:LYS:O	4:N:32:THR:HG22	2.08	0.52
3:D:468:VAL:N	3:D:469:LYS:HA	2.25	0.52
2:B:116:ASP:OD2	4:N:118:ARG:NH1	2.42	0.52
3:D:467:ILE:O	3:D:468:VAL:HB	2.10	0.51
2:B:12:THR:O	2:B:14:HIS:N	2.38	0.51
1:A:38:CYS:SG	2:B:65:VAL:HG23	2.51	0.50



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
4:N:129:TYR:O	4:N:131:LYS:N	2.44	0.49
2:B:72:LEU:HA	2:B:75:VAL:HG22	1.94	0.49
4:N:143:LEU:O	4:N:146:LYS:N	2.43	0.48
4:N:193:ILE:HG22	4:N:194:PRO:HD3	1.96	0.48
4:N:40:ASP:HB3	4:N:43:VAL:HG13	1.95	0.48
2:B:62:GLN:O	2:B:65:VAL:HG12	2.14	0.47
4:N:40:ASP:O	4:N:41:ASP:HB2	2.14	0.47
1:A:108:HIS:CE1	4:N:119:ARG:HB3	2.50	0.47
1:A:113:VAL:HG12	2:B:126:GLY:HA3	1.95	0.47
3:D:467:ILE:C	3:D:469:LYS:HA	2.35	0.47
1:A:173:ASP:OD2	1:A:177:THR:HB	2.15	0.47
1:A:119:THR:O	1:A:123:THR:CB	2.64	0.46
3:D:382:ILE:HG21	4:N:110:GLU:HB3	1.97	0.46
4:N:214:VAL:O	4:N:214:VAL:HG13	2.16	0.46
4:N:116:LYS:O	4:N:120:GLU:HB3	2.16	0.46
4:N:8:LEU:HB2	4:N:66:MET:HG3	1.97	0.46
2:B:63:GLN:O	2:B:67:LYS:HG2	2.16	0.45
2:B:12:THR:O	2:B:12:THR:HG22	2.17	0.44
4:N:137:LEU:O	4:N:141:ASP:HB2	2.17	0.44
2:B:59:THR:O	2:B:62:GLN:HG2	2.17	0.44
1:A:113:VAL:HG12	2:B:126:GLY:CA	2.48	0.44
3:D:467:ILE:O	3:D:468:VAL:CB	2.66	0.44
4:N:119:ARG:HG3	4:N:120:GLU:N	2.33	0.44
4:N:119:ARG:O	4:N:123:GLN:OE1	2.36	0.44
1:A:97:ASP:O	1:A:101:GLU:HG2	2.17	0.44
2:B:14:HIS:O	2:B:15:ILE:HB	2.17	0.44
4:N:10:ILE:HG22	4:N:11:THR:N	2.32	0.44
4:N:75:THR:O	4:N:76:THR:HB	2.17	0.44
3:D:362:ASN:HB3	4:N:90:GLU:HA	1.99	0.43
4:N:10:ILE:HD11	4:N:66:MET:SD	2.57	0.43
3:D:266:TRP:CZ2	3:D:270:ARG:HD2	2.54	0.43
1:A:79:ASN:ND2	2:B:31:SER:OG	2.49	0.43
4:N:120:GLU:O	4:N:124:VAL:HG22	2.19	0.43
1:A:187:LYS:N	1:A:188:PRO:HD2	2.33	0.43
3:D:352:UNK:N	3:D:353:PRO:HD3	2.33	0.43
2:B:62:GLN:HA	2:B:65:VAL:HG12	2.00	0.43
2:B:108:SER:HB2	2:B:111:LYS:HD3	2.00	0.43
3:D:366:ILE:HD11	4:N:45:GLN:HE21	1.83	0.43
4:N:193:ILE:N	4:N:194:PRO:CD	2.82	0.43
4:N:89:THR:C	4:N:91:PRO:HD3	2.39	0.43
1:A:183:TRP:CD1	4:N:183:LEU:HB3	2.54	0.43

Continued from previous page...



		Interatomic	Clash	
Atom-1	Atom-2	distance $(Å)$	overlap (Å)	
4:N:138:GLU:O	4:N:142:SER:CB	2.67	0.43	
2:B:121:LEU:O	2:B:124:ILE:N	2.52	0.42	
1:A:193:VAL:O	1:A:197:THR:OG1	2.29	0.42	
1:A:101:GLU:O	1:A:102:ASN:HB2	2.19	0.42	
2:B:32:ILE:O	2:B:34:LYS:N	2.52	0.42	
4:N:89:THR:HG22	4:N:90:GLU:H	1.85	0.42	
4:N:90:GLU:N	4:N:91:PRO:HD3	2.34	0.42	
2:B:65:VAL:O	2:B:69:TRP:CB	2.67	0.42	
2:B:119:ARG:N	2:B:120:PRO:CD	2.83	0.42	
3:D:425:GLU:O	3:D:429:SER:OG	2.29	0.42	
1:A:35:MET:O	1:A:39:THR:OG1	2.38	0.42	
2:B:187:ASP:O	2:B:191:LYS:HG3	2.20	0.42	
3:D:455:THR:O	3:D:459:LYS:HG2	2.20	0.42	
1:A:34:ILE:HG22	2:B:69:TRP:CE3	2.55	0.42	
3:D:363:GLU:HA	3:D:366:ILE:HD13	2.02	0.41	
1:A:231:ILE:HG22	1:A:232:VAL:N	2.35	0.41	
3:D:293:ILE:O	3:D:297:PHE:HB2	2.18	0.41	
2:B:172:GLN:HA	2:B:175:TYR:CE2	2.55	0.41	
4:N:170:GLU:O	4:N:174:GLN:HG2	2.21	0.41	
1:A:125:THR:O	1:A:129:GLN:OE1	2.39	0.41	
4:N:89:THR:HG22	4:N:90:GLU:N	2.35	0.41	
3:D:259:LYS:HE2	4:N:8:LEU:HD21	2.02	0.41	
1:A:140:LEU:HD23	1:A:144:LEU:HD13	2.03	0.41	
3:D:363:GLU:HA	3:D:366:ILE:CD1	2.51	0.41	
4:N:120:GLU:O	4:N:124:VAL:HG13	2.20	0.41	
4:N:15:LEU:HD12	4:N:16:ARG:N	2.36	0.41	
4:N:30:VAL:O	4:N:34:LEU:HG	2.21	0.41	
1:A:168:LEU:HA	1:A:173:ASP:OD1	2.21	0.40	
1:A:230:SER:HA	1:A:231:ILE:HA	1.89	0.40	
1:A:213:ILE:HG23	1:A:214:ASP:N	2.36	0.40	
1:A:198:THR:O	1:A:201:LYS:HB2	2.21	0.40	
1:A:209:GLU:O	1:A:210:PHE:O	2.39	0.40	
2:B:143:LEU:O	2:B:144:GLN:C	2.60	0.40	
4:N:136:LEU:O	4:N:140:ILE:HG12	2.21	0.40	
1:A:120:ARG:HG3	1:A:121:ALA:N	2.36	0.40	
3:D:257:ASP:OD1	3:D:258:LEU:N	2.54	0.40	
1:A:159:VAL:HG13	1:A:160:GLN:N	2.37	0.40	
2:B:129:GLN:O	2:B:133:GLU:HG3	2.22	0.40	
4:N:123:GLN:OE1	4:N:123:GLN:N	2.55	0.40	

Continued from previous page...

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:107:SER:OG	4:N:95:GLU:OE2[1_455]	1.86	0.34
2:B:107:SER:HG	4:N:95:GLU:OE2[1_455]	1.58	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Per	centil	es
1	А	231/233 (99%)	185 (80%)	37~(16%)	9 (4%)	3	20	
2	В	$186/205 \ (91\%)$	156 (84%)	23~(12%)	7 (4%)	3	21	
3	D	160/250~(64%)	137 (86%)	19~(12%)	4 (2%)	5	31	
4	N	192/216~(89%)	160 (83%)	24 (12%)	8 (4%)	3	19	
All	All	769/904~(85%)	638 (83%)	103 (13%)	28 (4%)	3	22	

All (28) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	115	THR
1	А	210	PHE
2	В	14	HIS
2	В	15	ILE
2	В	117	ASP
3	D	468	VAL
4	Ν	92	PHE
1	А	114	LEU
2	В	115	ILE
4	Ν	41	ASP
4	N	130	THR
2	В	105	THR
3	D	354	PRO
4	N	93	ASP
1	А	2	THR
1	А	111	LYS



Mol	Chain	Res	Type
1	А	228	ASN
1	А	230	SER
2	В	109	SER
4	Ν	133	GLU
1	А	125	THR
4	Ν	39	GLU
4	Ν	200	GLN
1	А	3	THR
2	В	33	ARG
3	D	353	PRO
3	D	361	PRO
4	N	194	PRO

Continued from previous page...

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	222/222~(100%)	222 (100%)	0	100	100
2	В	178/193~(92%)	178 (100%)	0	100	100
3	D	154/198~(78%)	154 (100%)	0	100	100
4	Ν	184/203~(91%)	184 (100%)	0	100	100
All	All	738/816~(90%)	738 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	233/233~(100%)	0.52	13 (5%) 24 14	33, 83, 118, 138	0
2	В	190/205~(92%)	0.63	19 (10%) 7 4	38, 80, 118, 144	0
3	D	164/250~(65%)	0.27	2 (1%) 79 68	35, 74, 103, 116	0
4	Ν	196/216~(90%)	0.43	13 (6%) 18 11	35, 78, 115, 134	0
All	All	783/904~(86%)	0.47	47 (6%) 21 13	33, 79, 115, 144	0

All (47) RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
4	Ν	37	SER	6.7
1	А	57	PHE	6.4
1	А	45	TYR	4.3
2	В	53	GLU	4.0
1	А	53	GLY	3.7
2	В	56	GLU	3.7
4	N	73	LYS	3.6
2	В	60	VAL	3.4
4	N	197	GLU	3.0
1	А	50	ASN	2.9
1	А	142	VAL	2.9
1	А	1	ALA	2.9
2	В	157	LEU	2.8
4	N	150	SER	2.8
4	N	36	GLU	2.8
2	В	37	SER	2.7
1	А	36	TYR	2.7
1	А	54	LYS	2.7
4	N	211	GLU	2.6
2	В	109	SER	2.6
2	В	105	THR	2.6



Mol	Chain	\mathbf{Res}	Type	RSRZ
2	В	115	ILE	2.5
1	А	157	GLU	2.4
2	В	165	PHE	2.4
2	В	54	GLY	2.4
2	В	97	ILE	2.4
2	В	57	ASN	2.3
2	В	129	GLN	2.3
2	В	186	PRO	2.3
2	В	110	LYS	2.3
4	Ν	20	LYS	2.3
2	В	68	LEU	2.3
1	А	102	ASN	2.3
1	А	156	THR	2.3
4	Ν	187	LYS	2.2
4	Ν	209	LEU	2.2
1	А	183	TRP	2.2
2	В	108	SER	2.2
3	D	429	SER	2.2
4	Ν	145	ALA	2.2
1	А	103	ARG	2.1
4	N	207	LEU	2.1
2	В	52	ARG	2.1
4	Ν	86	LYS	2.0
2	В	153	LEU	2.0
4	N	172	TRP	2.0
3	D	403	LEU	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.



6.5 Other polymers (i)

There are no such residues in this entry.

