



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 10:19 PM EDT

PDB ID : 5YAB
Title : Crystal structure of scyllo-inositol dehydrogenase with L-glucose dehydrogenase activity
Authors : Fukano, K.; Shimizu, T.; Sasaki, Y.; Nakamura, A.; Yajima, S.
Deposited on : 2017-08-31
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	 3% 88% 7% ..
1	B	380	 8% 87% 9% ..
1	C	380	 4% 90% 6% ..
1	D	380	 5% 88% 8% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Scyllo-inositol dehydrogenase with L-glucose dehydrogenase activity.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	367	2800	1753	513	515	19	0	1	0
1	B	366	2785	1744	510	513	18	0	0	0
1	C	368	2802	1756	513	515	18	0	0	0
1	D	368	2808	1759	514	516	19	0	1	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	72	SER	ASN	engineered mutation	UNP K7ZP76
A	373	LEU	-	expression tag	UNP K7ZP76
A	374	GLU	-	expression tag	UNP K7ZP76
A	375	HIS	-	expression tag	UNP K7ZP76
A	376	HIS	-	expression tag	UNP K7ZP76
A	377	HIS	-	expression tag	UNP K7ZP76
A	378	HIS	-	expression tag	UNP K7ZP76
A	379	HIS	-	expression tag	UNP K7ZP76
A	380	HIS	-	expression tag	UNP K7ZP76
B	72	SER	ASN	engineered mutation	UNP K7ZP76
B	373	LEU	-	expression tag	UNP K7ZP76
B	374	GLU	-	expression tag	UNP K7ZP76
B	375	HIS	-	expression tag	UNP K7ZP76
B	376	HIS	-	expression tag	UNP K7ZP76
B	377	HIS	-	expression tag	UNP K7ZP76
B	378	HIS	-	expression tag	UNP K7ZP76
B	379	HIS	-	expression tag	UNP K7ZP76
B	380	HIS	-	expression tag	UNP K7ZP76
C	72	SER	ASN	engineered mutation	UNP K7ZP76
C	373	LEU	-	expression tag	UNP K7ZP76

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	374	GLU	-	expression tag	UNP K7ZP76
C	375	HIS	-	expression tag	UNP K7ZP76
C	376	HIS	-	expression tag	UNP K7ZP76
C	377	HIS	-	expression tag	UNP K7ZP76
C	378	HIS	-	expression tag	UNP K7ZP76
C	379	HIS	-	expression tag	UNP K7ZP76
C	380	HIS	-	expression tag	UNP K7ZP76
D	72	SER	ASN	engineered mutation	UNP K7ZP76
D	373	LEU	-	expression tag	UNP K7ZP76
D	374	GLU	-	expression tag	UNP K7ZP76
D	375	HIS	-	expression tag	UNP K7ZP76
D	376	HIS	-	expression tag	UNP K7ZP76
D	377	HIS	-	expression tag	UNP K7ZP76
D	378	HIS	-	expression tag	UNP K7ZP76
D	379	HIS	-	expression tag	UNP K7ZP76
D	380	HIS	-	expression tag	UNP K7ZP76

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	D	1	4	2	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	233	Total 233	O 233	0	0
3	B	177	Total 177	O 177	0	0
3	C	227	Total 227	O 227	0	0
3	D	215	Total 215	O 215	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Scyllo-inositol dehydrogenase with L-glucose dehydrogenase activity




- Molecule 1: Scyllo-inositol dehydrogenase with L-glucose dehydrogenase activity




- Molecule 1: Scyllo-inositol dehydrogenase with L-glucose dehydrogenase activity



- Molecule 1: Scyllo-inositol dehydrogenase with L-glucose dehydrogenase activity





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.97Å 127.45Å 137.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.30 – 1.75 28.30 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.3 (28.30-1.75) 99.3 (28.30-1.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.10 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.177 , 0.192 0.186 , 0.201	Depositor DCC
R_{free} test set	7983 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	12.3	Xtrriage
Anisotropy	0.239	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12063	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2866	0.74	3/3884 (0.1%)
1	B	0.47	0/2851	0.74	4/3865 (0.1%)
1	C	0.49	0/2868	0.73	2/3887 (0.1%)
1	D	0.49	0/2874	0.78	3/3895 (0.1%)
All	All	0.49	0/11459	0.75	12/15531 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	ARG	NE-CZ-NH2	10.76	125.68	120.30
1	B	334	ARG	NE-CZ-NH1	9.62	125.11	120.30
1	B	334	ARG	NE-CZ-NH2	-9.18	115.71	120.30
1	B	160	ARG	NE-CZ-NH1	7.59	124.09	120.30
1	A	229	ARG	NE-CZ-NH1	6.77	123.69	120.30
1	D	160	ARG	NE-CZ-NH1	6.72	123.66	120.30
1	C	160	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	160	ARG	NE-CZ-NH1	-5.54	117.53	120.30
1	D	27	ARG	NE-CZ-NH2	5.52	123.06	120.30
1	B	160	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	D	127	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	C	275	ARG	NE-CZ-NH1	5.16	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2800	0	2719	23	0
1	B	2785	0	2702	25	0
1	C	2802	0	2726	16	0
1	D	2808	0	2730	27	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
2	C	4	0	3	0	0
2	D	4	0	3	0	0
3	A	233	0	0	4	0
3	B	177	0	0	0	0
3	C	227	0	0	2	0
3	D	215	0	0	2	0
All	All	12063	0	10889	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:194[B]:CYS:SG	1:D:357:ILE:HD13	1.81	1.19
1:B:223:HIS:N	1:B:226:GLY:O	1.81	1.11
1:C:62:ARG:NH1	1:C:69:GLU:OE2	1.82	1.10
1:A:106:LYS:HD3	1:A:194[A]:CYS:SG	2.09	0.93
1:A:194[B]:CYS:SG	1:A:357:ILE:HD13	2.12	0.90
1:D:106:LYS:HD2	1:D:194[A]:CYS:SG	2.18	0.83
1:A:194[B]:CYS:SG	1:A:357:ILE:CD1	2.73	0.76
1:D:194[B]:CYS:SG	1:D:357:ILE:CD1	2.70	0.75
1:B:222:PRO:HA	1:B:226:GLY:O	1.86	0.74
1:C:101:HIS:HD2	1:C:128:ARG:H	1.37	0.72
1:D:27:ARG:NH2	1:D:59:GLY:HA3	2.08	0.69
1:A:22:HIS:HE1	1:A:105:GLU:OE2	1.85	0.60
1:C:281:MET:HG2	1:D:319:ASN:HD22	1.67	0.59
1:A:168:MET:HE3	1:A:221:ARG:NH2	2.18	0.59
1:A:22:HIS:HD2	1:A:80:SER:OG	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:LYS:HE3	3:A:669:HOH:O	2.03	0.58
1:D:106:LYS:CD	1:D:194[A]:CYS:SG	2.92	0.56
1:B:167:TYR:O	1:B:168:MET:HE2	2.06	0.56
1:B:101:HIS:HD2	1:B:128:ARG:H	1.56	0.54
1:B:17:PHE:CD1	1:B:17:PHE:C	2.81	0.53
1:D:224:GLN:NE2	1:D:224:GLN:HA	2.22	0.53
1:B:71:VAL:CG1	1:B:95:ALA:HA	2.38	0.53
1:D:27:ARG:HH22	1:D:59:GLY:HA3	1.72	0.53
1:B:22:HIS:HE1	1:B:105:GLU:OE2	1.91	0.53
1:B:17:PHE:C	1:B:17:PHE:HD1	2.12	0.52
1:B:83:THR:HB	1:B:84:PRO:HD2	1.90	0.52
1:B:83:THR:HB	1:B:84:PRO:CD	2.40	0.52
1:B:17:PHE:CE1	1:B:18:MET:CE	2.93	0.51
1:B:27:ARG:HD2	1:B:40:LEU:HD11	1.93	0.51
1:C:358:HIS:HE1	3:C:627:HOH:O	1.93	0.50
1:B:222:PRO:CA	1:B:226:GLY:O	2.58	0.50
1:B:17:PHE:CE1	1:B:18:MET:HE1	2.46	0.50
1:D:83:THR:HB	1:D:84:PRO:HD2	1.94	0.49
1:A:106:LYS:HD2	1:A:106:LYS:C	2.32	0.49
1:B:18:MET:HG3	1:B:105:GLU:OE1	2.13	0.49
1:D:83:THR:HB	1:D:84:PRO:CD	2.43	0.49
1:A:336:ALA:HA	1:A:341:GLY:O	2.13	0.48
1:D:39:ARG:NH1	1:D:41:GLU:OE2	2.44	0.48
1:A:209:ARG:NH1	1:C:211:TYR:OH	2.47	0.48
1:B:220:ASP:HB3	1:B:227:THR:OG1	2.14	0.47
1:D:138:SER:HB3	1:D:325:GLN:HE21	1.78	0.47
1:D:43:LEU:C	1:D:43:LEU:HD23	2.35	0.47
1:A:12:LEU:HD23	1:A:43:LEU:HD13	1.96	0.47
1:D:342:LYS:HE2	3:D:604:HOH:O	2.14	0.47
1:B:222:PRO:C	1:B:226:GLY:O	2.51	0.47
1:A:27:ARG:HG2	1:B:28:ASN:ND2	2.29	0.47
1:C:89:ARG:HE	1:C:93:GLU:CD	2.18	0.47
1:A:83:THR:HB	1:A:84:PRO:HD2	1.96	0.47
1:B:170:ASP:HB3	1:B:173:LEU:HD13	1.97	0.46
1:A:194[B]:CYS:SG	1:A:357:ILE:HD12	2.54	0.46
1:A:13:ILE:HG22	1:A:83:THR:CG2	2.45	0.46
1:C:345:PRO:HB2	1:C:349:ASP:HB2	1.97	0.46
1:C:141:PHE:CE1	1:C:199:VAL:HG13	2.51	0.45
1:D:85:ASN:HB2	1:D:110:LEU:HD21	1.98	0.45
1:D:130:ILE:HD12	1:D:343:ALA:HB1	1.98	0.45
1:D:336:ALA:HA	1:D:341:GLY:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:HIS:HD2	1:B:80:SER:OG	1.99	0.45
1:A:83:THR:HB	1:A:84:PRO:CD	2.47	0.44
1:C:20:LYS:HE3	1:D:31:THR:O	2.16	0.44
1:A:138:SER:HB3	1:A:325:GLN:HE21	1.81	0.44
1:D:291:ARG:NH2	1:D:293:GLU:OE1	2.50	0.44
1:D:106:LYS:HE3	1:D:194[A]:CYS:SG	2.57	0.44
1:A:27:ARG:NH2	3:A:604:HOH:O	2.50	0.44
1:C:87:LEU:HD22	1:C:87:LEU:N	2.33	0.44
1:A:130:ILE:HD12	1:A:343:ALA:HB1	2.00	0.43
1:B:130:ILE:HD12	1:B:343:ALA:HB1	2.01	0.43
1:C:37:PRO:HB3	1:D:27:ARG:HH12	1.83	0.43
1:C:182:LYS:HG2	1:C:183:ASP:OD1	2.18	0.43
1:C:123:ARG:NH1	3:C:603:HOH:O	2.52	0.43
1:C:85:ASN:HB3	1:C:87:LEU:H	1.83	0.43
1:D:194[B]:CYS:HG	1:D:357:ILE:HD13	1.76	0.43
1:D:345:PRO:HB2	1:D:349:ASP:HB2	2.02	0.42
1:D:345:PRO:CB	1:D:349:ASP:HB2	2.49	0.42
1:C:345:PRO:CB	1:C:349:ASP:HB2	2.50	0.42
1:B:39:ARG:NH1	1:B:41:GLU:OE2	2.52	0.42
1:A:44:ALA:HA	1:A:64:THR:O	2.20	0.42
1:B:13:ILE:HG22	1:B:83:THR:CG2	2.50	0.41
1:D:182:LYS:O	1:D:183:ASP:HB2	2.21	0.41
1:A:27:ARG:CZ	3:A:730:HOH:O	2.67	0.41
1:A:71:VAL:HG13	1:A:95:ALA:HA	2.03	0.41
1:B:211:TYR:OH	1:D:209:ARG:NH2	2.54	0.41
1:C:130:ILE:HD12	1:C:343:ALA:HB1	2.02	0.41
1:D:27:ARG:NH1	3:D:601:HOH:O	2.35	0.41
1:B:25:ALA:HA	1:B:327:VAL:HG22	2.03	0.40
1:A:213:GLN:NE2	3:A:603:HOH:O	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/380 (96%)	361 (99%)	5 (1%)	0	100	100
1	B	364/380 (96%)	351 (96%)	13 (4%)	0	100	100
1	C	366/380 (96%)	362 (99%)	4 (1%)	0	100	100
1	D	367/380 (97%)	356 (97%)	11 (3%)	0	100	100
All	All	1463/1520 (96%)	1430 (98%)	33 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/288 (96%)	271 (98%)	6 (2%)	52	29
1	B	275/288 (96%)	270 (98%)	5 (2%)	59	40
1	C	277/288 (96%)	273 (99%)	4 (1%)	67	52
1	D	278/288 (96%)	272 (98%)	6 (2%)	52	29
All	All	1107/1152 (96%)	1086 (98%)	21 (2%)	57	37

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	LYS
1	A	134	ASN
1	A	163	TYR
1	A	168	MET
1	A	209	ARG
1	A	227	THR
1	B	17	PHE
1	B	69	GLU
1	B	134	ASN
1	B	163	TYR
1	B	227	THR
1	C	62	ARG
1	C	89	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	134	ASN
1	C	163	TYR
1	D	85	ASN
1	D	106	LYS
1	D	163	TYR
1	D	180	THR
1	D	224	GLN
1	D	373	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	28	ASN
1	A	52	HIS
1	A	306	GLN
1	A	325	GLN
1	B	22	HIS
1	B	28	ASN
1	B	101	HIS
1	B	306	GLN
1	C	101	HIS
1	C	213	GLN
1	C	306	GLN
1	C	358	HIS
1	D	28	ASN
1	D	213	GLN
1	D	224	GLN
1	D	278	GLN
1	D	319	ASN
1	D	325	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	500	-	3,3,3	0.77	0	3,3,3	0.92	0
2	ACT	B	500	-	3,3,3	0.79	0	3,3,3	0.97	0
2	ACT	D	500	-	3,3,3	0.60	0	3,3,3	1.03	0
2	ACT	C	500	-	3,3,3	0.72	0	3,3,3	0.74	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	367/380 (96%)	0.05	10 (2%) 54 60	6, 11, 24, 35	0
1	B	366/380 (96%)	0.36	30 (8%) 11 15	6, 14, 39, 54	0
1	C	368/380 (96%)	0.09	15 (4%) 37 44	6, 11, 30, 42	0
1	D	368/380 (96%)	0.16	20 (5%) 25 32	6, 11, 29, 48	0
All	All	1469/1520 (96%)	0.16	75 (5%) 28 34	6, 12, 32, 54	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	224	GLN	8.7
1	B	226	GLY	7.9
1	D	177	TRP	7.3
1	B	17	PHE	6.0
1	A	177	TRP	5.7
1	B	177	TRP	5.5
1	D	373	LEU	5.0
1	B	175	TRP	4.9
1	B	86	GLY	4.8
1	D	84	PRO	4.7
1	B	181	ARG	4.7
1	A	173	LEU	4.2
1	B	49	ASP	4.2
1	B	224	GLN	4.2
1	B	46	MET	4.1
1	D	225	GLY	4.1
1	B	227	THR	3.9
1	D	340	ARG	3.9
1	B	174	PRO	3.8
1	B	48	ALA	3.8
1	B	225	GLY	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	46	MET	3.6
1	C	49	ASP	3.6
1	B	223	HIS	3.4
1	C	17	PHE	3.4
1	B	229	ARG	3.3
1	B	87	LEU	3.3
1	D	85	ASN	3.3
1	B	85	ASN	3.2
1	D	178	ARG	3.1
1	B	84	PRO	3.1
1	D	180	THR	3.0
1	C	6	LYS	3.0
1	B	52	HIS	3.0
1	C	87	LEU	2.9
1	A	6	LYS	2.9
1	D	6	LYS	2.9
1	B	340	ARG	2.8
1	B	182	LYS	2.7
1	C	229	ARG	2.6
1	B	47	PRO	2.6
1	C	85	ASN	2.6
1	B	125	SER	2.6
1	D	223	HIS	2.5
1	B	68	ARG	2.5
1	A	126	ASP	2.5
1	D	183	ASP	2.5
1	A	340	ARG	2.4
1	D	125	SER	2.4
1	D	342	LYS	2.4
1	C	224	GLN	2.4
1	B	173	LEU	2.3
1	B	365	GLN	2.3
1	B	50	LYS	2.3
1	D	113	GLU	2.2
1	C	373	LEU	2.2
1	D	181	ARG	2.2
1	C	125	SER	2.2
1	A	84	PRO	2.2
1	C	84	PRO	2.2
1	B	123	ARG	2.2
1	D	173	LEU	2.2
1	D	182	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	178	ARG	2.2
1	C	177	TRP	2.2
1	A	224	GLN	2.1
1	A	174	PRO	2.1
1	D	83	THR	2.1
1	A	47	PRO	2.1
1	C	225	GLY	2.1
1	D	123	ARG	2.1
1	C	47	PRO	2.1
1	B	183	ASP	2.1
1	C	340	ARG	2.0
1	B	128	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	B	500	4/4	0.96	0.07	14,15,15,15	0
2	ACT	D	500	4/4	0.97	0.08	12,12,13,14	0
2	ACT	C	500	4/4	0.98	0.09	14,14,14,14	0
2	ACT	A	500	4/4	0.99	0.07	13,13,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.