

# Full wwPDB X-ray Structure Validation Report (i)

Jun 22, 2024 – 06:46 PM EDT

PDB ID : 6BFA

Title : Calcium-Dependent Protein Kinase 1 from Toxoplasma gondii (TgCDPK1) in

complex with inhibitor UW1553

Authors : Merritt, E.A. Deposited on : 2017-10-26

Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Whole archive Similar resolution Metric (#Entries) (#Entries, resolution range(Å)) $R_{free}$ 3140 (2.80-2.80) 130704 Clashscore 141614 3569 (2.80-2.80) Ramachandran outliers 138981 3498 (2.80-2.80) Sidechain outliers 138945 3500 (2.80-2.80) RSRZ outliers 127900 3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			3%		
1	A	484	82%	14%	• •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UW5	A	601	-	-	X	-



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Calmodulin-domain protein kinase 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	٨	464	Total	С	N	О	S	0	7	0
1	A	404	3744	2373	626	725	20	0	1	

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	GLY	-	expression tag	UNP Q9BJF5
A	25	PRO	-	expression tag	UNP Q9BJF5
A	26	GLY	-	expression tag	UNP Q9BJF5
A	27	SER	-	expression tag	UNP Q9BJF5
A	28	MET	-	expression tag	UNP Q9BJF5
A	29	MET	-	expression tag	UNP Q9BJF5

• Molecule 2 is 1-{4-amino-3-[6-(cyclopropyloxy)naphthalen-2-yl]-1H-pyrazolo[3,4-d]pyrimidi n-1-yl}-2-methylpropan-2-ol (three-letter code: UW5) (formula: C<sub>22</sub>H<sub>23</sub>N<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf		
2	Λ	1	Total	С	N	О	0	0
	Α	1	29	22	5	2	0	0

### • Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	87	Total O 87 87	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Calmodulin-domain protein kinase 1





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	48.12Å 72.66Å 65.75Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 99.78° 90.00°	Depositor
Resolution (Å)	36.23 - 2.80	Depositor
Resolution (A)	36.23 - 2.80	EDS
% Data completeness	89.4 (36.23-2.80)	Depositor
(in resolution range)	89.4 (36.23-2.80)	EDS
$R_{merge}$	0.30	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.38 (at 2.81Å)	Xtriage
Refinement program	REFMAC REFMAC_5.8.0158	Depositor
D D.	0.199 , 0.253	Depositor
$R, R_{free}$	0.201 , 0.248	DCC
$R_{free}$ test set	475  reflections  (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.6	Xtriage
Anisotropy	0.500	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.30 , 47.2	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3860	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.96% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UW5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.65	0/3827	0.78	2/5145 (0.0%)	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	145	ARG	NE-CZ-NH1	6.20	123.40	120.30
1	A	434	ARG	N-CA-C	-5.38	96.48	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3744	0	3738	52	1
2	A	29	0	0	12	0
3	A	87	0	0	4	0
All	All	3860	0	3738	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash



### magnitude.

A	A	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$overlap(\AA)$
1:A:112:MET:HE1	2:A:601:UW5:C13	1.89	1.01
1:A:112:MET:HE1	2:A:601:UW5:C17	2.02	0.90
1:A:112:MET:HE1	2:A:601:UW5:C14	2.04	0.87
1:A:58:GLY:CA	2:A:601:UW5:C6	2.55	0.83
1:A:385:LYS:O	1:A:389:ARG:HG3	1.82	0.80
1:A:112:MET:CE	2:A:601:UW5:C14	2.66	0.73
1:A:410:VAL:HG11	1:A:427:PHE:CD2	2.24	0.73
1:A:112:MET:CE	2:A:601:UW5:C17	2.67	0.72
1:A:441:GLU:OE1	1:A:445:ARG:NH1	2.21	0.72
1:A:58:GLY:HA3	2:A:601:UW5:C6	2.21	0.70
1:A:410:VAL:HG11	1:A:427:PHE:CE2	2.28	0.68
1:A:481:LEU:O	3:A:701:HOH:O	2.11	0.68
1:A:345:LEU:HB2	1:A:442:ARG:HH21	1.59	0.67
1:A:58:GLY:N	2:A:601:UW5:C6	2.58	0.67
1:A:413:ALA:C	1:A:415:ASP:H	1.97	0.67
1:A:481:LEU:HD12	3:A:701:HOH:O	1.97	0.63
2:A:601:UW5:C16	3:A:749:HOH:O	2.45	0.63
1:A:438:LEU:HD23	1:A:438:LEU:N	2.16	0.61
1:A:410:VAL:CG1	1:A:427:PHE:CD2	2.84	0.59
1:A:410:VAL:CG1	1:A:427:PHE:CE2	2.86	0.59
1:A:345:LEU:HB2	1:A:442:ARG:NH2	2.17	0.58
1:A:320:VAL:HG12	1:A:320:VAL:O	2.06	0.56
1:A:58:GLY:C	2:A:601:UW5:C6	2.74	0.55
2:A:601:UW5:C16	2:A:601:UW5:C12	2.84	0.54
1:A:413:ALA:C	1:A:415:ASP:N	2.60	0.53
1:A:281:ALA:HB2	1:A:307:TRP:NE1	2.23	0.53
1:A:485:ASP:N	3:A:701:HOH:O	2.37	0.52
1:A:112:MET:HG3	1:A:131:TYR:HE2	1.75	0.51
1:A:397:MET:HE1	1:A:403:VAL:HA	1.92	0.51
1:A:386:GLU:HA	1:A:389:ARG:HD2	1.94	0.49
1:A:410:VAL:HB	1:A:427:PHE:HE2	1.78	0.49
1:A:345:LEU:HD12	1:A:442:ARG:HH21	1.78	0.49
1:A:350:LYS:HE3	1:A:507:ASN:ND2	2.27	0.48
1:A:386:GLU:O	1:A:390:MET:HG2	2.13	0.48
1:A:345:LEU:CB	1:A:442:ARG:HH21	2.27	0.48
1:A:47:PHE:HE2	1:A:398:LEU:HD13	1.79	0.47
1:A:345:LEU:CB	1:A:442:ARG:NH2	2.78	0.47
1:A:397:MET:CE	1:A:403:VAL:HA	2.45	0.46
1:A:108:HIS:CG	1:A:109:PRO:HD2	2.51	0.46
1:A:450:PHE:CE2	1:A:458:ILE:HD12	2.51	0.46
1:A:141:ILE:O	1:A:331:ARG:NH2	2.49	0.45

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:384:TYR:CE2	1:A:388:MET:CE	3.00	0.45
1:A:437:LEU:HA	1:A:440:ARG:HB3	1.98	0.45
1:A:395:ALA:O	1:A:398:LEU:HG	2.17	0.44
1:A:345:LEU:CD1	1:A:442:ARG:HH21	2.31	0.44
1:A:83:SER:HA	1:A:123:TYR:CD2	2.53	0.43
1:A:112:MET:HE2	2:A:601:UW5:C14	2.49	0.43
1:A:149:VAL:O	1:A:153:ARG:HG3	2.18	0.43
1:A:413:ALA:O	1:A:415:ASP:N	2.43	0.42
1:A:324:ASP:O	1:A:327:ILE:HG22	2.19	0.41
1:A:367:MET:HE3	1:A:379:GLU:HB3	2.03	0.41
1:A:317:SER:HB3	1:A:320:VAL:HG21	2.01	0.41
1:A:389:ARG:CB	1:A:390:MET:HE3	2.50	0.41
1:A:440:ARG:O	1:A:440:ARG:HG3	2.16	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\mathring{\mathbf{A}}) \end{aligned}$
1:A:98:ARG:NH2	1:A:483:GLU:OE2[2_544]	2.00	0.20

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	469/484 (97%)	446 (95%)	22 (5%)	1 (0%)	47 78

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	414	VAL



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric		Outliers	Percentiles	
1	A	413/427 (97%)	393 (95%)	20 (5%)	25 58	

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	LYS
1	A	98	ARG
1	A	144	LYS
1	A	156	ARG
1	A	196	PHE
1	A	315	GLN
1	A	316	ILE
1	A	358	LYS
1	A	404	GLU
1	A	417	ASP
1	A	418	LYS
1	A	425	SER
1	A	434	ARG
1	A	435	LYS
1	A	440	ARG
1	A	460	SER
1	A	473	ASP
1	A	479	SER
1	A	486	LYS
1	A	488	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UW5	A	601	-	31,33,33	1.04	3 (9%)	35,50,50	1.24	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UW5	A	601	-	-	4/11/15/15	0/5/5/5

#### All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	A	601	UW5	C7-N2	-3.11	1.32	1.35
2	A	601	UW5	C6-C4	-2.89	1.43	1.52
2	A	601	UW5	C3-C4	2.64	1.57	1.54

#### All (4) bond angle outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	A	601	UW5	C13-O1-C14	-4.38	111.26	119.18

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
2	A	601	UW5	C16-C15-C14	2.87	61.43	59.51
2	A	601	UW5	C3-N1-N2	2.80	122.06	117.25
2	A	601	UW5	O1-C13-C12	-2.64	111.75	123.82

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	UW5	C15-C14-O1-C13
2	A	601	UW5	C17-C13-O1-C14
2	A	601	UW5	C12-C13-O1-C14
2	A	601	UW5	C16-C14-O1-C13


There are no ring outliers.

1 monomer is involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	UW5	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	464/484 (95%)	-0.16	13 (2%) 53 43	36, 64, 119, 166	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	A	416 PHE		7.1	
1	A	393	GLN	4.4	
1	A	44	THR	3.6	
1	A	314	GLU	3.4	
1	A	369	LYS	3.2	
1	A	415	ASP	3.2	
1	A	417	ASP	3.2	
1	A	420	GLY	3.1	
1	A	428	VAL	2.9	
1	A	411	LEU	2.6	
1	A	188	ASP	2.3	
1	A	422	ILE	2.3	
1	1 A		ARG	2.0	

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	UW5	A	601	29/29	0.80	0.33	37,39,44,45	29

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers (i)

There are no such residues in this entry.

