

Full wwPDB X-ray Structure Validation Report (i)

Jun 23, 2024 – 01:01 AM EDT

PDB ID : 6ERO

Title: Structure of human TFB2M

Authors: Hillen, H.S.; Morozov, Y.I.; Sarfallah, A.; Temiakov, D.; Cramer, P.

Deposited on : 2017-10-18

Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : FAILED EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain	
1	A	313	90%	• 5%
1	В	313	85%	• 12%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9638 atoms, of which 4700 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dimethyladenosine transferase 2, mitochondrial, Dimethylade nosine transferase 2, mitochondrial.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace	
1	A	296		C 1537	H 2414	N 408	O 422	S 15	0	0	0
1	В	277	Total 4515	_	H 2270		O 393	S 15	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ASN	-	expression tag	UNP Q9H5Q4
A	2	ALA	-	expression tag	UNP Q9H5Q4
A	208	GLY	-	linker	UNP Q9H5Q4
A	209	SER	-	linker	UNP Q9H5Q4
A	210	SER	-	linker	UNP Q9H5Q4
A	211	GLY	-	linker	UNP Q9H5Q4
В	1	ASN	-	expression tag	UNP Q9H5Q4
В	2	ALA	-	expression tag	UNP Q9H5Q4
В	208	GLY	-	linker	UNP Q9H5Q4
В	209	SER	-	linker	UNP Q9H5Q4
В	210	SER	-	linker	UNP Q9H5Q4
В	211	GLY	-	linker	UNP Q9H5Q4

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cl 2 2	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Λ	1	Total	С	Н	О	0	0	
3	3 A	1	14	3	8	3	0		
2	D	1	Total	С	Н	О	0	0	
3	Б	1	14	3	8	3		U	

• Molecule 4 is water.

Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	190	Total O 190 190	0	0
4	В	107	Total O 107 107	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

• Molecule 1: Dimethyladenosine transferase 2, mitochondrial,Dimethyladenosine transferase 2, mitochondrial



 \bullet Molecule 1: Dimethyladenosine transferase 2, mitochondrial, Dimethyladenosine transferase 2, mitochondrial





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	43.95Å 165.65Å 44.73Å	Depositor	
a, b, c, α , β , γ	90.00° 97.97° 90.00°	Depositor	
Resolution (Å)	43.52 - 1.75	Depositor	
% Data completeness	97.8 (43.52-1.75)	Depositor	
(in resolution range)	31.0 (40.02 1.10)	Depositor	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor	
R, R_{free}	0.192 , 0.214	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	9638	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	57.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.26	0/2437	0.48	0/3290	
1	В	0.27	0/2297	0.47	0/3100	
All	All	0.26	0/4734	0.48	0/6390	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2382	2414	2426	10	0
1	В	2245	2270	2285	11	0
2	A	2	0	0	0	0
3	A	6	8	8	0	0
3	В	6	8	8	0	0
4	A	190	0	0	1	0
4	В	107	0	0	5	0
All	All	4938	4700	4727	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
7100111 1	7100H1 2	$\operatorname{distance} \left(\operatorname{\AA} \right)$	overlap (Å)
1:A:104:PRO:HB2	1:A:105:PRO:HD3	1.73	0.70
1:B:222:GLN:NE2	4:B:501:HOH:O	2.29	0.64
1:B:18:ARG:NH2	4:B:505:HOH:O	2.33	0.61
1:B:247:ARG:NH2	4:B:506:HOH:O	2.35	0.60
1:B:141:LYS:NZ	4:B:502:HOH:O	2.29	0.60
1:B:124:THR:O	1:B:222:GLN:NE2	2.39	0.56
1:A:104:PRO:CB	1:A:105:PRO:HD3	2.36	0.54
1:A:40:LEU:HD13	1:A:129:LEU:HD11	1.92	0.51
1:B:29:LEU:HD11	1:B:56:ALA:HB1	1.94	0.49
1:B:67:LYS:HG3	1:B:88:HIS:CE1	2.48	0.48
1:B:67:LYS:HE3	1:B:88:HIS:CE1	2.49	0.47
1:A:184:PRO:O	1:A:248:ARG:NH2	2.48	0.47
1:B:130:LYS:HE3	1:B:164:GLU:OE2	2.17	0.45
1:A:180:ASP:HB2	1:A:181:PRO:CD	2.47	0.44
1:B:67:LYS:HE3	1:B:88:HIS:ND1	2.34	0.42
1:A:103:LYS:O	1:A:106:ALA:O	2.38	0.42
1:B:274:GLN:NE2	4:B:515:HOH:O	2.53	0.41
1:A:101:VAL:HG12	1:A:101:VAL:O	2.20	0.41
1:A:40:LEU:HD22	1:A:129:LEU:HD21	2.02	0.41
1:A:29:LEU:HD21	1:A:58:ALA:HB2	2.03	0.40
1:A:188:HIS:HB2	4:A:633:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	$292/313 \ (93\%)$	283 (97%)	9 (3%)	0	100	100
1	В	269/313~(86%)	265 (98%)	4 (2%)	0	100	100
All	All	561/626 (90%)	548 (98%)	13 (2%)	0	100	100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	$261/275 \ (95\%)$	259 (99%)	2 (1%)	81 72		
1	В	246/275 (90%)	246 (100%)	0	100 100		
All	All	507/550~(92%)	505 (100%)	2 (0%)	91 87		

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	SER
1	A	188	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain Dag	T in le	Bond lengths			Bond angles		
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	В	401	-	5,5,5	0.41	0	5,5,5	0.17	0
3	GOL	A	403	-	5,5,5	0.38	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	В	401	-	-	1/4/4/4	-
3	GOL	A	403	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	GOL	O1-C1-C2-C3
3	A	403	GOL	O1-C1-C2-O2
3	В	401	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

6.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

