



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 10:24 PM EDT

PDB ID : 6F5H
Title : Crystal structure of USP7 in complex with a 4-hydroxypiperidine based inhibitor
Authors : Harrison, T.; Gavory, G.; O'Dowd, C.; Helm, M.; Flasz, J.; Dossang, A.; Hughes, C.; Cassidy, E.; McClelland, K.; Odrzywol, E.; Page, N.; Barker, O.; Miel, H.; Feutron-Burton, S.; Rountree, J.S.S.
Deposited on : 2017-12-01
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 14% 89% 8%
1	B	357	 20% 81% 11% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6093 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	350	2859	1803	497	543	16	0	0	0
1	B	337	2741	1735	467	523	16	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	561	HIS	-	expression tag	UNP Q93009
A	562	HIS	-	expression tag	UNP Q93009
A	563	HIS	-	expression tag	UNP Q93009
B	561	HIS	-	expression tag	UNP Q93009
B	562	HIS	-	expression tag	UNP Q93009
B	563	HIS	-	expression tag	UNP Q93009

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is 3-[[4-oxidanyl-1-[(3 {R})-3-phenylbutanoyl]piperidin-4-yl]methyl]-6-(2-pyrrolidin-1-ylethylamino)pyrimidin-4-one (three-letter code: CQ5) (formula: C₂₆H₃₇N₅O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	34	26	5	3	0	0
4	B	1	34	26	5	3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	222	222	222	0	0
5	B	182	182	182	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7



- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.86Å 67.33Å 80.69Å 90.00° 105.06° 90.00°	Depositor
Resolution (Å)	29.95 – 2.16 28.22 – 2.16	Depositor EDS
% Data completeness (in resolution range)	95.5 (29.95-2.16) 95.6 (28.22-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.202 , 0.262 0.210 , 0.265	Depositor DCC
R_{free} test set	1958 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtrriage
Anisotropy	0.510	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6093	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CQ5, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2920	0.69	1/3933 (0.0%)
1	B	0.54	0/2800	0.71	1/3776 (0.0%)
All	All	0.53	0/5720	0.70	2/7709 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	454	LEU	CA-CB-CG	5.12	127.09	115.30
1	B	454	LEU	CA-CB-CG	5.11	127.05	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	510	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2859	0	2794	11	0
1	B	2741	0	2677	18	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	6	0	8	0	0
4	A	34	0	0	0	0
4	B	34	0	0	0	0
5	A	222	0	0	0	0
5	B	182	0	0	1	0
All	All	6093	0	5479	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:241:ALA:HA	1:B:316:VAL:HG21	1.74	0.69
1:A:424:ARG:NH1	1:A:426:GLU:OE2	2.32	0.62
1:A:227:SER:HB3	1:A:454:LEU:HD13	1.86	0.58
1:A:524:LEU:HD22	1:A:528:LEU:HD12	1.85	0.57
1:B:350:ILE:HD12	1:B:402:LEU:HD21	1.86	0.57
1:B:250:ASP:OD2	5:B:1101:HOH:O	2.18	0.57
1:B:231:THR:CG2	1:B:517:VAL:HG21	2.38	0.53
1:B:431:LEU:HG	1:B:433:LEU:HD13	1.91	0.52
1:B:228:LEU:HD23	1:B:299:LEU:HD13	1.92	0.52
1:B:453:VAL:HG12	1:B:455:VAL:HG23	1.93	0.50
1:B:237:GLN:HA	1:B:240:LYS:HD3	1.95	0.49
1:B:234:PHE:CD1	1:B:471:PRO:HB3	2.47	0.49
1:A:285:TRP:CZ3	1:A:299:LEU:HG	2.48	0.48
1:B:241:ALA:HB2	1:B:319:THR:HG21	1.96	0.48
1:B:285:TRP:CZ3	1:B:299:LEU:HG	2.50	0.47
1:B:279:LEU:HD22	1:B:283:PHE:HE2	1.80	0.47
1:A:241:ALA:HA	1:A:316:VAL:HG21	1.98	0.45
1:B:247:THR:HG22	1:B:310:LYS:HE3	1.98	0.44
1:A:316:VAL:HG22	1:A:319:THR:OG1	2.17	0.44
1:A:448:TYR:HB3	1:A:518:TYR:HB3	2.00	0.44
1:B:329:VAL:HG13	1:B:396:LEU:HD11	1.99	0.43
1:B:457:SER:HB2	1:B:464:HIS:HB3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:279:LEU:HD22	1:B:283:PHE:CE2	2.53	0.43
1:A:250:ASP:O	1:A:543:ARG:NH2	2.48	0.43
1:B:536:ILE:HG22	1:B:541:VAL:HG23	2.01	0.43
1:B:354:ILE:HG22	1:B:425:PHE:CE1	2.54	0.42
1:A:260:LEU:HD12	1:A:307:VAL:HG11	2.01	0.42
1:A:288:LEU:HD22	1:A:292:MET:HG3	2.03	0.41
1:A:297:GLN:OE1	1:A:405:GLN:HG2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/357 (97%)	338 (98%)	8 (2%)	0	100	100
1	B	333/357 (93%)	323 (97%)	10 (3%)	0	100	100
All	All	679/714 (95%)	661 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/324 (98%)	297 (94%)	20 (6%)	18	13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	305/324 (94%)	284 (93%)	21 (7%)	15	10
All	All	622/648 (96%)	581 (93%)	41 (7%)	16	11

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	SER
1	A	209	LYS
1	A	258	LEU
1	A	260	LEU
1	A	279	LEU
1	A	288	LEU
1	A	299	LEU
1	A	316	VAL
1	A	322	LYS
1	A	402	LEU
1	A	409	PHE
1	A	419	ILE
1	A	454	LEU
1	A	457	SER
1	A	469	LEU
1	A	470	ASN
1	A	524	LEU
1	A	540	LEU
1	A	543	ARG
1	A	561	HIS
1	B	208	LYS
1	B	219	GLN
1	B	252	SER
1	B	258	LEU
1	B	260	LEU
1	B	279	LEU
1	B	288	LEU
1	B	299	LEU
1	B	373	LEU
1	B	390	GLU
1	B	402	LEU
1	B	409	PHE
1	B	433	LEU
1	B	454	LEU
1	B	457	SER
1	B	469	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	470	ASN
1	B	496	HIS
1	B	510	CYS
1	B	540	LEU
1	B	549	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	GLN
1	A	309	ASN
1	A	438	GLN
1	B	403	HIS
1	B	545	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	1001	-	4,4,4	0.45	0	6,6,6	0.17	0
2	SO4	A	1001	-	4,4,4	0.41	0	6,6,6	0.14	0
4	CQ5	A	1003	-	34,37,37	0.93	2 (5%)	41,51,51	1.79	11 (26%)
2	SO4	B	1002	-	4,4,4	0.42	0	6,6,6	0.09	0
4	CQ5	B	1003	-	34,37,37	1.01	2 (5%)	41,51,51	1.83	12 (29%)
3	GOL	A	1002	-	5,5,5	0.32	0	5,5,5	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CQ5	B	1003	-	-	2/23/42/42	0/4/4/4
4	CQ5	A	1003	-	-	4/23/42/42	0/4/4/4
3	GOL	A	1002	-	-	0/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1003	CQ5	C12-C5	2.83	1.55	1.52
4	A	1003	CQ5	C12-C5	2.67	1.55	1.52
4	B	1003	CQ5	C16-N4	2.30	1.39	1.35
4	A	1003	CQ5	C16-N4	2.29	1.39	1.35

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1003	CQ5	C2-N1-C3	-4.58	119.48	123.00
4	A	1003	CQ5	N1-C2-N	-4.10	122.45	126.45
4	A	1003	CQ5	C17-C16-N4	3.96	125.67	118.37
4	B	1003	CQ5	C3-C-C1	-3.96	118.94	121.48
4	A	1003	CQ5	C2-N1-C3	-3.92	119.98	123.00
4	A	1003	CQ5	C3-C-C1	-3.82	119.03	121.48
4	B	1003	CQ5	N1-C2-N	-3.82	122.72	126.45
4	B	1003	CQ5	C17-C16-N4	3.68	125.16	118.37
4	B	1003	CQ5	C5-C4-N1	-3.56	107.11	113.99
4	A	1003	CQ5	C5-C4-N1	-3.19	107.83	113.99
4	B	1003	CQ5	O2-C16-C17	-3.04	116.39	122.07
4	A	1003	CQ5	O2-C16-N4	-2.91	116.94	122.12
4	B	1003	CQ5	C2-N-C1	2.89	120.78	113.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	CQ5	C2-N-C1	2.79	120.53	113.53
4	A	1003	CQ5	O-C3-C	-2.56	119.30	125.61
4	A	1003	CQ5	O2-C16-C17	-2.52	117.36	122.07
4	B	1003	CQ5	O-C3-C	-2.42	119.64	125.61
4	A	1003	CQ5	C15-C14-N4	2.31	115.14	110.82
4	B	1003	CQ5	C8-N3-C11	2.26	106.71	104.04
4	A	1003	CQ5	O-C3-N1	2.22	121.63	119.51
4	B	1003	CQ5	C10-C11-N3	2.10	106.39	103.90
4	B	1003	CQ5	O2-C16-N4	-2.06	118.45	122.12
4	B	1003	CQ5	C6-N2-C1	2.05	126.37	122.81

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1003	CQ5	O2-C16-C17-C18
4	A	1003	CQ5	O2-C16-N4-C13
4	A	1003	CQ5	N4-C16-C17-C18
4	B	1003	CQ5	O2-C16-C17-C18
4	B	1003	CQ5	N4-C16-C17-C18
4	A	1003	CQ5	O2-C16-N4-C14

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/357 (98%)	0.84	50 (14%) 2 3	35, 51, 79, 115	0
1	B	337/357 (94%)	0.99	70 (20%) 1 1	36, 53, 98, 116	0
All	All	687/714 (96%)	0.92	120 (17%) 1 1	35, 52, 92, 116	0

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	561	HIS	8.3
1	A	558	ARG	6.8
1	B	376	ASP	6.1
1	A	554	LYS	5.8
1	A	376	ASP	5.5
1	A	228	LEU	5.3
1	B	338	ASP	5.2
1	B	509	HIS	5.1
1	A	377	ASN	5.0
1	B	454	LEU	4.9
1	B	382	GLY	4.8
1	A	560	GLU	4.7
1	B	232	LEU	4.6
1	B	373	LEU	4.6
1	B	336	GLU	4.6
1	A	454	LEU	4.6
1	B	377	ASN	4.5
1	B	253	SER	4.4
1	A	562	HIS	4.4
1	A	563	HIS	4.4
1	B	414	GLN	4.3
1	A	404	LEU	4.3
1	B	344	ARG	4.2
1	B	517	VAL	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	229	LEU	4.1
1	A	231	THR	4.1
1	B	413	PRO	4.1
1	B	416	ASP	4.1
1	B	389	ALA	4.0
1	A	516	LEU	4.0
1	B	228	LEU	3.9
1	B	411	TYR	3.9
1	A	232	LEU	3.9
1	B	452	ALA	3.8
1	A	517	VAL	3.8
1	B	412	ASP	3.6
1	B	501	HIS	3.6
1	B	415	THR	3.5
1	B	515	MET	3.5
1	A	559	GLN	3.4
1	A	515	MET	3.4
1	A	338	ASP	3.4
1	B	339	TYR	3.3
1	B	453	VAL	3.3
1	A	227	SER	3.2
1	B	300	CYS	3.2
1	B	335	LYS	3.2
1	B	417	GLN	3.2
1	A	467	VAL	3.2
1	A	336	GLU	3.1
1	B	467	VAL	3.1
1	B	374	ASP	3.1
1	B	248	GLU	3.0
1	A	339	TYR	3.0
1	B	375	GLY	3.0
1	B	271	ASP	3.0
1	B	337	VAL	3.0
1	B	329	VAL	2.9
1	A	442	PRO	2.9
1	A	556	LYS	2.9
1	A	229	LEU	2.9
1	B	466	VAL	2.9
1	A	264	PHE	2.9
1	B	469	LEU	2.9
1	B	249	GLY	2.8
1	B	251	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	557	GLU	2.8
1	A	555	ARG	2.8
1	B	442	PRO	2.8
1	A	453	VAL	2.8
1	B	334	CYS	2.8
1	A	452	ALA	2.7
1	B	343	ARG	2.7
1	A	509	HIS	2.6
1	A	293	GLN	2.6
1	A	441	ASP	2.6
1	B	233	PHE	2.6
1	A	469	LEU	2.6
1	B	516	LEU	2.6
1	B	340	ARG	2.6
1	B	342	ASP	2.6
1	B	341	SER	2.6
1	B	404	LEU	2.6
1	B	441	ASP	2.5
1	A	233	PHE	2.5
1	A	375	GLY	2.5
1	A	235	THR	2.4
1	A	337	VAL	2.4
1	A	234	PHE	2.4
1	B	551	GLU	2.4
1	B	402	LEU	2.4
1	A	455	VAL	2.4
1	B	210	HIS	2.4
1	A	386	LEU	2.3
1	A	514	TYR	2.3
1	B	455	VAL	2.3
1	A	220	GLY	2.3
1	B	231	THR	2.2
1	B	350	ILE	2.2
1	B	227	SER	2.2
1	B	296	VAL	2.2
1	B	394	LYS	2.2
1	B	393	VAL	2.2
1	B	384	HIS	2.1
1	B	410	MET	2.1
1	A	374	ASP	2.1
1	A	304	LEU	2.1
1	A	553	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	238	LEU	2.1
1	B	273	PRO	2.1
1	B	254	LYS	2.1
1	B	230	GLN	2.1
1	B	372	GLN	2.0
1	A	466	VAL	2.0
1	B	331	TYR	2.0
1	A	242	VAL	2.0
1	B	264	PHE	2.0
1	B	549	ARG	2.0
1	A	405	GLN	2.0
1	B	250	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	1001	5/5	0.88	0.18	67,81,94,94	0
3	GOL	A	1002	6/6	0.88	0.15	62,64,66,69	0
2	SO4	B	1002	5/5	0.90	0.38	99,103,105,110	0
4	CQ5	A	1003	34/34	0.90	0.20	36,44,51,53	0
4	CQ5	B	1003	34/34	0.91	0.17	39,45,56,61	0
2	SO4	B	1001	5/5	0.96	0.11	64,65,70,75	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CQ5 A 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CQ5 B 1003:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.