



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 01:24 am GMT

PDB ID : 6FGE
Title : Crystal structure of human ZUFSP/ZUP1 in complex with ubiquitin
Authors : Kwasna, D.; Abdul Rehman, S.A.; Kulathu, Y.
Deposited on : 2018-01-10
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1043 (1.74-1.74)
Clashscore	180529	1119 (1.74-1.74)
Ramachandran outliers	177936	1112 (1.74-1.74)
Sidechain outliers	177891	1112 (1.74-1.74)
RSRZ outliers	164620	1043 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	 13% (red), 88% (green), 11% (yellow), . (grey)
2	C	76	 4% (red), 97% (green), . (grey)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	604	-	-	X	-
7	PEG	A	613	-	-	X	-

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 3607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zinc finger with UFM1-specific peptidase domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	343	2745	1721	496	509	19	0	5	0


- Molecule 2 is a protein called Polyubiquitin-B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	76	601	379	105	116	1	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	76	AYE	GLY	engineered mutation	UNP P0CG47

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			3	1	2		
6	A	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			3	1	2		
6	A	1	Total	C	O	0	0
			3	1	2		
6	A	1	Total	C	O	0	0
			3	1	2		
6	C	1	Total	C	O	0	0
			3	1	2		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is AMMONIUM ION (three-letter code: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total N 1 1	0	0
8	A	1	Total N 1 1	0	0

- Molecule 9 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	C	1	Total C O 4 2 2	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	160	Total 160	O 160	0	0
10	C	36	Total 36	O 36	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Zinc finger with UFM1-specific peptidase domain protein



- Molecule 2: Polyubiquitin-B



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	84.48Å 84.48Å 201.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	73.16 – 1.74 73.16 – 1.74	Depositor EDS
% Data completeness (in resolution range)	100.0 (73.16-1.74) 100.0 (73.16-1.74)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 1.74Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.174 , 0.209 0.186 , 0.222	Depositor DCC
R_{free} test set	2308 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtrriage
Anisotropy	0.003	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3607	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NH4, FMT, AYE, EDO, MLI, ACT, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	0/2806	0.84	0/3773
2	C	0.87	0/603	0.86	0/811
All	All	0.83	0/3409	0.84	0/4584

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	0	2718	38	0
2	C	601	0	630	2	0
3	A	7	0	2	0	0
4	A	12	0	18	9	0
5	A	12	0	16	1	0
6	A	18	0	6	0	0
6	C	3	0	1	0	0
7	A	7	0	10	4	0
8	A	2	0	0	0	0
9	C	4	0	3	1	0
10	A	160	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	C	36	0	0	0	0
All	All	3607	0	3404	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:455:HIS:CD2	4:A:604:EDO:H22	1.77	1.18
1:A:520:MET:HE3	1:A:523:LEU:HD12	1.35	1.09
1:A:455:HIS:HD2	4:A:604:EDO:H22	0.98	1.07
1:A:455:HIS:HD2	4:A:604:EDO:C2	1.85	0.89
1:A:520:MET:CE	1:A:523:LEU:HD12	2.02	0.89
1:A:374[B]:GLN:HG3	10:A:738:HOH:O	1.75	0.86
1:A:455:HIS:CD2	4:A:604:EDO:C2	2.65	0.77
1:A:246:ARG:HG2	1:A:246:ARG:HH11	1.52	0.74
1:A:445:PHE:HB3	4:A:604:EDO:H11	1.71	0.72
1:A:246:ARG:HG2	1:A:246:ARG:NH1	2.04	0.72
1:A:534:LYS:HD2	7:A:613:PEG:O4	1.90	0.70
1:A:445:PHE:HD2	4:A:604:EDO:H11	1.55	0.70
1:A:445:PHE:CD2	4:A:604:EDO:H11	2.28	0.69
1:A:362:TYR:CB	1:A:390:ILE:HD12	2.28	0.64
1:A:362:TYR:HB3	1:A:390:ILE:HD12	1.80	0.62
1:A:243:GLU:OE1	1:A:243:GLU:HA	2.02	0.59
1:A:241:GLN:O	1:A:244:GLU:HG3	2.03	0.58
1:A:537:ARG:O	7:A:613:PEG:H41	2.05	0.57
1:A:246:ARG:HH11	1:A:246:ARG:CG	2.17	0.53
1:A:348:HIS:ND1	1:A:520:MET:HG3	2.23	0.53
1:A:322:ILE:HD11	1:A:520:MET:HE2	1.89	0.53
1:A:445:PHE:CB	4:A:604:EDO:H11	2.38	0.52
1:A:455:HIS:HE1	1:A:545:HIS:O	1.94	0.50
1:A:567:GLN:O	1:A:570:GLN:HG2	2.12	0.49
1:A:276:LYS:HE3	1:A:305:MET:SD	2.52	0.49
1:A:308:LEU:HD22	1:A:391:PRO:HG2	1.95	0.49
1:A:292:MET:HE3	1:A:297:PHE:HB2	1.94	0.48
1:A:539:SER:HB3	7:A:613:PEG:H31	1.95	0.48
1:A:275:TYR:OH	5:A:605:GOL:H12	2.13	0.47
1:A:503:ASN:O	1:A:504:ARG:HB2	2.13	0.47
1:A:539:SER:OG	7:A:613:PEG:H21	2.14	0.47
1:A:286:GLU:OE1	1:A:291:ARG:HD2	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:GLU:HG2	2:C:68:HIS:CE1	2.51	0.46
1:A:284:GLU:HG3	1:A:297:PHE:CE2	2.51	0.45
2:C:71:LEU:H	9:C:101:ACT:H3	1.82	0.45
1:A:390:ILE:HB	1:A:391:PRO:HD3	1.98	0.44
1:A:445:PHE:HB3	4:A:604:EDO:C1	2.45	0.43
1:A:344:SER:O	1:A:346:VAL:HG23	2.17	0.43
1:A:301:LYS:O	1:A:305:MET:HG2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/348 (99%)	339 (98%)	7 (2%)	0	100	100
2	C	73/76 (96%)	72 (99%)	1 (1%)	0	100	100
All	All	419/424 (99%)	411 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/305 (96%)	292 (99%)	2 (1%)	81	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	68/68 (100%)	68 (100%)	0	100	100
All	All	362/373 (97%)	360 (99%)	2 (1%)	84	78

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	SER
1	A	471	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	364	ASN
1	A	441	HIS
1	A	455	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are modelled with single atom - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FMT	A	608	-	2,2,2	0.65	0	1,1,1	0.56	0
6	FMT	A	612	-	2,2,2	0.70	0	1,1,1	0.86	0
6	FMT	A	611	-	2,2,2	0.77	0	1,1,1	0.67	0
6	FMT	A	609	-	2,2,2	0.59	0	1,1,1	0.83	0
4	EDO	A	602	-	3,3,3	0.60	0	2,2,2	0.21	0
4	EDO	A	603	-	3,3,3	0.46	0	2,2,2	0.34	0
3	MLI	A	601	-	6,6,6	1.17	0	7,7,7	0.83	0
4	EDO	A	604	-	3,3,3	0.46	0	2,2,2	0.34	0
5	GOL	A	605	-	5,5,5	0.41	0	5,5,5	0.88	0
5	GOL	A	606	-	5,5,5	0.46	0	5,5,5	0.52	0
6	FMT	A	607	-	2,2,2	0.81	0	1,1,1	0.51	0
6	FMT	A	610	-	2,2,2	0.80	0	1,1,1	0.70	0
6	FMT	C	102	-	2,2,2	0.80	0	1,1,1	0.83	0
7	PEG	A	613	-	6,6,6	0.39	0	5,5,5	0.26	0
9	ACT	C	101	-	3,3,3	0.85	0	3,3,3	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	602	-	-	0/1/1/1	-
4	EDO	A	603	-	-	1/1/1/1	-
3	MLI	A	601	-	-	0/4/4/4	-
4	EDO	A	604	-	-	0/1/1/1	-
5	GOL	A	605	-	-	2/4/4/4	-
5	GOL	A	606	-	-	0/4/4/4	-
7	PEG	A	613	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	605	GOL	O1-C1-C2-O2
5	A	605	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	603	EDO	O1-C1-C2-O2
7	A	613	PEG	C1-C2-O2-C3

There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	604	EDO	9	0
5	A	605	GOL	1	0
7	A	613	PEG	4	0
9	C	101	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/348 (98%)	0.56	46 (13%) 8 11	12, 28, 75, 120	5 (1%)
2	C	75/76 (98%)	0.19	3 (4%) 43 50	18, 32, 52, 60	0
All	All	418/424 (98%)	0.49	49 (11%) 10 14	12, 29, 74, 120	5 (1%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	LEU	8.2
1	A	237	ALA	6.3
1	A	470	GLY	5.3
1	A	471	GLU	5.1
1	A	238	HIS	5.1
1	A	530	ALA	5.0
1	A	533	LEU	4.8
1	A	529	GLU	4.7
1	A	271	ASN	4.6
1	A	290	GLY	4.1
1	A	384	GLY	4.0
1	A	528	ILE	4.0
1	A	240	LEU	3.9
1	A	449	THR	3.7
1	A	239	GLN	3.4
1	A	531	SER	3.3
1	A	273	GLY	3.3
1	A	532	SER	3.3
1	A	272	SER	3.3
1	A	526	GLN	3.2
1	A	527	ASP	3.0
1	A	334	ALA	3.0
1	A	294	PRO	2.9
1	A	298	HIS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	289	ARG	2.8
1	A	523	LEU	2.8
1	A	251	GLU	2.7
1	A	472	GLY	2.7
1	A	468	SER	2.7
1	A	248	ARG	2.7
1	A	241	GLN	2.6
1	A	469	GLU	2.6
1	A	242	GLN	2.6
1	A	440	CYS	2.6
1	A	270	ASP	2.6
1	A	295	SER	2.5
1	A	380	ASP	2.5
1	A	473	SER	2.5
1	A	246	ARG	2.5
1	A	335	THR	2.4
2	C	61	ILE	2.4
2	C	24	GLU	2.4
2	C	33	LYS	2.3
1	A	535	GLN	2.3
1	A	244	GLU	2.2
1	A	381	CYS	2.2
1	A	293	PRO	2.2
1	A	247	LYS	2.1
1	A	261	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	PEG	A	613	7/7	0.56	0.24	56,60,74,76	0
9	ACT	C	101	4/4	0.72	0.20	32,34,48,54	0
6	FMT	A	609	3/3	0.75	0.16	48,48,55,59	0
6	FMT	A	611	3/3	0.76	0.16	54,54,54,68	0
4	EDO	A	603	4/4	0.77	0.18	33,34,46,53	0
5	GOL	A	605	6/6	0.82	0.21	31,52,57,78	0
6	FMT	C	102	3/3	0.83	0.14	46,46,46,58	0
6	FMT	A	610	3/3	0.84	0.14	46,46,51,52	0
5	GOL	A	606	6/6	0.84	0.15	32,43,53,76	0
6	FMT	A	612	3/3	0.84	0.13	34,34,40,43	0
4	EDO	A	604	4/4	0.85	0.29	26,26,31,45	0
8	NH4	A	615	1/1	0.86	0.24	47,47,47,47	0
8	NH4	A	614	1/1	0.86	0.15	31,31,31,31	0
6	FMT	A	608	3/3	0.88	0.19	37,37,43,46	0
4	EDO	A	602	4/4	0.90	0.10	24,26,27,27	0
6	FMT	A	607	3/3	0.90	0.15	44,44,47,51	0
3	MLI	A	601	7/7	0.96	0.07	23,24,26,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.