



# Full wwPDB X-ray Structure Validation Report i

Jun 22, 2024 – 01:34 PM EDT

PDB ID : 6FZY  
Title : PPAR mutant  
Authors : Rochel, N.  
Deposited on : 2018-03-15  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 4307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C 2136	N 1375	O 347	S 404	10	0	0
1	B	269	Total	C 2124	N 1369	O 345	S 399	11	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	227	GLY	-	expression tag	UNP P37231
A	228	SER	-	expression tag	UNP P37231
A	229	HIS	-	expression tag	UNP P37231
A	230	MET	-	expression tag	UNP P37231
A	475	MET	THR	engineered mutation	UNP P37231
B	227	GLY	-	expression tag	UNP P37231
B	228	SER	-	expression tag	UNP P37231
B	229	HIS	-	expression tag	UNP P37231
B	230	MET	-	expression tag	UNP P37231
B	475	MET	THR	engineered mutation	UNP P37231


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	22	Total O 22 22	0	0
2	B	25	Total O 25 25	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.19 Å 62.67 Å 165.53 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.65 – 3.10 42.65 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.65-3.10) 98.1 (42.65-2.85)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.10 (at 2.86 Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
$R$ , $R_{free}$	0.195, 0.235 0.196, 0.244	Depositor DCC
$R_{free}$ test set	778 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	97.2	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33, 108.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.177 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4307	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.46	0/2172	0.64	0/2931
1	B	0.48	0/2159	0.66	0/2912
All	All	0.47	0/4331	0.65	0/5843

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2136	0	2156	15	0
1	B	2124	0	2150	10	0
2	A	22	0	0	0	0
2	B	25	0	0	0	0
All	All	4307	0	4306	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:428:LEU:HD12	1:A:431:VAL:HG22	1.72	0.69

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:ILE:HG23	1:A:416:ILE:HD12	1.77	0.66
1:B:458:GLN:OE1	1:B:458:GLN:HA	1.99	0.63
1:A:423:GLY:O	1:A:428:LEU:HD11	2.04	0.58
1:A:425:ARG:O	1:A:428:LEU:HD21	2.09	0.52
1:A:425:ARG:HB2	1:A:428:LEU:HD21	1.93	0.50
1:A:407:LEU:HD11	1:A:463:LEU:HD13	1.95	0.48
1:A:428:LEU:HD12	1:A:431:VAL:CG2	2.43	0.48
1:A:429:LEU:H	1:A:429:LEU:HD23	1.79	0.47
1:B:407:LEU:HD11	1:B:463:LEU:HD13	1.95	0.47
1:B:467:MET:HA	1:B:470:LEU:HD12	1.96	0.46
1:A:429:LEU:HD23	1:A:429:LEU:N	2.31	0.46
1:A:467:MET:HA	1:A:470:LEU:HD12	1.98	0.46
1:A:317:SER:O	1:A:321:VAL:HG23	2.16	0.45
1:A:456:SER:HB3	1:A:459:LEU:HB2	1.99	0.45
1:A:353:ILE:HD12	1:A:416:ILE:HG23	1.98	0.45
1:B:317:SER:O	1:B:321:VAL:HG23	2.17	0.45
1:B:258:LYS:HG3	1:B:360:SER:HB3	1.98	0.44
1:B:353:ILE:HD12	1:B:416:ILE:HG23	1.99	0.44
1:B:230:MET:SD	1:B:450:LYS:HD2	2.58	0.43
1:A:398:PHE:HB2	1:A:473:ILE:HD11	2.02	0.42
1:B:481:LEU:HA	1:B:484:ILE:HD12	2.00	0.42
1:B:398:PHE:HB2	1:B:473:ILE:HD11	2.02	0.41
1:B:470:LEU:O	1:B:474:VAL:HG23	2.21	0.41
1:A:425:ARG:C	1:A:428:LEU:HD21	2.41	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	266/279 (95%)	255 (96%)	11 (4%)	0	100 100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	265/279 (95%)	255 (96%)	9 (3%)	1 (0%)	34 69
All	All	531/558 (95%)	510 (96%)	20 (4%)	1 (0%)	47 79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	421	LEU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	235/251 (94%)	219 (93%)	16 (7%)	16 45
1	B	233/251 (93%)	220 (94%)	13 (6%)	21 52
All	All	468/502 (93%)	439 (94%)	29 (6%)	18 49

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	271	ASP
1	A	284	MET
1	A	290	ILE
1	A	308	ARG
1	A	315	PHE
1	A	316	ARG
1	A	338	ASP
1	A	397	GLU
1	A	410	SER
1	A	425	ARG
1	A	428	LEU
1	A	429	LEU
1	A	440	ASN
1	A	480	LEU
1	A	488	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	503	ASP
1	B	272	LYS
1	B	284	MET
1	B	285	MET
1	B	308	ARG
1	B	315	PHE
1	B	338	ASP
1	B	364	LYS
1	B	397	GLU
1	B	410	SER
1	B	440	ASN
1	B	457	SER
1	B	480	LEU
1	B	487	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	336	ASN
1	A	373	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	270/279 (96%)	-0.14	5 (1%) 66 46	66, 121, 192, 221	0
1	B	269/279 (96%)	-0.12	3 (1%) 80 64	72, 117, 200, 208	0
All	All	539/558 (96%)	-0.13	8 (1%) 73 54	66, 119, 197, 221	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	288	ASP	4.8
1	A	489	THR	4.4
1	B	300	GLU	4.3
1	A	299	GLN	3.1
1	A	334	PHE	2.6
1	A	309	ILE	2.2
1	A	288	ASP	2.1
1	B	490	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no monosaccharides in this entry.

### 6.4 Ligands i

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.