



Full wwPDB NMR Structure Validation Report ⓘ

Jun 22, 2024 – 07:58 PM EDT

PDB ID : 6G4A
BMRB ID : 34249
Title : FLN5 (full length)
Authors : Waudby, C.A.; Wlodarski, T.; Karyadi, M.-E.; Cassaignau, A.M.E.; Chan, S.H.S.; Wentink, A.S.; Schmidt-Engler, J.M.; Camilloni, C.; Vendruscolo, M.; Cabrita, L.D.; Christodoulou, J.
Deposited on : 2018-03-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 43%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	114	

2 Ensemble composition and analysis

This entry contains 10 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:11-A:62, A:66-A:114 (101)	0.55	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 4, 5, 6, 7, 10
2	8, 9
3	2, 3

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1659 atoms, of which 804 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Gelation factor.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	114	1659	534	804	149	169	3	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P13466
A	2	HIS	-	expression tag	UNP P13466
A	3	HIS	-	expression tag	UNP P13466
A	4	HIS	-	expression tag	UNP P13466
A	5	HIS	-	expression tag	UNP P13466
A	6	HIS	-	expression tag	UNP P13466
A	7	HIS	-	expression tag	UNP P13466
A	8	ALA	-	expression tag	UNP P13466
A	9	SER	-	expression tag	UNP P13466

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Gelation factor




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Gelation factor




4.2.2 Score per residue for model 2

- Molecule 1: Gelation factor




4.2.3 Score per residue for model 3

- Molecule 1: Gelation factor




4.2.4 Score per residue for model 4

- Molecule 1: Gelation factor




4.2.5 Score per residue for model 5

- Molecule 1: Gelation factor




4.2.6 Score per residue for model 6

- Molecule 1: Gelation factor




4.2.7 Score per residue for model 7

- Molecule 1: Gelation factor




4.2.8 Score per residue for model 8

- Molecule 1: Gelation factor




4.2.9 Score per residue for model 9

- Molecule 1: Gelation factor



4.2.10 Score per residue for model 10

- Molecule 1: Gelation factor



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 572 calculated structures, 10 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
GROMACS	structure calculation	4.6.5

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	556
Number of shifts mapped to atoms	556
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	43%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.57±0.02	0±0/763 (0.0± 0.0%)	1.99±0.09	22±4/1043 (2.1± 0.4%)
All	All	0.57	0/7630 (0.0%)	2.00	215/10430 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	5.8±1.4
All	All	0	58

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	98	ARG	NE-CZ-NH1	20.40	130.50	120.30	10	4
1	A	48	ARG	NE-CZ-NH1	15.06	127.83	120.30	6	8
1	A	83	TYR	CB-CG-CD1	-14.23	112.46	121.00	6	2
1	A	48	ARG	NE-CZ-NH2	13.46	127.03	120.30	10	4
1	A	79	TYR	CB-CG-CD1	-13.11	113.13	121.00	7	4
1	A	19	TYR	CB-CG-CD2	-13.04	113.17	121.00	10	4
1	A	91	TYR	CB-CG-CD2	-11.02	114.39	121.00	10	8
1	A	83	TYR	CB-CG-CD2	-10.14	114.92	121.00	10	3
1	A	19	TYR	CB-CG-CD1	9.72	126.83	121.00	10	6
1	A	73	ASP	CB-CG-OD1	8.87	126.28	118.30	7	4
1	A	73	ASP	CB-CG-OD2	-8.82	110.36	118.30	1	5
1	A	91	TYR	CB-CG-CD1	8.67	126.20	121.00	2	3
1	A	81	VAL	CG1-CB-CG2	-8.64	97.08	110.90	6	1
1	A	48	ARG	NH1-CZ-NH2	-8.61	109.93	119.40	10	4
1	A	76	ASP	CB-CG-OD1	8.60	126.04	118.30	9	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	30	ASP	CB-CG-OD2	8.41	125.87	118.30	2	1
1	A	67	VAL	CA-CB-CG1	8.34	123.42	110.90	5	2
1	A	43	THR	CA-CB-CG2	8.28	124.00	112.40	6	1
1	A	55	PHE	CB-CG-CD2	-8.24	115.03	120.80	2	4
1	A	32	ALA	CB-CA-C	8.14	122.31	110.10	6	2
1	A	82	VAL	CA-CB-CG2	8.13	123.09	110.90	5	2
1	A	39	PHE	CB-CG-CD1	-8.10	115.13	120.80	3	2
1	A	36	PHE	CB-CG-CD1	-8.01	115.19	120.80	2	3
1	A	84	ASP	CB-CG-OD2	7.95	125.45	118.30	9	4
1	A	66	VAL	CA-CB-CG1	7.93	122.80	110.90	4	1
1	A	42	ASP	CB-CG-OD1	-7.84	111.24	118.30	6	4
1	A	50	ASP	CB-CG-OD1	7.83	125.35	118.30	1	1
1	A	98	ARG	NE-CZ-NH2	-7.75	116.42	120.30	10	2
1	A	83	TYR	CG-CD1-CE1	-7.75	115.10	121.30	6	2
1	A	41	VAL	CA-CB-CG1	7.58	122.27	110.90	4	5
1	A	23	GLU	OE1-CD-OE2	-7.43	114.38	123.30	4	1
1	A	91	TYR	CG-CD2-CE2	-7.43	115.36	121.30	6	1
1	A	29	PHE	CB-CG-CD1	7.25	125.87	120.80	6	2
1	A	36	PHE	CB-CG-CD2	7.20	125.84	120.80	2	1
1	A	19	TYR	CG-CD1-CE1	-7.16	115.57	121.30	7	1
1	A	19	TYR	CG-CD2-CE2	7.05	126.94	121.30	2	2
1	A	71	VAL	CG1-CB-CG2	-7.01	99.68	110.90	5	3
1	A	96	THR	CA-CB-CG2	7.00	122.20	112.40	6	2
1	A	98	ARG	CD-NE-CZ	6.98	133.37	123.60	8	2
1	A	29	PHE	CB-CG-CD2	-6.93	115.95	120.80	10	2
1	A	55	PHE	CB-CG-CD1	-6.81	116.03	120.80	1	2
1	A	15	ALA	CB-CA-C	6.80	120.30	110.10	10	2
1	A	78	THR	CA-CB-CG2	6.75	121.85	112.40	4	2
1	A	56	GLU	OE1-CD-OE2	-6.73	115.23	123.30	8	2
1	A	21	GLU	OE1-CD-OE2	-6.71	115.25	123.30	1	1
1	A	53	ASP	CB-CG-OD2	6.71	124.34	118.30	1	1
1	A	71	VAL	CA-CB-CG1	6.59	120.79	110.90	3	2
1	A	95	VAL	CA-CB-CG1	6.56	120.73	110.90	7	1
1	A	81	VAL	CA-CB-CG2	6.49	120.63	110.90	6	3
1	A	46	VAL	CA-CB-CG1	6.42	120.53	110.90	8	2
1	A	84	ASP	OD1-CG-OD2	-6.35	111.24	123.30	9	2
1	A	108	ASP	CB-CG-OD2	6.34	124.01	118.30	1	1
1	A	46	VAL	CG1-CB-CG2	-6.34	100.76	110.90	2	3
1	A	87	VAL	CA-CB-CG2	6.33	120.39	110.90	8	3
1	A	88	GLU	OE1-CD-OE2	-6.25	115.79	123.30	1	3
1	A	79	TYR	CB-CG-CD2	6.19	124.72	121.00	9	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	83	TYR	CD1-CE1-CZ	6.16	125.34	119.80	6	2
1	A	30	ASP	CB-CG-OD1	6.12	123.81	118.30	1	1
1	A	82	VAL	CA-CB-CG1	6.09	120.04	110.90	3	4
1	A	112	ILE	CA-CB-CG1	6.07	122.53	111.00	6	1
1	A	112	ILE	CG1-CB-CG2	-6.06	98.06	111.40	6	1
1	A	48	ARG	CD-NE-CZ	6.04	132.05	123.60	6	1
1	A	55	PHE	CG-CD2-CE2	-6.02	114.18	120.80	3	1
1	A	109	VAL	CA-CB-CG1	6.00	119.90	110.90	2	1
1	A	26	VAL	CA-CB-CG1	5.98	119.86	110.90	6	1
1	A	91	TYR	CA-CB-CG	5.94	124.68	113.40	4	1
1	A	19	TYR	CZ-CE2-CD2	-5.92	114.47	119.80	2	1
1	A	69	ALA	O-C-N	-5.92	113.23	122.70	10	1
1	A	48	ARG	C-N-CA	5.91	136.48	121.70	6	2
1	A	42	ASP	CB-CG-OD2	5.89	123.60	118.30	4	2
1	A	79	TYR	CG-CD1-CE1	-5.89	116.59	121.30	5	2
1	A	62	PRO	N-CA-CB	5.88	110.35	103.30	4	2
1	A	110	LYS	O-C-N	5.81	131.99	122.70	3	1
1	A	93	VAL	CA-CB-CG1	5.80	119.60	110.90	9	1
1	A	98	ARG	NH1-CZ-NH2	-5.79	113.03	119.40	10	1
1	A	53	ASP	CB-CG-OD1	5.73	123.45	118.30	3	1
1	A	113	GLU	OE1-CD-OE2	-5.71	116.44	123.30	1	1
1	A	58	ALA	CB-CA-C	5.70	118.65	110.10	8	1
1	A	19	TYR	CD1-CE1-CZ	5.70	124.93	119.80	7	1
1	A	97	LEU	CB-CG-CD1	5.69	120.67	111.00	4	1
1	A	17	HIS	O-C-N	-5.68	113.61	122.70	8	1
1	A	95	VAL	CG1-CB-CG2	-5.65	101.85	110.90	3	1
1	A	41	VAL	CG1-CB-CG2	-5.64	101.88	110.90	3	1
1	A	74	ASN	CA-CB-CG	5.62	125.77	113.40	4	1
1	A	67	VAL	CG1-CB-CG2	-5.61	101.93	110.90	5	2
1	A	79	TYR	CZ-CE2-CD2	-5.57	114.78	119.80	10	1
1	A	84	ASP	CB-CG-OD1	5.54	123.29	118.30	9	1
1	A	86	PRO	N-CD-CG	5.47	111.40	103.20	5	1
1	A	79	TYR	CG-CD2-CE2	-5.46	116.93	121.30	5	1
1	A	91	TYR	CZ-CE2-CD2	5.40	124.66	119.80	9	1
1	A	101	PRO	N-CA-CB	5.40	109.78	103.30	1	1
1	A	109	VAL	CA-CB-CG2	5.38	118.97	110.90	7	1
1	A	18	SER	CB-CA-C	5.35	120.27	110.10	5	1
1	A	42	ASP	C-N-CA	5.34	135.04	121.70	6	2
1	A	50	ASP	CB-CG-OD2	-5.33	113.50	118.30	5	1
1	A	13	PRO	N-CD-CG	5.32	111.18	103.20	10	1
1	A	76	ASP	C-N-CA	5.29	133.41	122.30	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	78	THR	OG1-CB-CG2	-5.27	97.87	110.00	5	1
1	A	57	VAL	CA-CB-CG1	-5.24	103.04	110.90	5	1
1	A	105	MET	CA-CB-CG	-5.24	104.40	113.30	3	1
1	A	68	ASP	CB-CG-OD1	5.23	123.00	118.30	6	1
1	A	15	ALA	N-CA-CB	-5.22	102.80	110.10	8	1
1	A	112	ILE	O-C-N	-5.17	114.43	122.70	1	1
1	A	68	ASP	CB-CG-OD2	5.16	122.94	118.30	1	2
1	A	88	GLU	O-C-N	-5.14	114.45	123.20	1	1
1	A	104	ASN	C-N-CA	5.14	134.55	121.70	8	1
1	A	68	ASP	CB-CA-C	5.11	120.63	110.40	8	1
1	A	15	ALA	O-C-N	-5.08	114.57	122.70	7	1
1	A	82	VAL	O-C-N	5.08	130.82	122.70	2	1
1	A	43	THR	N-CA-CB	5.05	119.89	110.30	4	1
1	A	79	TYR	CD1-CG-CD2	5.01	123.42	117.90	5	1
1	A	113	GLU	O-C-N	-5.01	114.68	123.20	7	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	87	VAL	Peptide	6
1	A	96	THR	Peptide	6
1	A	19	TYR	Sidechain	5
1	A	91	TYR	Sidechain,Mainchain	5
1	A	79	TYR	Sidechain	4
1	A	50	ASP	Peptide	4
1	A	83	TYR	Sidechain	3
1	A	29	PHE	Sidechain	2
1	A	48	ARG	Sidechain	2
1	A	57	VAL	Mainchain	1
1	A	86	PRO	Peptide	1
1	A	89	GLY	Mainchain	1
1	A	36	PHE	Sidechain	1
1	A	58	ALA	Peptide,Mainchain	1
1	A	103	LYS	Mainchain	1
1	A	24	GLY	Mainchain	1
1	A	105	MET	Peptide	1
1	A	55	PHE	Sidechain	1
1	A	59	ILE	Peptide	1
1	A	16	GLU	Peptide	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	17	HIS	Sidechain	1
1	A	32	ALA	Peptide	1
1	A	21	GLU	Peptide	1
1	A	69	ALA	Mainchain	1
1	A	67	VAL	Peptide	1
1	A	92	ASN	Peptide	1
1	A	97	LEU	Peptide,Mainchain	1
1	A	39	PHE	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	747	710	710	1±1
All	All	7470	7100	7100	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:85:ALA:HB2	1:A:91:TYR:CE2	0.52	2.39	8	1
1:A:107:ILE:HD12	1:A:107:ILE:C	0.46	2.30	8	1
1:A:24:GLY:HA3	1:A:36:PHE:CD2	0.44	2.47	2	1
1:A:19:TYR:O	1:A:39:PHE:CD2	0.44	2.70	7	1
1:A:105:MET:HG3	1:A:106:PRO:HA	0.42	1.92	4	1
1:A:91:TYR:CD2	1:A:111:CYS:SG	0.41	3.13	3	1
1:A:25:LEU:HD21	1:A:36:PHE:CE1	0.41	2.50	6	1
1:A:29:PHE:HA	1:A:112:ILE:O	0.41	2.16	3	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/114 (88%)	90±2 (90±2%)	8±2 (8±2%)	2±1 (2±1%)	13	56
All	All	1000/1140 (88%)	899 (90%)	84 (8%)	17 (2%)	13	56

All 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	43	THR	5
1	A	47	ALA	4
1	A	45	GLY	3
1	A	113	GLU	2
1	A	42	ASP	2
1	A	26	VAL	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/91 (88%)	79±1 (99±1%)	1±1 (1±1%)	82	97
All	All	800/910 (88%)	794 (99%)	6 (1%)	82	97

All 4 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	83	TYR	3
1	A	42	ASP	1
1	A	55	PHE	1
1	A	75	ASN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation i

The completeness of assignment taking into account all chemical shift lists is 43% for the well-defined parts and 39% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list*

7.1.1 Bookkeeping i

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	556
Number of shifts mapped to atoms	556
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing i

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	91	-0.18 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	82	-0.09 ± 0.26	None needed (< 0.5 ppm)
$^{13}\text{C}'$	93	0.41 ± 0.17	None needed (< 0.5 ppm)
^{15}N	94	-0.23 ± 0.42	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments i

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 43%, i.e. 538 atoms were assigned a chemical shift out of a possible 1257. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	458/500 (92%)	189/205 (92%)	178/202 (88%)	91/93 (98%)
Sidechain	80/674 (12%)	0/438 (0%)	80/216 (37%)	0/20 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/83 (0%)	0/40 (0%)	0/42 (0%)	0/1 (0%)
Overall	538/1257 (43%)	189/683 (28%)	258/460 (56%)	91/114 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 39%, i.e. 556 atoms were assigned a chemical shift out of a possible 1430. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	474/566 (84%)	196/232 (84%)	184/228 (81%)	94/106 (89%)
Sidechain	82/739 (11%)	0/481 (0%)	82/237 (35%)	0/21 (0%)
Aromatic	0/125 (0%)	0/64 (0%)	0/54 (0%)	0/7 (0%)
Overall	556/1430 (39%)	196/777 (25%)	266/519 (51%)	94/134 (70%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

