



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 05:30 PM EDT

PDB ID : 6MUL
Title : Murine PI3K delta kinsae domain - cpd 1
Authors : Fischmann, T.O.
Deposited on : 2018-10-23
Resolution : 3.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	940	 3% 80% 7% 13%
1	B	940	 3% 81% 6% 13%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 13340 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	822	6640	4256	1128	1201	55	0	2	1
1	B	822	6632	4253	1126	1199	54	0	1	1


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	105	GLY	-	expression tag	UNP Q3UDT3
B	105	GLY	-	expression tag	UNP Q3UDT3

- Molecule 2 is 1-{1-[8-(1-ethyl-5-methyl-1H-pyrazol-4-yl)-9-methyl-9H-purin-6-yl]piperidin-4-yl}-1,3-dihydro-2H-imidazo[4,5-b]pyridin-2-one (three-letter code: K4A) (formula: C₂₃H₂₆N₁₀O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	34	23	10	1	0	0
2	B	1	34	23	10	1	0	0



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.03Å 219.11Å 78.24Å 90.00° 113.62° 90.00°	Depositor
Resolution (Å)	36.52 – 3.09 36.52 – 3.09	Depositor EDS
% Data completeness (in resolution range)	99.4 (36.52-3.09) 99.4 (36.52-3.09)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 3.12Å)	Xtrriage
Refinement program	BUSTER-TNT 2.11.7	Depositor
R, R_{free}	0.248 , 0.292 0.260 , 0.307	Depositor DCC
R_{free} test set	1776 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	71.1	Xtrriage
Anisotropy	0.218	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 34.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.117 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	13340	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K4A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/6783	0.62	3/9151 (0.0%)
1	B	0.50	0/6775	0.62	3/9141 (0.0%)
All	All	0.51	0/13558	0.62	6/18292 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	368	GLU	N-CA-C	-8.96	86.80	111.00
1	B	367	SER	C-N-CA	7.55	140.58	121.70
1	A	368	GLU	N-CA-C	-6.90	92.38	111.00
1	A	367	SER	C-N-CA	6.01	136.72	121.70
1	B	367	SER	CA-C-N	-5.94	104.12	117.20
1	A	366	CYS	N-CA-C	5.22	125.09	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	434	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6640	0	6615	14	0
1	B	6632	0	6612	11	0
2	A	34	0	0	0	0
2	B	34	0	0	0	0
All	All	13340	0	13227	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:THR:O	1:A:475:LEU:O	1.90	0.90
1:B:434:THR:O	1:B:475:LEU:O	2.06	0.72
1:A:434:THR:O	1:A:475:LEU:CB	2.44	0.64
1:A:270:LEU:HD11	1:A:770:SER:HB2	1.82	0.61
1:A:434:THR:O	1:A:475:LEU:HB2	2.06	0.55
1:A:335:ALA:HB3	1:B:126:HIS:CG	2.43	0.53
1:B:752:MET:HG3	1:B:758:PRO:HG2	1.91	0.52
1:A:434:THR:O	1:A:475:LEU:HB3	2.12	0.48
1:B:621:TYR:CE1	1:B:983:ALA:HB2	2.48	0.48
1:A:335:ALA:HB3	1:B:126:HIS:ND1	2.28	0.48
1:A:621:TYR:CE1	1:A:983:ALA:HB2	2.49	0.47
1:B:435:GLY:O	1:B:474:TYR:HA	2.15	0.46
1:A:637:ALA:HB1	1:A:644:GLY:HA2	2.00	0.44
1:A:859:LEU:HD21	1:A:901:ILE:HD11	1.99	0.44
1:B:637:ALA:HB1	1:B:644:GLY:HA2	2.00	0.44
1:A:435:GLY:O	1:A:474:TYR:HA	2.19	0.42
1:A:358:VAL:HG22	1:A:377:PHE:CE1	2.54	0.42
1:B:358:VAL:HG22	1:B:377:PHE:CE1	2.54	0.42
1:B:434:THR:O	1:B:475:LEU:CB	2.68	0.42
1:B:937:ASP:O	1:B:941:VAL:HG23	2.20	0.41
1:A:937:ASP:O	1:A:941:VAL:HG23	2.20	0.41
1:B:387:MET:HE3	1:B:590:CYS:HB3	2.03	0.41
1:A:387:MET:HE3	1:A:590:CYS:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	802/940 (85%)	748 (93%)	48 (6%)	6 (1%)	22	57
1	B	801/940 (85%)	744 (93%)	51 (6%)	6 (1%)	22	57
All	All	1603/1880 (85%)	1492 (93%)	99 (6%)	12 (1%)	19	57

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	434	THR
1	B	535	ARG
1	A	328	ILE
1	A	434	THR
1	A	535	ARG
1	A	755	LYS
1	B	755	LYS
1	A	730	HIS
1	B	331	ARG
1	B	730	HIS
1	A	331	ARG
1	B	328	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	729/827 (88%)	684 (94%)	45 (6%)	18	49
1	B	728/827 (88%)	685 (94%)	43 (6%)	19	50
All	All	1457/1654 (88%)	1369 (94%)	88 (6%)	18	49

All (88) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	LYS
1	A	118	SER
1	A	190	LEU
1	A	229	ARG
1	A	257	CYS
1	A	270	LEU
1	A	275	THR
1	A	316	LEU
1	A	329	GLU
1	A	332	LYS
1	A	340	LYS
1	A	342	VAL
1	A	365	VAL
1	A	423	LEU
1	A	427	ASP
1	A	467	SER
1	A	472	VAL
1	A	477	GLU
1	A	511	LEU
1	A	512	ARG
1	A	517	ARG
1	A	523	LEU
1	A	530	LEU
1	A	548	ARG
1	A	553	THR
1	A	565	LEU
1	A	589	ASP
1	A	631	LYS
1	A	653	SER
1	A	675	SER
1	A	696	ASN
1	A	705	LYS
1	A	714	MET
1	A	753	ASP
1	A	755	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	792	GLN
1	A	851	ASN
1	A	855	LEU
1	A	893	ASP
1	A	895	HIS
1	A	911	ASP
1	A	915	PHE
1	A	930	VAL
1	A	998	LEU
1	A	1009	GLU
1	B	190	LEU
1	B	229	ARG
1	B	257	CYS
1	B	275	THR
1	B	316	LEU
1	B	332	LYS
1	B	334	ASN
1	B	340	LYS
1	B	342	VAL
1	B	366	CYS
1	B	367	SER
1	B	368	GLU
1	B	423	LEU
1	B	427	ASP
1	B	445	VAL
1	B	467	SER
1	B	477	GLU
1	B	511	LEU
1	B	512	ARG
1	B	517	ARG
1	B	523	LEU
1	B	530	LEU
1	B	548	ARG
1	B	553	THR
1	B	565	LEU
1	B	589	ASP
1	B	631	LYS
1	B	653	SER
1	B	675	SER
1	B	696	ASN
1	B	705	LYS
1	B	714	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	753	ASP
1	B	792	GLN
1	B	851	ASN
1	B	855	LEU
1	B	893	ASP
1	B	895	HIS
1	B	911	ASP
1	B	915	PHE
1	B	930	VAL
1	B	937	ASP
1	B	1009	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	792	GLN
1	B	792	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	K4A	A	9001	-	31,39,39	0.84	1 (3%)	40,58,58	1.65	5 (12%)
2	K4A	B	9001	-	31,39,39	0.81	1 (3%)	40,58,58	1.79	8 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	K4A	A	9001	-	-	2/10/24/24	0/6/6/6
2	K4A	B	9001	-	-	2/10/24/24	0/6/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	9001	K4A	C19-N18	3.42	1.46	1.38
2	B	9001	K4A	C19-N18	3.36	1.46	1.38

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	9001	K4A	C26-N18-C19	-7.63	103.71	109.25
2	A	9001	K4A	C26-N18-C19	-7.05	104.13	109.25
2	A	9001	K4A	O27-C19-N18	-3.83	122.55	125.91
2	B	9001	K4A	O27-C19-N18	-3.30	123.01	125.91
2	B	9001	K4A	C33-N29-N30	3.15	123.00	117.08
2	B	9001	K4A	C33-N29-C28	-2.78	125.64	128.97
2	A	9001	K4A	C33-N29-N30	2.56	121.88	117.08
2	B	9001	K4A	C04-C05-N06	2.53	126.03	118.92
2	B	9001	K4A	C21-C26-N18	2.52	110.02	106.94
2	A	9001	K4A	N20-C19-N18	2.51	109.92	106.66
2	A	9001	K4A	C33-N29-C28	-2.29	126.22	128.97
2	B	9001	K4A	C10-N01-C02	2.24	129.60	126.14
2	B	9001	K4A	N20-C19-N18	2.17	109.48	106.66

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	9001	K4A	C34-C33-N29-N30
2	B	9001	K4A	C34-C33-N29-N30
2	A	9001	K4A	C34-C33-N29-C28
2	B	9001	K4A	C34-C33-N29-C28

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	822/940 (87%)	0.05	24 (2%)	51 28	32, 82, 129, 168	0
1	B	822/940 (87%)	0.04	32 (3%)	39 20	25, 78, 134, 165	0
All	All	1644/1880 (87%)	0.05	56 (3%)	45 24	25, 80, 131, 168	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	840	ASN	9.2
1	B	366	CYS	7.2
1	A	317	TRP	5.9
1	B	840	ASN	5.7
1	B	317	TRP	5.2
1	A	843	ASN	4.9
1	B	842	SER	4.8
1	A	842	SER	4.2
1	B	330	GLY	3.8
1	A	197	GLU	3.7
1	A	366	CYS	3.7
1	B	110	LYS	3.7
1	B	109	VAL	3.6
1	B	843	ASN	3.6
1	B	343	VAL	3.5
1	B	344	GLN	3.3
1	A	415	ASP	3.3
1	B	346	GLY	3.2
1	B	345	ALA	3.1
1	A	471	LEU	2.9
1	B	394	LEU	2.9
1	A	192	VAL	2.9
1	A	470	ALA	2.9
1	B	347	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	235	GLN	2.8
1	B	334	ASN	2.8
1	B	342	VAL	2.8
1	B	522	GLU	2.7
1	A	591	TYR	2.7
1	B	360	SER	2.7
1	B	378	ASP	2.7
1	A	511	LEU	2.6
1	A	433	LYS	2.6
1	B	230	GLN	2.6
1	B	377	PHE	2.5
1	A	334	ASN	2.5
1	A	515	LEU	2.5
1	B	432	LEU	2.4
1	A	203	PHE	2.3
1	A	332	LYS	2.3
1	B	371	TRP	2.3
1	B	205	PHE	2.3
1	A	568	LEU	2.2
1	A	1027	TRP	2.2
1	A	187	ASN	2.2
1	A	375	LEU	2.1
1	B	323	PHE	2.1
1	B	362	GLU	2.1
1	B	397	VAL	2.1
1	B	203	PHE	2.1
1	B	229	ARG	2.1
1	B	361	SER	2.0
1	B	197	GLU	2.0
1	B	433	LYS	2.0
1	A	919	PHE	2.0
1	A	209	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	K4A	A	9001	34/34	0.93	0.31	35,45,61,66	0
2	K4A	B	9001	34/34	0.95	0.25	31,42,58,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.