



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:50 PM UTC

PDB ID : 6OEZ / pdb_00006oez
Title : Crystal structure of Trypanothione Reductase from Trypanosoma brucei in complex with inhibitor (+)-N-(Cyclobutylmethyl)-3-{5-[1-(pyrrolidin-1-yl)cyclohexyl]-2-(1-[[{(2S)-pyrrolidin-2-yl]methyl}-1H-indol-5-yl]-1,3-thiazol-4-yl]prop-2-yn-1-amine
Authors : Halgas, O.; De Gasparo, R.; Harangozo, D.; Krauth-Siegel, R.L.; Diederich, F.; Pai, E.F.
Deposited on : 2019-03-28
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	 2% 92% 5%
1	B	495	 2% 96%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Trypanothione reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	486	Total 3692	C 2350	N 627	O 696	S 19	0	0	0
1	A	485	Total 3683	C 2344	N 625	O 695	S 19	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP Q389T8
B	-1	SER	-	expression tag	UNP Q389T8
B	0	HIS	-	expression tag	UNP Q389T8
A	-2	GLY	-	expression tag	UNP Q389T8
A	-1	SER	-	expression tag	UNP Q389T8
A	0	HIS	-	expression tag	UNP Q389T8

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	53	27	9	15	2	0	0
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	B	1	15	8	2	4	1	0	0
3	A	1	15	8	2	4	1	0	0

- Molecule 4 is N-(cyclobutylmethyl)-3-{5-[1-(pyrrolidin-1-yl)cyclohexyl]-2-(1-[[2-(2S)-pyrrolidin-2-yl]methyl]-1H-indol-5-yl)-1,3-thiazol-4-yl}prop-2-yn-1-amine (CCD ID: M9J) (formula: C₃₄H₄₅N₅S).

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	117.53Å 117.53Å 224.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.10 – 2.50 29.10 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.10-2.50) 94.0 (29.10-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.217 , 0.235 0.222 , 0.241	Depositor DCC
R_{free} test set	2724 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 28.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7795	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M9J, FAD, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	2/3761 (0.1%)	0.34	0/5104
1	B	0.17	0/3770	0.35	0/5115
All	All	0.22	2/7531 (0.0%)	0.34	0/10219

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	287	ARG	CZ-NH2	6.17	1.41	1.33
1	A	287	ARG	CZ-NH1	5.63	1.40	1.32

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3683	0	3690	17	1
1	B	3692	0	3703	4	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	15	0	17	1	0
3	B	15	0	17	1	0
4	A	40	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	40	0	0	0	0
5	A	94	0	0	1	0
5	B	110	0	0	1	1
All	All	7795	0	7489	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:MET:HA	4:A:503:M9J:C26	2.24	0.66
4:A:503:M9J:C23	4:A:503:M9J:C18	2.86	0.54
1:A:116:ASP:HB2	4:A:503:M9J:C25	2.38	0.53
1:A:399:LEU:N	3:A:502:EPE:O2S	2.41	0.47
1:A:266:LYS:N	1:A:266:LYS:HD2	2.32	0.45
1:B:99:LYS:HD2	1:B:99:LYS:C	2.43	0.44
1:A:289:PRO:HG3	1:A:330:ASP:HB2	2.00	0.44
1:A:231:ASP:OD2	5:A:601:HOH:O	2.21	0.43
1:A:113:MET:HG3	4:A:503:M9J:C28	2.47	0.43
1:A:113:MET:HA	4:A:503:M9J:C27	2.48	0.43
1:A:341:GLU:HA	1:A:357:THR:HG21	2.00	0.43
1:A:370:PRO:HG2	1:A:430:LEU:HD11	2.00	0.43
1:A:163:TRP:CG	1:A:164:PRO:HD2	2.54	0.42
1:A:219:LEU:O	1:A:250:MET:N	2.48	0.42
1:A:221:TYR:CE2	1:A:223:ASN:HB2	2.54	0.42
1:A:479:VAL:HG22	1:A:484:MET:HE3	2.01	0.42
1:A:164:PRO:HD3	1:A:287:ARG:NH1	2.35	0.42
1:B:351:GLY:O	1:B:352:ASN:C	2.62	0.41
1:B:60:LYS:HD2	1:B:60:LYS:C	2.46	0.41
3:B:502:EPE:O1S	5:B:601:HOH:O	2.22	0.41
1:A:392:TYR:O	1:A:414:LYS:HA	2.21	0.40
1:B:440:ALA:HB3	1:A:440:ALA:HB3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:GLU:OE2	5:B:684:HOH:O[8_554]	1.99	0.21

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	483/495 (98%)	467 (97%)	16 (3%)	0	100	100
1	B	484/495 (98%)	471 (97%)	12 (2%)	1 (0%)	43	63
All	All	967/990 (98%)	938 (97%)	28 (3%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	45	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	398/407 (98%)	394 (99%)	4 (1%)	68	86
1	B	399/407 (98%)	396 (99%)	3 (1%)	73	88
All	All	797/814 (98%)	790 (99%)	7 (1%)	70	87

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	57	CYS
1	B	216	LYS
1	B	335	THR
1	A	57	CYS
1	A	61	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	288	ILE
1	A	335	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	224	ASN
1	B	252	ASN
1	B	340	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	M9J	A	503	-	43,46,46	0.35	0	48,65,65	0.78	2 (4%)
4	M9J	B	503	-	43,46,46	0.35	0	48,65,65	0.77	2 (4%)
3	EPE	B	502	-	15,15,15	0.74	1 (6%)	19,20,20	1.45	4 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EPE	A	502	-	15,15,15	0.73	1 (6%)	19,20,20	1.50	2 (10%)
2	FAD	A	501	-	58,58,58	1.47	10 (17%)	85,89,89	1.57	16 (18%)
2	FAD	B	501	-	58,58,58	1.47	10 (17%)	85,89,89	1.58	15 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	M9J	A	503	-	-	7/20/58/58	0/7/7/7
4	M9J	B	503	-	-	4/20/58/58	0/7/7/7
3	EPE	B	502	-	-	4/9/19/19	0/1/1/1
3	EPE	A	502	-	-	0/9/19/19	0/1/1/1
2	FAD	A	501	-	-	7/34/50/50	0/6/6/6
2	FAD	B	501	-	-	5/34/50/50	0/6/6/6

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	C9A-C5X	5.65	1.50	1.41
2	B	501	FAD	C9A-C5X	5.46	1.49	1.41
2	A	501	FAD	C5A-C4A	4.37	1.46	1.39
2	B	501	FAD	C5A-C4A	4.32	1.46	1.39
2	A	501	FAD	C8-C7	3.41	1.49	1.40
2	B	501	FAD	C8-C7	3.39	1.49	1.40
2	B	501	FAD	C5A-C6A	2.63	1.48	1.41
2	A	501	FAD	C5A-C6A	2.63	1.48	1.41
2	B	501	FAD	C4-N3	-2.53	1.34	1.38
3	A	502	EPE	C10-S	2.48	1.81	1.77
2	A	501	FAD	C4-N3	-2.46	1.34	1.38
2	A	501	FAD	C8A-N7A	2.42	1.36	1.31
3	B	502	EPE	C10-S	2.41	1.81	1.77
2	B	501	FAD	C5A-N7A	-2.36	1.34	1.39
2	B	501	FAD	C8A-N7A	2.35	1.36	1.31
2	A	501	FAD	C5A-N7A	-2.23	1.35	1.39
2	A	501	FAD	C4X-N5	2.16	1.35	1.30
2	A	501	FAD	C4A-N9A	-2.12	1.33	1.37
2	B	501	FAD	C4X-N5	2.11	1.35	1.30
2	B	501	FAD	C4A-N9A	-2.06	1.33	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	FAD	C2-N3	-2.05	1.34	1.39
2	A	501	FAD	C2-N3	-2.02	1.34	1.39

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	C5A-C4A-N3A	-5.63	118.96	126.72
2	A	501	FAD	C5A-C4A-N3A	-5.30	119.42	126.72
2	B	501	FAD	N3A-C4A-N9A	4.46	134.75	127.17
2	A	501	FAD	N3A-C4A-N9A	4.16	134.24	127.17
2	B	501	FAD	C2A-N3A-C4A	3.92	121.40	111.83
2	B	501	FAD	N3A-C2A-N1A	-3.90	122.68	128.58
3	A	502	EPE	C5-N4-C3	3.83	117.09	108.84
2	A	501	FAD	N3A-C2A-N1A	-3.82	122.81	128.58
4	A	503	M9J	C5-C6-N2	3.81	115.78	109.99
4	B	503	M9J	C5-C6-N2	3.78	115.73	109.99
2	A	501	FAD	C2A-N3A-C4A	3.77	121.05	111.83
2	A	501	FAD	C4A-C5A-N7A	-3.62	106.44	110.58
2	B	501	FAD	C4A-C5A-N7A	-3.50	106.58	110.58
3	B	502	EPE	C5-N4-C3	3.28	115.90	108.84
2	A	501	FAD	C4A-N9A-C8A	3.24	109.14	105.74
2	A	501	FAD	C4-C4X-N5	3.17	122.59	118.21
2	B	501	FAD	C4-C4X-N5	3.13	122.53	118.21
2	B	501	FAD	C4A-N9A-C8A	3.08	108.98	105.74
2	A	501	FAD	C5A-N7A-C8A	2.79	107.84	103.45
2	B	501	FAD	C5A-N7A-C8A	2.75	107.78	103.45
2	A	501	FAD	C4X-C10-N10	2.67	120.30	116.48
3	B	502	EPE	C7-N4-C5	2.65	118.31	111.24
3	A	502	EPE	C7-N4-C5	2.61	118.20	111.24
2	A	501	FAD	N9A-C8A-N7A	-2.60	110.24	113.94
2	B	501	FAD	N9A-C8A-N7A	-2.51	110.38	113.94
2	A	501	FAD	C6A-C5A-N7A	2.49	136.89	132.09
2	A	501	FAD	C4X-C10-N1	-2.43	118.62	124.59
2	B	501	FAD	C4X-C10-N10	2.43	119.95	116.48
2	B	501	FAD	C6A-C5A-N7A	2.30	136.53	132.09
2	B	501	FAD	C4X-C10-N1	-2.30	118.95	124.59
3	B	502	EPE	O2S-S-C10	2.27	110.16	106.73
2	B	501	FAD	O4-C4-C4X	-2.23	120.64	126.53
4	B	503	M9J	C7-C6-N2	-2.22	115.77	120.99
4	A	503	M9J	C7-C6-N2	-2.21	115.78	120.99
2	A	501	FAD	O4-C4-C4X	-2.21	120.69	126.53
2	A	501	FAD	C2A-N1A-C6A	2.18	122.31	118.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	EPE	C7-N4-C3	2.17	117.01	111.24
2	A	501	FAD	C10-N1-C2	2.16	121.52	116.85
2	B	501	FAD	C2A-N1A-C6A	2.15	122.25	118.73
2	B	501	FAD	C10-N1-C2	2.05	121.28	116.85
2	A	501	FAD	C4X-C4-N3	2.03	118.43	113.25

There are no chirality outliers.

All (27) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	FAD	O4'-C4'-C5'-O5'
2	A	501	FAD	C5'-O5'-P-O2P
2	A	501	FAD	C5'-O5'-P-O3P
3	B	502	EPE	C9-C10-S-O1S
3	B	502	EPE	C9-C10-S-O2S
4	B	503	M9J	N1-C10-C11-C14
4	B	503	M9J	N1-C10-C11-C12
4	A	503	M9J	C5-C4-N-C3
4	A	503	M9J	C5-C4-N-C
4	A	503	M9J	N3-C20-C21-N4
3	B	502	EPE	C8-C7-N4-C5
2	A	501	FAD	C3'-C4'-C5'-O5'
2	A	501	FAD	O4B-C4B-C5B-O5B
4	A	503	M9J	C29-C4-N-C3
3	B	502	EPE	C9-C10-S-O3S
2	A	501	FAD	C3B-C4B-C5B-O5B
2	B	501	FAD	PA-O3P-P-O5'
2	A	501	FAD	PA-O3P-P-O5'
4	B	503	M9J	N3-C20-C21-C24
4	A	503	M9J	N3-C20-C21-C24
2	B	501	FAD	O4B-C4B-C5B-O5B
2	B	501	FAD	PA-O3P-P-O1P
4	A	503	M9J	C33-C4-N-C3
4	B	503	M9J	C11-C10-N1-C9
4	A	503	M9J	C11-C10-N1-C9
2	B	501	FAD	P-O3P-PA-O1A
2	B	501	FAD	P-O3P-PA-O2A


There are no ring outliers.


3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	M9J	5	0
3	B	502	EPE	1	0
3	A	502	EPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	485/495 (97%)	0.12	9 (1%) 66 62	20, 30, 49, 67	0
1	B	486/495 (98%)	0.03	10 (2%) 63 59	19, 27, 42, 61	0
All	All	971/990 (98%)	0.07	19 (1%) 65 61	19, 28, 48, 67	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4	ALA	4.0
1	A	488	PRO	4.0
1	B	3	LYS	3.9
1	A	352	ASN	3.5
1	B	488	PRO	2.6
1	B	352	ASN	2.6
1	B	353	LYS	2.5
1	A	446	ARG	2.5
1	A	4	ALA	2.4
1	B	399	LEU	2.3
1	B	402	ASN	2.2
1	B	118	GLU	2.1
1	A	487	LEU	2.1
1	A	482	GLU	2.1
1	A	30	ARG	2.1
1	B	420	SER	2.1
1	B	168	ALA	2.1
1	A	351	GLY	2.1
1	A	118	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	M9J	A	503	40/40	0.70	0.26	56,62,65,67	0
4	M9J	B	503	40/40	0.76	0.21	41,45,49,50	0
3	EPE	B	502	15/15	0.88	0.14	26,33,43,43	0
3	EPE	A	502	15/15	0.88	0.17	51,53,58,58	0
2	FAD	A	501	53/53	0.95	0.09	24,29,33,34	0
2	FAD	B	501	53/53	0.96	0.10	27,28,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around M9J B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.