

Full wwPDB X-ray Structure Validation Report (i)

Jun 25, 2024 – 01:49 PM EDT

PDB ID	:	60KN
Title	:	OX40R (TNFRSF4) bound to Fab $1A7$
Authors	:	Ultsch, M.H.; Boenig, G.; Harris, S.F.
Deposited on	:	2019-04-14
Resolution	:	3.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	1191 (3.30-3.22)
Clashscore	141614	1251 (3.30-3.22)
Ramachandran outliers	138981	1229 (3.30-3.22)
Sidechain outliers	138945	1228 (3.30-3.22)
RSRZ outliers	127900	1154 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	225	.% 81%	14%	•••	
1	С	225	78%	16%	•••	
2	В	214	80%	17%	•	
2	D	214	2% 79 %	17%	•	
3	Е	163	14% 39% 10% • 50%			



Mol	Chain	Length		Quality of chain					
			13%						
3	R	163		47%	9%	•	42%		



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7978 atoms, of which 51 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Fab 1A7 heavy chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	217	Total	С	Ν	Ο	\mathbf{S}	0	0	0
1		211	1638	1034	270	328	6	0	0	0
1	C	220	Total	С	Ν	0	S	0	0	0
1		220	1659	1046	274	332	7	0	0	0

• Molecule 2 is a protein called Fab 1A7 light chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
0	Р	012	Total	С	Ν	0	S	0	0	0
	D	213	1647	1029	280	333	5	0		
0	П	012	Total	С	Ν	0	S	0	0	0
	D	213	1647	1029	280	333	5	0	0	0

• Molecule 3 is a protein called Tumor necrosis factor receptor superfamily member 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Е	81	Total 595	C 359	N 111	O 115	S 10	0	0	0
3	R	94	Total 693	C 420	N 127	0 134	S 12	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Е	8	MET	-	initiating methionine	UNP P43489
Е	9	GLY	-	expression tag	UNP P43489
Е	10	SER	-	expression tag	UNP P43489
Е	11	SER	-	expression tag	UNP P43489
Е	12	HIS	-	expression tag	UNP P43489
Е	13	HIS	-	expression tag	UNP P43489
Е	14	HIS	-	expression tag	UNP P43489
Е	15	HIS	-	expression tag	UNP P43489



Chain	Residue	Modelled	Actual	Comment	Reference
Е	16	HIS	-	expression tag	UNP P43489
Е	17	HIS	-	expression tag	UNP P43489
Е	18	SER	-	expression tag	UNP P43489
Е	19	SER	-	expression tag	UNP P43489
Е	20	GLY	-	expression tag	UNP P43489
Е	21	LEU	-	expression tag	UNP P43489
Е	22	VAL	-	expression tag	UNP P43489
Е	23	PRO	-	expression tag	UNP P43489
Е	24	ARG	-	expression tag	UNP P43489
Е	25	GLY	-	expression tag	UNP P43489
Е	26	SER	-	expression tag	UNP P43489
Е	27	HIS	-	expression tag	UNP P43489
Е	28	MET	-	expression tag	UNP P43489
R	8	MET	-	initiating methionine	UNP P43489
R	9	GLY	-	expression tag	UNP P43489
R	10	SER	-	expression tag	UNP P43489
R	11	SER	-	expression tag	UNP P43489
R	12	HIS	-	expression tag	UNP P43489
R	13	HIS	-	expression tag	UNP P43489
R	14	HIS	-	expression tag	UNP P43489
R	15	HIS	-	expression tag	UNP P43489
R	16	HIS	-	expression tag	UNP P43489
R	17	HIS	-	expression tag	UNP P43489
R	18	SER	-	expression tag	UNP P43489
R	19	SER	-	expression tag	UNP P43489
R	20	GLY	-	expression tag	UNP P43489
R	21	LEU	-	expression tag	UNP P43489
R	22	VAL	-	expression tag	UNP P43489
R	23	PRO	-	expression tag	UNP P43489
R	24	ARG	-	expression tag	UNP P43489
R	25	GLY	-	expression tag	UNP P43489
R	26	SER	-	expression tag	UNP P43489
R	27	HIS	-	expression tag	UNP P43489
R	28	MET	-	expression tag	UNP P43489

• Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C H N O S 32 8 17 2 4 1	0	0
4	В	1	Total C H N O S 32 8 17 2 4 1	0	0
4	D	1	Total C H N O S 32 8 17 2 4 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	2	Total O 2 2	0	0
5	С	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Fab 1A7 heavy chain

CYS • Molecule 3: Tumor necrosis factor receptor superfamily member 4 14% Chain E: 39% 10% 50% GLY PRO GLY PHE PHE ASN VAL VAL VAL VAL LYS SER LYS SER LYS SPRO CYS PRO C87 N86 R96 S9 GLU ASP ASP ASP • Molecule 3: Tumor necrosis factor receptor superfamily member 4 13% Chain R: 47% 42% 9% VAL VAL SER SER LYS PRO



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	121.59Å 121.59Å 198.08Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{Posolution} \left(\overset{\circ}{\mathbf{A}} \right)$	29.49 - 3.25	Depositor
Resolution (A)	29.49 - 3.25	EDS
% Data completeness	99.8 (29.49-3.25)	Depositor
(in resolution range)	99.9 (29.49-3.25)	EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.88 (at 3.24 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
P. P.	0.192 , 0.228	Depositor
n, n_{free}	0.211 , 0.245	DCC
R_{free} test set	1213 reflections (5.05%)	wwPDB-VP
Wilson B-factor $(Å^2)$	98.6	Xtriage
Anisotropy	0.474	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.29 , 83.1	EDS
L-test for twinning ²	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7978	wwPDB-VP
Average B, all atoms $(Å^2)$	120.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 14.85% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.40	0/1679	0.71	0/2292
1	С	0.41	0/1700	0.72	0/2319
2	В	0.37	0/1683	0.63	0/2285
2	D	0.37	0/1683	0.63	0/2285
3	Е	0.45	0/609	0.66	0/829
3	R	0.44	0/711	0.64	0/968
All	All	0.40	0/8065	0.67	0/10978

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1638	0	1598	13	0
1	С	1659	0	1621	24	0
2	В	1647	0	1604	19	0
2	D	1647	0	1604	23	0
3	Е	595	0	550	4	0
3	R	693	0	634	6	0
4	А	15	17	17	0	0



	3	1	1 5			
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	В	15	17	17	1	0
4	D	15	17	17	0	0
5	А	2	0	0	0	0
5	С	1	0	0	0	0
All	All	7927	51	7662	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1 Atom-2		distance (Å)	overlap (Å)
2:B:123:GLU:N	4:B:301:EPE:O2S	2.15	0.78
1:C:11:VAL:HB	1:C:151:PRO:HG3	1.74	0.69
3:R:155:LEU:HB2	3:R:166:CYS:HA	1.74	0.69
1:A:11:VAL:HB	1:A:151:PRO:HG3	1.75	0.69
3:R:97:GLN:HB2	3:R:106:VAL:HB	1.74	0.69
1:C:102:TRP:CD1	2:D:91:GLY:HA2	2.28	0.68
1:C:133:LYS:HG2	2:D:117:ILE:HG22	1.76	0.68
1:C:51:MET:HG2	1:C:57:ASP:HB2	1.77	0.67
2:D:2:ILE:HG23	2:D:26:SER:HB3	1.78	0.66
2:D:37:GLN:HB2	2:D:47:LEU:HD11	1.78	0.65
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.79	0.64
1:C:102:TRP:CD1	2:D:96:PRO:HB3	2.34	0.63
1:C:91:THR:HG23	1:C:114:THR:HA	1.81	0.63
2:B:113:PRO:HD2	2:B:201:LEU:HD13	1.80	0.63
2:D:113:PRO:HD2	2:D:201:LEU:HD13	1.81	0.62
1:C:101:ARG:HD2	1:C:102:TRP:CZ3	2.35	0.61
1:A:91:THR:HG23	1:A:114:THR:HA	1.82	0.60
1:C:102:TRP:HD1	2:D:91:GLY:HA2	1.70	0.56
2:D:2:ILE:HG12	2:D:27:GLN:HB2	1.88	0.55
3:E:146:ASN:HD21	3:E:149:LEU:HB2	1.72	0.52
2:D:155:GLN:HB3	2:D:158:ASN:HD21	1.75	0.52
1:C:30:THR:HA	1:C:53:PRO:HB2	1.91	0.52
2:D:14:SER:O	2:D:17:ASP:HB2	2.10	0.52
2:D:22:THR:HG22	2:D:72:THR:HG23	1.92	0.52
2:D:197:THR:HG23	2:D:204:PRO:HG3	1.91	0.52
2:B:197:THR:HG23	2:B:204:PRO:HG3	1.92	0.52
2:D:21:ILE:HG12	2:D:102:THR:HG21	1.93	0.51
1:C:142:LEU:HD13	1:C:215:VAL:HG21	1.94	0.50
2:B:155:GLN:HB3	2:B:158:ASN:HD21	1.75	0.50



	lo uo pugom	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:B:21:ILE:HG12	2:B:102:THR:HG21	1.93	0.50
2:B:90:GLN:HG2	2:B:92:HIS:H	1.77	0.50
1:A:142:LEU:HD13	1:A:215:VAL:HG21	1.94	0.48
1:C:40:ALA:HB3	1:C:43:GLN:HB2	1.96	0.48
2:D:120:PRO:HD3	2:D:132:VAL:HG22	1.97	0.47
2:B:48:ILE:HD13	2:B:54:LEU:HB3	1.96	0.47
1:C:146:VAL:HG11	1:C:154:VAL:HG11	1.97	0.47
1:C:133:LYS:HG2	2:D:117:ILE:CG2	2.43	0.47
2:B:34:ASN:HD22	2:B:89:GLN:HE21	1.62	0.46
1:A:134:SER:HA	2:B:116:PHE:HD1	1.80	0.46
3:R:99:CYS:HB2	3:R:105:THR:HG23	1.98	0.46
2:B:120:PRO:HD3	2:B:132:VAL:HG22	1.97	0.46
2:D:33:LEU:HD21	2:D:88:CYS:HB2	1.98	0.46
1:C:19:LYS:HG3	1:C:82:GLU:HB3	1.97	0.46
2:B:47:LEU:HA	2:B:58:VAL:HG21	1.97	0.45
1:A:146:VAL:HG11	1:A:154:VAL:HG11	1.98	0.45
2:D:47:LEU:HA	2:D:58:VAL:HG21	1.98	0.45
2:B:3:GLN:HG3	2:B:26:SER:HB2	1.98	0.45
2:D:5:THR:HA	2:D:100:GLN:HE22	1.80	0.45
1:C:29:PHE:O	1:C:53:PRO:HG2	2.17	0.45
1:A:7:SER:HB3	1:A:21:SER:H	1.82	0.44
2:B:36:TYR:HE1	2:B:89:GLN:CG	2.31	0.44
1:C:2:VAL:HG23	3:E:119:TYR:HB3	1.97	0.44
2:D:36:TYR:HE1	2:D:89:GLN:CG	2.30	0.44
1:C:2:VAL:HG22	1:C:27:TYR:HB3	1.99	0.44
1:C:101:ARG:HD2	1:C:102:TRP:HZ3	1.81	0.44
1:A:2:VAL:HG22	1:A:27:TYR:HB3	2.00	0.44
2:B:34:ASN:HD22	2:B:89:GLN:NE2	2.16	0.44
1:A:41:PRO:HD3	1:A:92:ALA:HA	2.00	0.44
2:B:90:GLN:HE21	2:B:93:THR:H	1.64	0.43
2:D:146:VAL:HG11	2:D:177:SER:HB2	2.00	0.43
2:B:52:SER:HA	2:B:64:GLY:HA3	2.00	0.43
1:C:60:TYR:HB2	1:C:65:ARG:HG3	2.00	0.43
3:R:96:LYS:HD2	3:R:108:ARG:HD3	2.01	0.43
2:B:33:LEU:HD21	2:B:88:CYS:HB2	2.00	0.43
1:C:7:SER:HB3	1:C:21:SER:H	1.82	0.43
1:A:101:ARG:HA	1:A:102:TRP:HA	1.84	0.42
2:D:18:ARG:HA	2:D:75:ILE:O	2.20	0.42
1:A:170:PHE:HB3	2:B:162:SER:OG	2.19	0.42
3:R:100:THR:HB	3:R:103:GLN:H	1.85	0.42
2:D:17:ASP:O	2:D:18:ARG:HB3	2.19	0.41



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:154:THR:HA	3:E:166:CYS:HB3	2.02	0.41
1:A:72:ARG:HD3	1:A:74:THR:HG23	2.02	0.41
1:A:60:TYR:HB2	1:A:65:ARG:HG3	2.01	0.41
1:C:134:SER:HA	2:D:116:PHE:HD1	1.86	0.41
3:R:155:LEU:CB	3:R:166:CYS:HA	2.45	0.41
1:C:131:SER:C	1:C:133:LYS:H	2.24	0.40
1:C:218:LYS:HE3	1:C:219:SER:H	1.86	0.40
3:E:158:ALA:HB2	3:E:164:ALA:N	2.37	0.40
1:C:99:ALA:HA	1:C:100:PRO:HD2	1.91	0.40
1:A:30:THR:HG23	1:A:54:ASP:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Per	centiles
1	А	215/225~(96%)	192 (89%)	21 (10%)	2(1%)	1'	7 50
1	С	218/225~(97%)	196 (90%)	15~(7%)	7 (3%)	4	23
2	В	211/214~(99%)	192 (91%)	15 (7%)	4 (2%)	8	34
2	D	211/214~(99%)	188 (89%)	18 (8%)	5(2%)	6	28
3	Е	77/163~(47%)	63 (82%)	11 (14%)	3 (4%)	3	18
3	R	90/163~(55%)	78 (87%)	10 (11%)	2 (2%)	6	31
All	All	1022/1204 (85%)	909 (89%)	90 (9%)	23~(2%)	6	29

All (23) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	219	SER
3	R	123	VAL



Mol	Chain	Res	Type
2	В	30	SER
2	В	138	ASN
2	D	18	ARG
2	D	30	SER
2	D	138	ASN
1	А	65	ARG
1	С	58	SER
1	С	136	SER
1	А	105	SER
2	В	84	ALA
1	С	65	ARG
1	С	100	PRO
1	С	101	ARG
1	С	105	SER
3	Е	88	ASN
3	R	158	ALA
2	В	211	ARG
2	D	50	TYR
2	D	211	ARG
3	Е	163	ASP
3	Е	164	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	187/195~(96%)	175~(94%)	12 (6%)	17 47
1	С	190/195~(97%)	173 (91%)	17 (9%)	9 32
2	В	189/190~(100%)	177~(94%)	12 (6%)	18 47
2	D	189/190~(100%)	175~(93%)	14 (7%)	13 40
3	Ε	67/142~(47%)	59~(88%)	8 (12%)	5 20
3	R	78/142~(55%)	71 (91%)	7 (9%)	9 32
All	All	900/1054~(85%)	830~(92%)	70 (8%)	12 38



All ((70)	residues	with a	non-rotameric	sidechain	are listed	below:
-------	------	----------	--------	---------------	-----------	------------	--------

Mol	Chain	Res	Type
1	А	30	THR
1	А	50	ASP
1	А	55	ASN
1	А	67	ARG
1	А	97	VAL
1	А	109	GLN
1	А	132	SER
1	А	152	GLU
1	А	154	VAL
1	А	174	LEU
1	А	182	LEU
1	А	201	ASN
2	В	1	ASP
2	В	7	SER
2	В	33	LEU
2	В	54	LEU
2	В	90	GLN
2	В	137	ASN
2	В	142	ARG
2	В	152	ASN
2	В	161	GLU
2	В	180	THR
2	В	197	THR
2	В	201	LEU
1	С	31	ASP
1	С	34	MET
1	С	50	ASP
1	С	58	SER
1	С	69	THR
1	С	72	ARG
1	С	101	ARG
1	С	102	TRP
1	С	131	SER
1	С	132	SER
1	С	133	LYS
1	С	152	GLU
1	С	154	VAL
1	С	174	LEU
1	С	182	LEU
1	С	201	ASN
1	С	218	LYS
2	D	7	SER



Mol	Chain	Res	Type
2	D	17	ASP
2	D	33	LEU
2	D	51	THR
2	D	55	ARG
2	D	72	THR
2	D	89	GLN
2	D	100	GLN
2	D	137	ASN
2	D	152	ASN
2	D	161	GLU
2	D	180	THR
2	D	197	THR
2	D	201	LEU
3	Е	84	CYS
3	Е	87	CYS
3	Е	104	ASP
3	Е	116	LEU
3	Е	124	ASP
3	Е	145	THR
3	Е	155	LEU
3	Е	162	SER
3	R	107	CYS
3	R	108	ARG
3	R	116	LEU
3	R	117	ASP
3	R	125	CYS
3	R	155	LEU
3	R	156	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	175	GLN
1	А	203	ASN
2	В	89	GLN
2	В	137	ASN
1	С	203	ASN
2	D	100	GLN
2	D	124	GLN
2	D	137	ASN
3	Е	97	GLN
3	Е	138	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Dog	Tink	Bo	ond leng	$_{\rm sths}$	B	ond ang	gles
WIOI	туре	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	D	301	-	15,15,15	0.90	1 (6%)	18,20,20	0.30	0
4	EPE	А	301	-	15,15,15	1.25	1 (6%)	18,20,20	0.46	0
4	EPE	В	301	-	15,15,15	0.90	1 (6%)	18,20,20	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	D	301	-	-	6/9/19/19	0/1/1/1
4	EPE	А	301	-	-	1/9/19/19	0/1/1/1
4	EPE	В	301	-	-	1/9/19/19	0/1/1/1

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
4	А	301	EPE	C10-S	-4.72	1.70	1.77
4	D	301	EPE	C10-S	-3.39	1.72	1.77
4	В	301	EPE	C10-S	-3.19	1.73	1.77

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	301	EPE	S-C10-C9-N1
4	В	301	EPE	N4-C7-C8-O8
4	D	301	EPE	N4-C7-C8-O8
4	D	301	EPE	C9-C10-S-O3S
4	D	301	EPE	C10-C9-N1-C2
4	D	301	EPE	C10-C9-N1-C6
4	D	301	EPE	C9-C10-S-O1S
4	D	301	EPE	C9-C10-S-O2S

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	301	EPE	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	217/225~(96%)	-0.18	2 (0%) 84 84	67, 90, 131, 148	0
1	С	220/225~(97%)	0.03	3 (1%) 75 74	71, 118, 176, 185	0
2	В	213/214~(99%)	-0.18	0 100 100	74, 103, 129, 156	0
2	D	213/214~(99%)	0.08	5 (2%) 60 58	89, 127, 164, 181	0
3	E	81/163~(49%)	1.30	23 (28%) 0 0	126, 157, 227, 245	0
3	R	94/163~(57%)	0.88	21 (22%) 0 1	109, 139, 169, 183	0
All	All	1038/1204 (86%)	0.13	54 (5%) 27 25	67, 115, 174, 245	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Е	85	THR	7.5
3	R	81	CYS	7.4
3	Е	86	TRP	5.3
3	Е	97	GLN	4.6
3	Е	107	CYS	4.5
3	R	74	ASP	4.4
3	R	102	THR	4.3
2	D	68	GLY	4.3
3	Е	84	CYS	4.2
3	Е	119	TYR	4.2
3	Е	104	ASP	4.1
3	Е	148	THR	4.1
3	R	103	GLN	4.0
3	Е	122	GLY	3.7
3	R	148	THR	3.7
3	R	151	GLY	3.5
3	R	147	CYS	3.5
3	Е	150	ALA	3.4
3	Е	106	VAL	3.4



Mol	Chain	Res	Type	RSRZ
2	D	27	GLN	3.3
3	R	154	THR	3.1
3	R	73	ASN	3.0
3	Е	151	GLY	3.0
3	R	83	PRO	2.9
3	R	150	ALA	2.9
3	Е	99	CYS	2.9
3	R	82	LYS	2.9
3	Е	101	ALA	2.7
1	А	42	GLY	2.7
3	R	166	CYS	2.7
1	С	35	SER	2.6
3	Е	156	GLN	2.6
3	Е	100	THR	2.6
3	Е	165	ILE	2.6
2	D	32	TYR	2.5
3	R	67	CYS	2.5
3	R	96	LYS	2.5
3	R	70	GLY	2.4
3	R	97	GLN	2.4
3	R	156	GLN	2.3
2	D	91	GLY	2.3
1	С	65	ARG	2.3
2	D	28	ASP	2.2
3	Е	92	GLY	2.2
3	R	101	ALA	2.2
3	Е	96	LYS	2.2
3	Е	137	ASP	2.2
3	Е	108	ARG	2.2
3	R	90	ARG	2.1
3	Е	123	VAL	2.1
1	А	41	PRO	2.1
3	Е	90	ARG	2.0
1	С	49	GLY	2.0
3	R	92	GLY	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	EPE	А	301	15/15	0.86	0.46	168,172,173,173	0
4	EPE	D	301	15/15	0.86	0.24	131,133,135,136	0
4	EPE	В	301	15/15	0.91	0.23	113,136,139,139	0

6.5 Other polymers (i)

There are no such residues in this entry.

