



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2024 – 12:18 AM EST

PDB ID : 6P3W
Title : Crystal structure of the Cyclin A-CDK2-ORC1 complex
Authors : Wang, B.; Song, J.
Deposited on : 2019-05-24
Resolution : 2.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">3% 88% 12%</p>
1	C	298	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">7% 83% 17% .</p>
2	B	257	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 89%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">7% 89% 10% .</p>
2	D	257	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 91%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">6% 91% 9%</p>
3	E	7	<div style="display: flex; align-items: center;"> <div style="width: 14%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 71%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 29%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">14% 71% 29%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	7	<div style="text-align: center;"> <p>57% 57% 43%</p> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	303	-	-	-	X
4	PO4	C	301	-	-	-	X
4	PO4	C	303	-	-	-	X
4	PO4	D	502	-	-	-	X
4	PO4	D	503	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8967 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	Total 2357	C 1533	N 401	O 414	S 9	1	1	0
1	C	296	Total 2303	C 1495	N 393	O 407	S 8	0	1	0

- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	257	Total 2051	C 1331	N 329	O 380	S 11	0	0	0
2	D	256	Total 2049	C 1329	N 333	O 376	S 11	0	0	0

- Molecule 3 is a protein called ORC1 Peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	7	Total 69	C 43	N 17	O 9	0	1	0
3	F	4	Total 28	C 20	N 4	O 4	0	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	26	Total 26	O 26	0	0
5	B	5	Total 5	O 5	0	0
5	C	4	Total 4	O 4	0	0
5	D	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent kinase 2



- Molecule 1: Cyclin-dependent kinase 2



- Molecule 2: Cyclin-A2



- Molecule 2: Cyclin-A2






- Molecule 3: ORC1 Peptide



- Molecule 3: ORC1 Peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	186.40Å 186.40Å 214.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.07 – 2.54 20.07 – 2.54	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.07-2.54) 99.8 (20.07-2.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.53Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.198 , 0.237 0.205 , 0.239	Depositor DCC
R_{free} test set	3709 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	72.0	Xtrriage
Anisotropy	0.280	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 60.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8967	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2418	0.41	0/3288
1	C	0.27	0/2362	0.46	0/3218
2	B	0.22	0/2101	0.39	0/2859
2	D	0.22	0/2098	0.38	0/2851
3	E	0.24	0/71	0.50	0/91
3	F	0.25	0/27	0.51	0/35
All	All	0.24	0/9077	0.41	0/12342

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2357	0	2364	22	0
1	C	2303	0	2276	28	0
2	B	2051	0	2039	20	0
2	D	2049	0	2052	14	0
3	E	69	0	84	3	0
3	F	28	0	29	0	0
4	A	15	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	5	0	0	0	0
4	C	20	0	0	0	0
4	D	20	0	0	1	0
5	A	26	0	0	1	0
5	B	5	0	0	0	0
5	C	4	0	0	0	0
5	D	15	0	0	0	0
All	All	8967	0	8844	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:39:THR:O	2:D:292:LEU:HD12	1.60	1.00
1:C:39:THR:O	2:D:292:LEU:CD1	2.34	0.76
2:D:263:LEU:HD21	2:D:295:GLU:HG3	1.76	0.67
1:C:248:PHE:HA	1:C:251:VAL:HG22	1.80	0.62
1:C:72:THR:HG23	1:C:75:LYS:H	1.68	0.59
2:B:210:MET:HG2	3:E:239:LEU:HB3	1.85	0.59
1:C:72:THR:OG1	1:C:73:GLU:N	2.36	0.58
1:A:157:ARG:NH2	4:A:301:PO4:O4	2.38	0.57
1:C:284:PRO:HA	1:C:287:GLN:HG3	1.88	0.56
1:A:122:ARG:HG2	1:A:122:ARG:O	2.05	0.55
1:C:160:THR:O	1:C:163:VAL:HG22	2.07	0.55
2:D:391:LEU:HD23	2:D:432:LEU:HD21	1.89	0.55
1:C:124:LEU:HG	1:C:152:PHE:CD1	2.42	0.54
1:C:104:ILE:HG12	1:C:196:MET:HG2	1.88	0.54
2:B:211:ARG:NH2	2:B:341:LEU:O	2.41	0.54
1:A:24:LYS:NZ	4:D:502:PO4:O1	2.41	0.52
2:D:364:LEU:HD21	2:D:370:GLN:HB2	1.91	0.52
2:B:338:GLU:HA	2:B:341:LEU:HD12	1.92	0.52
1:A:72:THR:HG23	1:A:75:LYS:H	1.74	0.52
1:C:231:THR:HA	1:C:236:TYR:CD1	2.46	0.51
2:B:361:HIS:CD2	2:B:391:LEU:HD21	2.46	0.51
1:A:195:GLU:HB2	5:A:410:HOH:O	2.12	0.50
1:A:227:TRP:O	1:A:230:VAL:HG23	2.11	0.50
2:B:280:TYR:OH	3:E:234[B]:ARG:NH2	2.45	0.50
1:A:72:THR:OG1	1:A:73:GLU:N	2.46	0.49
1:A:124:LEU:HG	1:A:152:PHE:CD1	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:158:THR:HG21	1:C:177:CYS:O	2.13	0.48
1:C:161:HIS:HB2	2:D:271:TYR:OH	2.13	0.48
1:C:97:THR:O	1:C:199:ARG:NH2	2.47	0.48
1:A:139:GLY:HA2	1:A:294:PRO:HD3	1.96	0.47
1:C:115:LEU:HD21	1:C:185:ASP:HB3	1.95	0.47
1:A:253:PRO:HG2	1:A:254:PRO:HD3	1.97	0.47
2:B:275:VAL:HG21	2:B:292:LEU:HD21	1.96	0.46
2:D:275:VAL:HG21	2:D:292:LEU:HD21	1.98	0.46
2:B:289:LYS:O	2:B:293:ARG:HG3	2.15	0.46
1:C:241:PRO:HG2	1:C:243:TRP:CZ3	2.51	0.46
1:C:133:LEU:HB3	1:C:141:ILE:HD11	1.96	0.46
2:D:311:VAL:HG12	2:D:352:PRO:HA	1.98	0.46
1:A:231:THR:HA	1:A:236:TYR:CD1	2.51	0.45
1:A:247:ASP:OD2	1:A:249:SER:OG	2.28	0.45
2:B:270:ILE:HG13	2:B:271:TYR:N	2.31	0.45
2:B:211:ARG:O	2:B:215:VAL:HG12	2.17	0.45
1:A:270:ASP:HB3	1:A:273:LYS:HG3	1.99	0.45
2:B:348:LEU:HD23	2:B:348:LEU:HA	1.84	0.45
1:C:203:PHE:CE1	1:C:215:ILE:HA	2.52	0.45
1:C:256:ASP:OD1	1:C:257:GLU:N	2.51	0.44
1:A:94:SER:O	1:A:98:GLY:N	2.45	0.44
1:A:56:LYS:NZ	2:B:303:THR:O	2.50	0.44
2:B:270:ILE:HG13	2:B:271:TYR:H	1.84	0.43
2:D:207:THR:OG1	2:D:208:ASN:N	2.52	0.43
1:C:88:LYS:HA	1:C:91:MET:HE2	2.01	0.43
1:C:105:LYS:HD3	1:C:285:PHE:CE2	2.53	0.43
1:A:40:GLU:O	1:A:42:GLU:HG2	2.19	0.43
1:C:62:ASN:HA	1:C:142:LYS:HG2	2.01	0.42
2:D:364:LEU:CD2	2:D:370:GLN:HB2	2.49	0.42
2:B:280:TYR:CZ	3:E:234[B]:ARG:HD3	2.54	0.42
2:D:217:TRP:HZ2	2:D:281:ILE:HG22	1.84	0.42
1:C:253:PRO:HG2	1:C:254:PRO:HD3	2.00	0.42
1:C:217:ARG:HG2	1:C:243:TRP:CE2	2.53	0.42
1:C:22:ARG:HG3	1:C:29:VAL:HG12	2.01	0.42
2:B:215:VAL:HG13	2:B:342:ILE:HD13	2.00	0.42
2:D:388:LYS:HB3	2:D:389:PRO:HD3	2.01	0.41
2:B:176:PRO:HB2	2:B:177:ASP:H	1.70	0.41
2:B:346:PRO:O	2:B:349:LYS:HG2	2.20	0.41
2:B:218:LEU:HA	2:B:221:VAL:HG22	2.03	0.41
2:B:308:ALA:HA	2:B:309:PRO:HD3	1.89	0.41
1:C:127:ASP:OD1	1:C:129:LYS:HD3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:197:VAL:HG11	1:C:255:LEU:HD13	2.02	0.41
2:D:345:ASP:HA	2:D:346:PRO:HA	1.88	0.41
1:A:10:ILE:HD12	1:A:82:PHE:HE1	1.85	0.40
1:A:197:VAL:HG11	1:A:255:LEU:HD13	2.03	0.40
1:A:237:LYS:HA	1:A:238:PRO:HD3	1.97	0.40
1:C:158:THR:HG22	1:C:178:LYS:O	2.21	0.40
1:A:65:LYS:HE2	1:A:65:LYS:HB2	1.92	0.40
1:A:240:PHE:HA	1:A:241:PRO:HD2	1.94	0.40
1:A:73:GLU:OE2	2:B:293:ARG:NH2	2.54	0.40
2:B:345:ASP:HB3	2:B:346:PRO:HD3	2.03	0.40
1:C:266:MET:HA	1:C:275:ILE:HG22	2.03	0.40
2:D:308:ALA:HA	2:D:309:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/298 (99%)	289 (98%)	7 (2%)	0	100	100
1	C	295/298 (99%)	287 (97%)	8 (3%)	0	100	100
2	B	255/257 (99%)	251 (98%)	3 (1%)	1 (0%)	34	46
2	D	254/257 (99%)	252 (99%)	1 (0%)	1 (0%)	34	46
3	E	6/7 (86%)	6 (100%)	0	0	100	100
3	F	2/7 (29%)	2 (100%)	0	0	100	100
All	All	1108/1124 (99%)	1087 (98%)	19 (2%)	2 (0%)	47	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	324	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	346	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/263 (95%)	248 (99%)	2 (1%)	81	88
1	C	240/263 (91%)	239 (100%)	1 (0%)	91	95
2	B	224/231 (97%)	222 (99%)	2 (1%)	78	86
2	D	224/231 (97%)	222 (99%)	2 (1%)	78	86
3	E	7/6 (117%)	7 (100%)	0	100	100
3	F	2/6 (33%)	2 (100%)	0	100	100
All	All	947/1000 (95%)	940 (99%)	7 (1%)	84	90

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ASP
1	A	252	VAL
2	B	289	LYS
2	B	398	TYR
1	C	84	HIS
2	D	296	HIS
2	D	432	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	D	501	-	4,4,4	0.93	0	6,6,6	0.43	0
4	PO4	C	301	-	4,4,4	0.92	0	6,6,6	0.43	0
4	PO4	D	502	-	4,4,4	0.91	0	6,6,6	0.42	0
4	PO4	A	303	-	4,4,4	0.94	0	6,6,6	0.40	0
4	PO4	B	501	-	4,4,4	0.92	0	6,6,6	0.42	0
4	PO4	D	503	-	4,4,4	0.93	0	6,6,6	0.43	0
4	PO4	A	301	-	4,4,4	0.93	0	6,6,6	0.43	0
4	PO4	A	302	-	4,4,4	0.92	0	6,6,6	0.43	0
4	PO4	C	302	-	4,4,4	0.97	0	6,6,6	0.35	0
4	PO4	C	303	-	4,4,4	0.93	0	6,6,6	0.46	0
4	PO4	C	304	-	4,4,4	0.93	0	6,6,6	0.44	0
4	PO4	D	504	-	4,4,4	0.92	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	502	PO4	1	0
4	A	301	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/298 (100%)	-0.05	8 (2%) 54 61	55, 75, 112, 178	0
1	C	296/298 (99%)	0.26	20 (6%) 17 20	61, 93, 128, 158	0
2	B	257/257 (100%)	0.08	17 (6%) 18 21	62, 83, 115, 159	0
2	D	256/257 (99%)	0.02	15 (5%) 22 26	59, 77, 119, 160	0
3	E	7/7 (100%)	1.72	1 (14%) 2 3	110, 116, 122, 130	0
3	F	4/7 (57%)	3.19	4 (100%) 0 0	118, 125, 128, 148	0
All	All	1118/1124 (99%)	0.10	65 (5%) 23 27	55, 82, 121, 178	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	298	LEU	5.8
1	C	161	HIS	5.6
2	D	323	GLN	5.6
2	B	176	PRO	5.3
1	A	39	THR	5.0
1	C	14	THR	4.6
3	E	234[A]	ARG	4.3
1	C	108	LEU	4.2
1	C	13	GLY	4.1
3	F	238	GLU	4.1
1	C	12	GLU	4.1
1	A	12	GLU	3.8
2	B	323	GLN	3.8
2	B	359	ALA	3.7
1	A	13	GLY	3.5
1	A	108	LEU	3.5
1	C	193	PHE	3.4
1	A	40	GLU	3.4
1	C	15	TYR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	239	LEU	3.2
1	C	293	VAL	3.2
3	F	236	ARG	3.1
2	D	360	PHE	3.0
2	D	359	ALA	3.0
2	D	181	ASP	3.0
2	D	271	TYR	3.0
2	B	362	LEU	2.9
2	B	257	GLY	2.9
1	C	290	THR	2.9
1	A	161	HIS	2.9
2	B	358	ALA	2.8
1	C	95	ALA	2.8
2	B	177	ASP	2.8
1	C	196	MET	2.7
2	D	358	ALA	2.7
2	B	363	ALA	2.6
2	D	257	GLY	2.6
2	B	256	VAL	2.6
2	B	255	LEU	2.5
2	B	324	PRO	2.5
1	C	287	GLN	2.5
2	B	432	LEU	2.4
1	C	294	PRO	2.4
3	F	237	LEU	2.4
2	D	256	VAL	2.4
2	D	324	PRO	2.3
1	C	97	THR	2.3
1	A	171	PRO	2.2
2	D	177	ASP	2.2
1	C	17	VAL	2.2
1	C	96	LEU	2.2
1	C	296	LEU	2.2
2	D	363	ALA	2.1
1	C	104	ILE	2.1
1	C	39	THR	2.1
2	D	325	ALA	2.1
2	B	336	LEU	2.1
2	D	180	GLU	2.1
2	B	325	ALA	2.1
2	D	378	ARG	2.1
2	B	357	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	360	PHE	2.0
2	B	345	ASP	2.0
2	D	357	GLY	2.0
1	C	107	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PO4	C	302	5/5	0.54	0.38	161,165,169,171	0
4	PO4	C	301	5/5	0.66	0.42	217,219,221,222	0
4	PO4	D	501	5/5	0.67	0.33	192,194,195,196	0
4	PO4	D	504	5/5	0.68	0.31	213,214,215,216	0
4	PO4	A	303	5/5	0.71	0.55	190,191,192,193	0
4	PO4	D	503	5/5	0.75	0.42	197,197,202,202	0
4	PO4	C	304	5/5	0.77	0.36	204,205,209,214	0
4	PO4	C	303	5/5	0.78	0.56	194,196,198,201	0
4	PO4	D	502	5/5	0.79	0.45	198,201,203,204	0
4	PO4	A	301	5/5	0.84	0.31	187,188,191,191	0
4	PO4	A	302	5/5	0.86	0.39	170,171,171,175	0
4	PO4	B	501	5/5	0.86	0.31	209,210,213,213	0

6.5 Other polymers [i](#)

There are no such residues in this entry.