



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 12:04 AM EDT

PDB ID : 6QSF  
Title : Crystal structure of Pizza6S in the presence of Keggin (STA)  
Authors : Noguchi, H.; Vandebroek, L.; Kamata, K.; Tame, J.R.H.; Van Meervelt, L.;  
Parac-Vogt, T.N.; Voet, A.R.D.  
Deposited on : 2019-02-20  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	 4% 85% 12% .
1	B	256	 6% 81% 17% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SIW	B	301[A]	-	-	X	-
2	SIW	B	301[B]	-	-	X	-




<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	134	Total 134	O 134	0	0
3	B	99	Total 99	O 99	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pizza6S



- Molecule 1: Pizza6S



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.24Å 69.78Å 62.67Å 90.00° 104.67° 90.00°	Depositor
Resolution (Å)	47.64 – 1.50 47.64 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.64-1.50) 99.8 (47.64-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.77 (at 1.50Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.224 , 0.238 0.227 , 0.239	Depositor DCC
$R_{free}$ test set	3244 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.5	Xtrriage
Anisotropy	0.600	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4046	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.4834e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/1839	0.81	0/2533
1	B	0.58	0/1873	0.78	0/2580
All	All	0.60	0/3712	0.80	0/5113

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1810	0	1789	19	0
1	B	1844	0	1821	39	0
2	A	53	0	0	7	0
2	B	106	0	0	18	0
3	A	134	0	0	3	0
3	B	99	0	0	3	0
All	All	4046	0	3610	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:199:HIS:HE2	1:B:224:THR:HG1	1.33	0.76
1:B:204:VAL:HG21	1:B:238[A]:VAL:HG11	1.71	0.72
1:A:120:VAL:HG21	1:A:154:VAL:HG21	1.71	0.71
1:A:36:VAL:HG21	1:A:70[B]:VAL:HG11	1.76	0.66
1:A:3:THR:N	3:A:402:HOH:O	2.29	0.65
1:A:143:GLY:N	1:A:155[A]:THR:OG1	2.31	0.64
1:B:203[B]:ARG:NH1	3:B:401:HOH:O	2.32	0.62
1:B:230[B]:VAL:HG12	1:B:236:VAL:HG22	1.81	0.62
1:B:224:THR:O	1:B:241:HIS:HB3	2.00	0.61
1:B:29[A]:THR:HG23	1:B:60:VAL:HB	1.83	0.60
1:B:157:HIS:CD2	2:B:301[B]:SIW:O13E	2.55	0.59
1:B:199:HIS:NE2	1:B:224:THR:OG1	2.32	0.59
1:A:224:THR:O	1:A:241:HIS:HB3	2.03	0.59
1:B:71[A]:THR:HG23	1:B:102:VAL:HB	1.84	0.59
1:B:201:ASN:HA	3:B:426:HOH:O	2.02	0.57
1:B:37[A]:VAL:HG12	1:B:48:VAL:HG22	1.86	0.57
1:B:78:VAL:HG21	1:B:112:VAL:HG21	1.88	0.56
1:B:197[A]:THR:HG23	1:B:228:VAL:HB	1.85	0.56
1:B:176:PRO:HD2	1:B:207:LEU:HD21	1.88	0.55
1:B:100:SER:HB2	2:B:301[B]:SIW:O14E	2.07	0.54
1:B:203[B]:ARG:HD3	1:B:216:VAL:HG11	1.89	0.53
1:A:36:VAL:HG21	1:A:70[A]:VAL:HG21	1.90	0.52
1:B:197[A]:THR:HG22	1:B:204:VAL:HG22	1.92	0.52
2:B:301[B]:SIW:O18	2:B:301[B]:SIW:O4	2.29	0.51
1:A:218:PRO:HG2	1:A:249:LEU:HD21	1.94	0.50
2:A:301:SIW:O18	2:A:301:SIW:O4	2.29	0.50
1:B:14:THR:HB	1:B:241:HIS:NE2	2.27	0.50
2:B:301[A]:SIW:O18	2:B:301[A]:SIW:O4	2.29	0.50
2:B:301[B]:SIW:O19E	2:B:301[B]:SIW:O9E	2.30	0.49
1:B:56:THR:O	1:B:73:HIS:HB3	2.12	0.49
1:B:199:HIS:CD2	1:B:224:THR:HG1	2.28	0.49
2:A:301:SIW:O19E	2:A:301:SIW:O9E	2.30	0.49
1:B:16:SER:HB3	2:B:301[A]:SIW:O6	2.13	0.49
1:B:50:PRO:HG2	1:B:81:LEU:HD21	1.94	0.49
1:B:98:THR:O	1:B:115:HIS:HB3	2.12	0.49
1:B:37[B]:VAL:HG12	1:B:46:GLN:HB2	1.95	0.48
1:B:240:ASP:HB3	1:B:245:ARG:HG2	1.95	0.48
2:B:301[A]:SIW:O23	2:B:301[A]:SIW:O22	2.32	0.48
1:B:195:TYR:CD1	1:B:230[A]:VAL:HG21	2.49	0.48
2:B:301[A]:SIW:O9E	2:B:301[A]:SIW:O19E	2.30	0.48
2:A:301:SIW:O23	2:A:301:SIW:O22	2.32	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:301[A]:SIW:O22	2:B:301[A]:SIW:O17	2.32	0.48
1:B:16:SER:CB	2:B:301[A]:SIW:O6	2.62	0.47
1:B:17:GLY:N	1:B:29[A]:THR:OG1	2.48	0.47
1:B:228:VAL:HG23	1:B:238[A]:VAL:HG12	1.95	0.47
2:B:301[B]:SIW:O17	2:B:301[B]:SIW:O22	2.32	0.47
1:A:151:THR:HG23	1:A:166:ALA:HA	1.96	0.47
2:A:301:SIW:O22	2:A:301:SIW:O17	2.32	0.47
1:B:134:PRO:HD2	1:B:165:LEU:HD21	1.96	0.47
2:B:301[B]:SIW:O22	2:B:301[B]:SIW:O23	2.32	0.47
2:A:301:SIW:O16	2:A:301:SIW:O21	2.34	0.46
1:A:182:THR:HG22	3:A:433:HOH:O	2.15	0.46
1:B:248:LYS:NZ	3:B:404:HOH:O	2.47	0.46
1:B:201:ASN:O	1:B:203[B]:ARG:HG3	2.17	0.45
2:B:301[A]:SIW:O16	2:B:301[A]:SIW:O21	2.34	0.45
2:B:301[B]:SIW:O16	2:B:301[B]:SIW:O21	2.34	0.45
1:A:155[A]:THR:HG21	1:A:184:SER:O	2.17	0.45
1:A:8:PRO:HD2	1:A:39[A]:LEU:HD21	1.98	0.44
1:B:1:SER:HB3	1:B:250:ALA:O	2.16	0.44
1:A:8:PRO:HG2	1:A:39[A]:LEU:HD21	1.99	0.44
1:A:115:HIS:HE1	2:A:301:SIW:O1E	2.01	0.44
1:A:199:HIS:NE2	1:A:224:THR:HB	2.34	0.42
1:A:37[A]:VAL:HG12	1:A:48:VAL:HG22	2.01	0.42
1:A:78:VAL:HG21	1:A:112:VAL:HG21	2.01	0.42
2:B:301[B]:SIW:O4	2:B:301[B]:SIW:O20	2.38	0.42
1:B:201:ASN:HB3	1:B:203[A]:ARG:HD3	2.01	0.42
1:B:1:SER:N	1:B:251:ALA:O	2.52	0.42
1:B:73:HIS:NE2	2:B:301[A]:SIW:O12E	2.52	0.42
1:B:14:THR:O	1:B:31:HIS:HB3	2.19	0.42
2:B:301[A]:SIW:O4	2:B:301[A]:SIW:O20	2.38	0.42
1:A:167:ALA:HA	3:A:517:HOH:O	2.19	0.41
1:A:206:LYS:HB3	1:A:217:LEU:HD11	2.03	0.41
2:A:301:SIW:O4	2:A:301:SIW:O20	2.38	0.41
1:B:204:VAL:HG21	1:B:238[B]:VAL:HG21	2.03	0.41
1:A:75:ASN:HB3	1:A:77:ARG:HD3	2.03	0.40
1:B:157:HIS:HD2	2:B:301[B]:SIW:O13E	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/256 (99%)	241 (95%)	11 (4%)	1 (0%)	34	13
1	B	258/256 (101%)	247 (96%)	11 (4%)	0	100	100
All	All	511/512 (100%)	488 (96%)	22 (4%)	1 (0%)	47	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	98	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/201 (101%)	202 (100%)	1 (0%)	88	78
1	B	207/201 (103%)	207 (100%)	0	100	100
All	All	410/402 (102%)	409 (100%)	1 (0%)	92	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	SIW	B	301[A]	3	28,76,76	4.39	19 (67%)	0,234,234	-	-
2	SIW	B	301[B]	-	28,76,76	4.39	19 (67%)	0,234,234	-	-
2	SIW	A	301	-	28,76,76	4.39	19 (67%)	0,234,234	-	-

All (57) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301[B]	SIW	SI1-O19E	-9.63	1.45	1.62
2	A	301	SIW	SI1-O19E	-9.63	1.45	1.62
2	B	301[A]	SIW	SI1-O19E	-9.61	1.45	1.62
2	B	301[A]	SIW	SI1-O4	-8.84	1.47	1.62
2	B	301[B]	SIW	SI1-O4	-8.83	1.47	1.62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	SIW	SI1-O4	-8.80	1.47	1.62
2	B	301[B]	SIW	W2E-O19E	7.73	2.63	2.33
2	A	301	SIW	W2E-O19E	7.72	2.63	2.33
2	B	301[A]	SIW	W2E-O19E	7.71	2.63	2.33
2	B	301[B]	SIW	W3E-O19E	7.67	2.63	2.33
2	A	301	SIW	W3E-O19E	7.67	2.63	2.33
2	B	301[A]	SIW	W3E-O19E	7.66	2.63	2.33
2	A	301	SIW	W4E-O4	7.65	2.63	2.33
2	B	301[A]	SIW	W4E-O4	7.65	2.63	2.33
2	B	301[B]	SIW	W4E-O4	7.63	2.63	2.33
2	A	301	SIW	W5E-O4	7.40	2.62	2.33
2	B	301[B]	SIW	W5E-O4	7.40	2.62	2.33
2	B	301[A]	SIW	W5E-O4	7.39	2.62	2.33
2	B	301[B]	SIW	W7-O21	7.09	2.61	2.33
2	A	301	SIW	W7-O21	7.07	2.61	2.33
2	B	301[A]	SIW	W7-O21	7.07	2.61	2.33
2	B	301[B]	SIW	W6-O22	4.08	2.49	2.33
2	B	301[A]	SIW	W6-O22	4.08	2.49	2.33
2	A	301	SIW	W6-O22	4.05	2.49	2.33
2	A	301	SIW	SI1-O21	2.98	1.68	1.62
2	B	301[B]	SIW	SI1-O21	2.98	1.68	1.62
2	B	301[A]	SIW	SI1-O21	2.94	1.67	1.62
2	B	301[B]	SIW	W6-O5	-2.40	1.59	2.16
2	B	301[A]	SIW	W6-O5	-2.40	1.59	2.16
2	A	301	SIW	W6-O5	-2.40	1.59	2.16
2	B	301[B]	SIW	W1-O12	-2.27	1.62	2.16
2	B	301[A]	SIW	W1-O12	-2.27	1.62	2.16
2	A	301	SIW	W1E-O12E	-2.27	1.62	2.16
2	B	301[B]	SIW	W1E-O12E	-2.27	1.62	2.16
2	A	301	SIW	W1-O12	-2.27	1.62	2.16
2	B	301[A]	SIW	W1E-O12E	-2.27	1.62	2.16
2	A	301	SIW	SI1-O22	2.19	1.66	1.62
2	B	301[A]	SIW	SI1-O22	2.17	1.66	1.62
2	B	301[B]	SIW	W3-O11	-2.15	1.65	2.16
2	B	301[B]	SIW	SI1-O22	2.15	1.66	1.62
2	A	301	SIW	W3E-O11E	-2.15	1.65	2.16
2	A	301	SIW	W3-O11	-2.15	1.65	2.16
2	B	301[A]	SIW	W3-O11	-2.15	1.65	2.16
2	B	301[B]	SIW	W3E-O11E	-2.15	1.65	2.16
2	B	301[A]	SIW	W3E-O11E	-2.15	1.65	2.16
2	B	301[A]	SIW	W4-O14	-2.11	1.66	2.16
2	A	301	SIW	W4-O14	-2.10	1.66	2.16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301[A]	SIW	W4E-O14E	-2.10	1.66	2.16
2	B	301[B]	SIW	W4E-O14E	-2.10	1.66	2.16
2	A	301	SIW	W4E-O14E	-2.10	1.66	2.16
2	B	301[B]	SIW	W4-O14	-2.10	1.66	2.16
2	B	301[B]	SIW	W5-O13	-2.02	1.68	2.16
2	B	301[A]	SIW	W5E-O13E	-2.02	1.68	2.16
2	B	301[A]	SIW	W5-O13	-2.02	1.68	2.16
2	A	301	SIW	W5E-O13E	-2.02	1.68	2.16
2	A	301	SIW	W5-O13	-2.02	1.68	2.16
2	B	301[B]	SIW	W5E-O13E	-2.02	1.68	2.16

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

3 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301[A]	SIW	9	0
2	B	301[B]	SIW	9	0
2	A	301	SIW	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	248/256 (96%)	0.71	10 (4%) 38 42	11, 19, 32, 45	0
1	B	251/256 (98%)	0.88	15 (5%) 21 23	13, 22, 35, 45	1 (0%)
All	All	499/512 (97%)	0.79	25 (5%) 28 31	11, 21, 34, 45	1 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	167	ALA	4.8
1	A	168	GLY	4.3
1	A	166	ALA	3.9
1	A	76[A]	ASN	3.6
1	B	203[A]	ARG	3.5
1	B	224	THR	3.2
1	A	74	GLY	2.9
1	B	41	ALA	2.8
1	A	169	SER	2.6
1	B	56	THR	2.5
1	B	184	SER	2.5
1	B	210	GLY	2.5
1	A	171	THR	2.3
1	B	74	GLY	2.3
1	B	79	VAL	2.3
1	B	52	THR	2.3
1	A	216	VAL	2.3
1	B	87	THR	2.2
1	B	202	ASN	2.2
1	B	84	GLY	2.2
1	B	123	LEU	2.2
1	A	170	ASN	2.2
1	B	116	GLY	2.1
1	B	170	ASN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	200	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SIW	B	301[A]	53/53	0.97	0.13	6,20,27,43	53
2	SIW	B	301[B]	53/53	0.97	0.13	8,18,25,29	53
2	SIW	A	301	53/53	0.99	0.10	11,17,25,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around SIW B 301 (A):**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SIW B 301 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.