



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 10:51 AM EDT

PDB ID : 6RPD
Title : Structure of ferrous KpDyP in complex with cyanide
Authors : Pfanzagl, V.; Beale, J.; Hofbauer, S.
Deposited on : 2019-05-14
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	 94%
1	B	303	 96%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10335 atoms, of which 4827 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iron-dependent peroxidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	299	4820	1529	2381	432	470	8	0	22	0
1	B	298	4705	1497	2330	417	453	8	0	13	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PRO	-	expression tag	UNP A0A0W8ATM9
A	-1	LEU	-	expression tag	UNP A0A0W8ATM9
A	0	GLY	-	expression tag	UNP A0A0W8ATM9
A	300	GLY	-	expression tag	UNP A0A0W8ATM9
B	-2	PRO	-	expression tag	UNP A0A0W8ATM9
B	-1	LEU	-	expression tag	UNP A0A0W8ATM9
B	0	GLY	-	expression tag	UNP A0A0W8ATM9
B	300	GLY	-	expression tag	UNP A0A0W8ATM9

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is CYANIDE ION (three-letter code: CYN) (formula: CN).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			2	1	1		
5	B	1	Total	C	N	0	0
			2	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	312	Total 312	O 312	0	0
6	B	248	Total 248	O 248	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Iron-dependent peroxidase



- Molecule 1: Iron-dependent peroxidase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.57Å 75.74Å 75.71Å 90.00° 107.85° 90.00°	Depositor
Resolution (Å)	40.62 – 1.52 40.63 – 1.52	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.62-1.52) 97.6 (40.63-1.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.52Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.171 , 0.189 0.170 , 0.191	Depositor DCC
R_{free} test set	4058 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtrriage
Anisotropy	0.198	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10335	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYN, HEM, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	10/2564 (0.4%)	0.67	3/3464 (0.1%)
1	B	0.52	1/2475 (0.0%)	0.63	1/3346 (0.0%)
All	All	0.60	11/5039 (0.2%)	0.65	4/6810 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	189	PRO	N-CA	12.17	1.68	1.47
1	B	3	GLN	C-N	-9.00	1.13	1.34
1	A	75	GLU	CD-OE1	-8.71	1.16	1.25
1	A	188	VAL	C-N	7.54	1.48	1.34
1	A	75	GLU	CD-OE2	-5.52	1.19	1.25
1	A	192	GLU	C-O	-5.43	1.13	1.23
1	A	144[A]	GLY	N-CA	5.31	1.54	1.46
1	A	144[B]	GLY	N-CA	5.31	1.54	1.46
1	A	192	GLU	CD-OE2	-5.08	1.20	1.25
1	A	143[A]	ASP	C-N	5.03	1.42	1.33
1	A	143[B]	ASP	C-N	5.03	1.42	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143[A]	ASP	C-N-CA	-6.92	107.76	122.30
1	A	143[B]	ASP	C-N-CA	-6.92	107.76	122.30
1	B	3	GLN	O-C-N	5.92	132.17	122.70
1	A	189	PRO	CA-N-CD	-5.25	104.15	111.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	189	PRO	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2439	2381	2298	4	0
1	B	2375	2330	2276	0	0
2	A	43	30	30	0	0
2	B	43	30	30	1	0
3	A	24	32	32	0	0
3	B	18	24	24	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	1	0
6	A	312	0	0	0	0
6	B	248	0	0	0	0
All	All	5508	4827	4690	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:PRO:N	1:A:189:PRO:CA	1.67	1.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:PRO:N	1:A:189:PRO:C	2.56	0.49
1:A:165:ASP:OD1	1:A:295[B]:ARG:NH2	2.47	0.48
2:B:401:HEM:NB	5:B:406:CYN:C	2.84	0.41
1:A:156:GLU:HA	1:A:161:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/303 (105%)	315 (99%)	3 (1%)	0	100	100
1	B	309/303 (102%)	302 (98%)	7 (2%)	0	100	100
All	All	627/606 (104%)	617 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/244 (107%)	256 (98%)	5 (2%)	57	26
1	B	252/244 (103%)	246 (98%)	6 (2%)	49	19
All	All	513/488 (105%)	502 (98%)	11 (2%)	60	22

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99[A]	LEU
1	A	99[B]	LEU
1	A	162[A]	ASP
1	A	162[B]	ASP
1	A	178	HIS
1	B	99[A]	LEU
1	B	99[B]	LEU
1	B	125	GLU
1	B	162	ASP
1	B	181	LYS
1	B	278	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	401	1	42,50,50	1.34	7 (16%)	46,82,82	1.52	10 (21%)
2	HEM	B	401	1	42,50,50	1.34	5 (11%)	46,82,82	1.63	11 (23%)
3	GOL	B	403	-	5,5,5	0.07	0	5,5,5	0.13	0
3	GOL	B	402	-	5,5,5	0.11	0	5,5,5	0.28	0
3	GOL	A	404	-	5,5,5	0.09	0	5,5,5	0.12	0
3	GOL	A	403	-	5,5,5	0.14	0	5,5,5	0.21	0
5	CYN	A	407	-	1,1,1	0.09	0	-		
3	GOL	B	404	-	5,5,5	0.14	0	5,5,5	0.23	0
3	GOL	A	402	-	5,5,5	0.05	0	5,5,5	0.06	0
3	GOL	A	405	-	5,5,5	0.28	0	5,5,5	0.19	0
5	CYN	B	406	-	1,1,1	0.14	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	401	1	-	4/12/54/54	-
2	HEM	B	401	1	-	4/12/54/54	-
3	GOL	B	403	-	-	0/4/4/4	-
3	GOL	B	402	-	-	0/4/4/4	-
3	GOL	A	404	-	-	0/4/4/4	-
3	GOL	A	403	-	-	2/4/4/4	-
3	GOL	B	404	-	-	4/4/4/4	-
3	GOL	A	402	-	-	0/4/4/4	-
3	GOL	A	405	-	-	2/4/4/4	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	HEM	C3C-C4C	3.83	1.47	1.41
2	A	401	HEM	C3C-C4C	3.66	1.46	1.41
2	B	401	HEM	C4B-NB	-2.76	1.33	1.38
2	B	401	HEM	CHB-C1B	2.65	1.41	1.34
2	A	401	HEM	CHB-C1B	2.60	1.41	1.34
2	A	401	HEM	C1D-C2D	2.56	1.49	1.44
2	B	401	HEM	C4D-ND	-2.34	1.36	1.40
2	A	401	HEM	C1D-ND	-2.22	1.34	1.38
2	A	401	HEM	C1B-NB	-2.19	1.36	1.40
2	A	401	HEM	CHA-C4D	2.15	1.39	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	HEM	CHA-C4D	2.08	1.39	1.34
2	A	401	HEM	C4D-ND	-2.06	1.36	1.40

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	HEM	O2A-CGA-CBA	4.01	126.68	114.00
2	B	401	HEM	CHC-C4B-NB	3.54	128.24	124.44
2	B	401	HEM	C2D-C1D-ND	3.08	113.46	109.90
2	A	401	HEM	O2A-CGA-O1A	-3.04	115.52	123.33
2	A	401	HEM	C2D-C1D-ND	3.01	113.38	109.90
2	A	401	HEM	CHC-C4B-NB	2.99	127.65	124.44
2	A	401	HEM	CHD-C1D-C2D	-2.92	120.42	125.03
2	A	401	HEM	C2B-C1B-NB	2.51	112.72	109.84
2	B	401	HEM	CHD-C1D-C2D	-2.46	121.14	125.03
2	B	401	HEM	O2A-CGA-O1A	-2.45	117.04	123.33
2	A	401	HEM	C1D-C2D-C3D	-2.38	104.48	106.98
2	B	401	HEM	C3D-C4D-ND	2.22	112.61	110.17
2	B	401	HEM	O1A-CGA-CBA	-2.15	116.28	123.09
2	A	401	HEM	CMA-C3A-C4A	-2.14	125.31	128.46
2	B	401	HEM	CBA-CAA-C2A	2.14	116.14	112.54
2	B	401	HEM	CMA-C3A-C4A	-2.13	125.34	128.46
2	A	401	HEM	CBA-CAA-C2A	2.11	116.09	112.54
2	B	401	HEM	C1D-C2D-C3D	-2.11	104.76	106.98
2	A	401	HEM	CHA-C4D-C3D	-2.08	121.38	125.23
2	B	401	HEM	CHA-C4D-C3D	-2.08	121.40	125.23
2	A	401	HEM	C3D-C4D-ND	2.03	112.40	110.17

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	GOL	O1-C1-C2-C3
3	B	404	GOL	O1-C1-C2-C3
3	B	404	GOL	C1-C2-C3-O3
3	A	403	GOL	O1-C1-C2-O2
3	B	404	GOL	O1-C1-C2-O2
3	A	405	GOL	C1-C2-C3-O3
3	A	405	GOL	O1-C1-C2-C3
2	B	401	HEM	CAD-CBD-CGD-O2D
3	B	404	GOL	O2-C2-C3-O3
2	B	401	HEM	CAD-CBD-CGD-O1D

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	401	HEM	CAD-CBD-CGD-O1D
2	A	401	HEM	CAD-CBD-CGD-O2D
2	A	401	HEM	CAA-CBA-CGA-O1A
2	B	401	HEM	CAA-CBA-CGA-O2A
2	A	401	HEM	CAA-CBA-CGA-O2A
2	B	401	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	HEM	1	0
5	B	406	CYN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	3:GLN	C	4[A]:VAL	N	1.13
1	B	3:GLN	C	4[B]:VAL	N	1.12

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/303 (98%)	-0.24	3 (1%) 82 85	13, 21, 40, 52	0
1	B	298/303 (98%)	-0.16	7 (2%) 60 65	15, 24, 47, 73	0
All	All	597/606 (98%)	-0.20	10 (1%) 70 74	13, 22, 44, 73	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	GLY	8.8
1	B	299	LEU	3.5
1	A	295[A]	ARG	3.5
1	B	27	VAL	2.8
1	A	294[A]	GLU	2.6
1	B	86	LEU	2.5
1	A	162[A]	ASP	2.4
1	B	26	ASP	2.3
1	B	85	GLY	2.3
1	B	84	LYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	404	6/6	0.69	0.30	20,20,20,20	0
3	GOL	A	405	6/6	0.76	0.25	20,20,20,20	0
3	GOL	B	402	6/6	0.85	0.15	31,41,45,45	0
3	GOL	A	404	6/6	0.85	0.25	27,30,35,35	0
3	GOL	A	402	6/6	0.86	0.10	40,42,45,45	0
3	GOL	A	403	6/6	0.90	0.13	27,35,38,38	0
3	GOL	B	403	6/6	0.91	0.32	33,40,43,43	0
5	CYN	B	406	2/2	0.93	0.14	22,22,22,27	0
4	MG	B	405	1/1	0.98	0.12	25,25,25,25	0
5	CYN	A	407	2/2	0.98	0.10	19,19,19,22	0
4	MG	A	406	1/1	0.98	0.07	25,25,25,25	0
2	HEM	B	401	43/43	0.99	0.06	16,18,22,27	0
2	HEM	A	401	43/43	0.99	0.07	14,16,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.