



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:08 AM UTC

PDB ID : 6UB1 / pdb_00006ub1
Title : Crystal structure of a GH128 (subgroup III) curdlan-specific exo-beta-1,3-galactanase from *Blastomyces gilchristii* (BgGH128_III) in complex with laminaribiose at -3 and -2 subsites
Authors : Costa, P.A.C.R.; Santos, C.R.; Domingues, M.N.; Lima, E.A.; Mandelli, F.; Murakami, M.T.
Deposited on : 2019-09-11
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	 82% 5% 12%
1	B	261	 83% 5% 12%
1	C	261	 80% 8% 12%
1	D	261	 82% 6% 12%
2	E	2	 50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 50% 50%
2	G	2	 50% 50%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8197 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCOSIDE HYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	1750	1120	295	329	6	0	2	0
1	B	230	1755	1122	297	330	6	0	2	0
1	C	230	1756	1123	296	331	6	0	2	0
1	D	230	1759	1125	296	332	6	0	3	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP A0A179UGT5
A	3	GLY	-	expression tag	UNP A0A179UGT5
A	4	SER	-	expression tag	UNP A0A179UGT5
A	5	SER	-	expression tag	UNP A0A179UGT5
A	6	HIS	-	expression tag	UNP A0A179UGT5
A	7	HIS	-	expression tag	UNP A0A179UGT5
A	8	HIS	-	expression tag	UNP A0A179UGT5
A	9	HIS	-	expression tag	UNP A0A179UGT5
A	10	HIS	-	expression tag	UNP A0A179UGT5
A	11	HIS	-	expression tag	UNP A0A179UGT5
A	12	SER	-	expression tag	UNP A0A179UGT5
A	13	SER	-	expression tag	UNP A0A179UGT5
A	14	GLY	-	expression tag	UNP A0A179UGT5
A	15	LEU	-	expression tag	UNP A0A179UGT5
A	16	VAL	-	expression tag	UNP A0A179UGT5
A	17	PRO	-	expression tag	UNP A0A179UGT5
A	18	ALA	-	expression tag	UNP A0A179UGT5
A	19	GLY	-	expression tag	UNP A0A179UGT5
A	20	SER	-	expression tag	UNP A0A179UGT5
A	21	HIS	-	expression tag	UNP A0A179UGT5
A	22	MET	-	expression tag	UNP A0A179UGT5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	MET	-	initiating methionine	UNP A0A179UGT5
B	3	GLY	-	expression tag	UNP A0A179UGT5
B	4	SER	-	expression tag	UNP A0A179UGT5
B	5	SER	-	expression tag	UNP A0A179UGT5
B	6	HIS	-	expression tag	UNP A0A179UGT5
B	7	HIS	-	expression tag	UNP A0A179UGT5
B	8	HIS	-	expression tag	UNP A0A179UGT5
B	9	HIS	-	expression tag	UNP A0A179UGT5
B	10	HIS	-	expression tag	UNP A0A179UGT5
B	11	HIS	-	expression tag	UNP A0A179UGT5
B	12	SER	-	expression tag	UNP A0A179UGT5
B	13	SER	-	expression tag	UNP A0A179UGT5
B	14	GLY	-	expression tag	UNP A0A179UGT5
B	15	LEU	-	expression tag	UNP A0A179UGT5
B	16	VAL	-	expression tag	UNP A0A179UGT5
B	17	PRO	-	expression tag	UNP A0A179UGT5
B	18	ALA	-	expression tag	UNP A0A179UGT5
B	19	GLY	-	expression tag	UNP A0A179UGT5
B	20	SER	-	expression tag	UNP A0A179UGT5
B	21	HIS	-	expression tag	UNP A0A179UGT5
B	22	MET	-	expression tag	UNP A0A179UGT5
C	2	MET	-	initiating methionine	UNP A0A179UGT5
C	3	GLY	-	expression tag	UNP A0A179UGT5
C	4	SER	-	expression tag	UNP A0A179UGT5
C	5	SER	-	expression tag	UNP A0A179UGT5
C	6	HIS	-	expression tag	UNP A0A179UGT5
C	7	HIS	-	expression tag	UNP A0A179UGT5
C	8	HIS	-	expression tag	UNP A0A179UGT5
C	9	HIS	-	expression tag	UNP A0A179UGT5
C	10	HIS	-	expression tag	UNP A0A179UGT5
C	11	HIS	-	expression tag	UNP A0A179UGT5
C	12	SER	-	expression tag	UNP A0A179UGT5
C	13	SER	-	expression tag	UNP A0A179UGT5
C	14	GLY	-	expression tag	UNP A0A179UGT5
C	15	LEU	-	expression tag	UNP A0A179UGT5
C	16	VAL	-	expression tag	UNP A0A179UGT5
C	17	PRO	-	expression tag	UNP A0A179UGT5
C	18	ALA	-	expression tag	UNP A0A179UGT5
C	19	GLY	-	expression tag	UNP A0A179UGT5
C	20	SER	-	expression tag	UNP A0A179UGT5
C	21	HIS	-	expression tag	UNP A0A179UGT5
C	22	MET	-	expression tag	UNP A0A179UGT5

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	2	MET	-	initiating methionine	UNP A0A179UGT5
D	3	GLY	-	expression tag	UNP A0A179UGT5
D	4	SER	-	expression tag	UNP A0A179UGT5
D	5	SER	-	expression tag	UNP A0A179UGT5
D	6	HIS	-	expression tag	UNP A0A179UGT5
D	7	HIS	-	expression tag	UNP A0A179UGT5
D	8	HIS	-	expression tag	UNP A0A179UGT5
D	9	HIS	-	expression tag	UNP A0A179UGT5
D	10	HIS	-	expression tag	UNP A0A179UGT5
D	11	HIS	-	expression tag	UNP A0A179UGT5
D	12	SER	-	expression tag	UNP A0A179UGT5
D	13	SER	-	expression tag	UNP A0A179UGT5
D	14	GLY	-	expression tag	UNP A0A179UGT5
D	15	LEU	-	expression tag	UNP A0A179UGT5
D	16	VAL	-	expression tag	UNP A0A179UGT5
D	17	PRO	-	expression tag	UNP A0A179UGT5
D	18	ALA	-	expression tag	UNP A0A179UGT5
D	19	GLY	-	expression tag	UNP A0A179UGT5
D	20	SER	-	expression tag	UNP A0A179UGT5
D	21	HIS	-	expression tag	UNP A0A179UGT5
D	22	MET	-	expression tag	UNP A0A179UGT5

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	2	Total	C	O	0	0	0
			23	12	11			
2	F	2	Total	C	O	0	0	0
			23	12	11			
2	G	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	C O	0	0
			12	6 6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	263	Total	O	0	0
			263	263		
4	B	273	Total	O	0	0
			273	273		
4	C	266	Total	O	0	0
			266	266		
4	D	294	Total	O	0	0
			294	294		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GLYCOSIDE HYDROLASE




- Molecule 1: GLYCOSIDE HYDROLASE



- Molecule 1: GLYCOSIDE HYDROLASE



- Molecule 1: GLYCOSIDE HYDROLASE




- Molecule 2: beta-D-glucopyranose-(1-3)-beta-D-glucopyranose

Chain E:  50% 50%

BGC1
BGC2

- Molecule 2: beta-D-glucopyranose-(1-3)-beta-D-glucopyranose

Chain F:  50% 50%

BGC1
BGC2

- Molecule 2: beta-D-glucopyranose-(1-3)-beta-D-glucopyranose

Chain G:  50% 50%

BGC1
BGC2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	56.32Å 60.80Å 63.55Å 69.67° 78.24° 77.86°	Depositor
Resolution (Å)	35.49 – 1.60 35.49 – 1.60	Depositor EDS
% Data completeness (in resolution range)	86.3 (35.49-1.60) 86.2 (35.49-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.60 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.160 , 0.197 0.172 , 0.207	Depositor DCC
R_{free} test set	4350 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	13.9	Xtrriage
Anisotropy	0.551	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.035 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8197	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 90.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2589e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.17	2/1806 (0.1%)	1.34	4/2456 (0.2%)
1	B	1.20	2/1807 (0.1%)	1.28	3/2457 (0.1%)
1	C	1.17	4/1808 (0.2%)	1.23	4/2458 (0.2%)
1	D	1.13	1/1815 (0.1%)	1.24	4/2468 (0.2%)
All	All	1.17	9/7236 (0.1%)	1.27	15/9839 (0.2%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	210	GLU	CD-OE1	6.72	1.38	1.25
1	A	168	ILE	C-O	5.90	1.30	1.24
1	C	164	GLY	C-O	5.78	1.32	1.24
1	C	62	GLU	N-CA	5.67	1.50	1.46
1	D	201	ILE	C-O	5.57	1.30	1.24
1	B	236	GLU	CD-OE1	5.46	1.35	1.25
1	A	75	VAL	N-CA	5.44	1.50	1.46
1	C	104	ALA	C-O	5.23	1.29	1.23
1	C	116	MET	C-O	-5.15	1.17	1.23

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	ARG	CB-CG-CD	7.81	129.25	111.30
1	B	119	GLU	CB-CA-C	7.57	123.72	110.85
1	C	181	ASN	CA-CB-CG	6.66	119.26	112.60
1	C	214	ASP	CA-CB-CG	6.41	119.00	112.60
1	C	181	ASN	N-CA-C	-5.76	106.16	112.72
1	C	187	ARG	NE-CZ-NH2	5.71	124.34	119.20
1	B	181	ASN	CA-CB-CG	5.65	118.25	112.60
1	D	214	ASP	CA-CB-CG	5.53	118.13	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	242	PHE	CA-CB-CG	5.51	119.31	113.80
1	D	87	TRP	CA-C-O	-5.49	115.06	120.82
1	A	191	GLU	N-CA-C	-5.24	105.65	111.36
1	B	37	ARG	CG-CD-NE	5.20	123.44	112.00
1	D	42	ASN	CA-CB-CG	5.12	117.72	112.60
1	A	213	GLY	CA-C-O	-5.06	116.50	121.05
1	A	220	ASN	CA-CB-CG	-5.00	107.60	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1750	0	1657	10	0
1	B	1755	0	1659	5	0
1	C	1756	0	1659	12	0
1	D	1759	0	1662	10	0
2	E	23	0	21	1	0
2	F	23	0	21	1	0
2	G	23	0	21	1	0
3	B	12	0	12	0	0
4	A	263	0	0	6	0
4	B	273	0	0	2	0
4	C	266	0	0	2	0
4	D	294	0	0	1	0
All	All	8197	0	6712	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:100:LYS:HE3	4:C:493:HOH:O	1.51	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:203:THR:HG21	1:D:236:GLU:OE2	1.63	0.98
1:C:203:THR:HG21	1:C:236[B]:GLU:OE2	1.64	0.95
1:A:212:GLN:HG3	4:A:573:HOH:O	1.81	0.80
1:D:82:ARG:NH2	2:G:2:BGC:O3	2.17	0.76
1:C:203:THR:OG1	1:C:236[B]:GLU:HG3	1.90	0.71
1:D:203:THR:HG21	1:D:236:GLU:CD	2.21	0.65
1:A:82:ARG:HG2	4:A:479:HOH:O	1.97	0.64
1:C:203:THR:HG21	1:C:236[B]:GLU:CD	2.27	0.60
1:C:82:ARG:NH1	2:F:2:BGC:O3	2.33	0.59
1:D:203:THR:OG1	1:D:236:GLU:HG3	2.02	0.59
1:C:137:ARG:NH1	4:C:401:HOH:O	2.38	0.57
1:A:216:GLU:HG3	4:A:557:HOH:O	2.07	0.55
1:A:119:GLU:HG3	4:A:404:HOH:O	2.07	0.54
1:B:182:VAL:HG22	1:B:221:PHE:CD1	2.42	0.54
1:A:82:ARG:NH2	2:E:2:BGC:O3	2.42	0.52
1:D:119[B]:GLU:HA	1:D:161:VAL:HG11	1.92	0.52
1:D:119[A]:GLU:HA	1:D:161:VAL:HG11	1.92	0.52
1:A:175:TRP:O	1:A:209:PHE:HA	2.10	0.52
1:D:33:GLY:N	4:D:407:HOH:O	2.44	0.51
1:C:175:TRP:O	1:C:209:PHE:HA	2.11	0.51
1:C:69:THR:HG21	1:C:72:ILE:HD12	1.95	0.49
1:B:175:TRP:O	1:B:209:PHE:HA	2.13	0.48
1:A:205:TRP:CZ2	1:A:237:ARG:HD2	2.49	0.48
1:D:175:TRP:O	1:D:209:PHE:HA	2.15	0.47
1:B:72:ILE:HD11	4:B:644:HOH:O	2.14	0.47
1:A:119:GLU:HA	1:A:161:VAL:HG11	1.97	0.46
1:C:62:GLU:HB3	1:C:63:PRO:HD2	1.98	0.46
1:C:69:THR:CG2	1:C:72:ILE:HD12	2.47	0.45
1:A:119:GLU:CG	4:A:404:HOH:O	2.65	0.45
1:C:241:PHE:HA	1:C:242:PHE:HB3	1.99	0.44
1:C:119:GLU:HA	1:C:161:VAL:HG11	2.01	0.42
1:B:216:GLU:HG2	4:B:530:HOH:O	2.19	0.41
1:D:142:ALA:HA	1:D:172:ALA:O	2.21	0.41
1:B:62:GLU:HB3	1:B:63:PRO:HD2	2.03	0.41
1:D:203:THR:HG21	1:D:236:GLU:CG	2.51	0.41
1:A:119:GLU:OE1	4:A:401:HOH:O	2.22	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/261 (88%)	223 (97%)	6 (3%)	0	100	100
1	B	229/261 (88%)	223 (97%)	6 (3%)	0	100	100
1	C	229/261 (88%)	224 (98%)	5 (2%)	0	100	100
1	D	230/261 (88%)	224 (97%)	6 (3%)	0	100	100
All	All	917/1044 (88%)	894 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/194 (88%)	167 (98%)	4 (2%)	44	20
1	B	171/194 (88%)	170 (99%)	1 (1%)	78	66
1	C	171/194 (88%)	170 (99%)	1 (1%)	78	66
1	D	172/194 (89%)	172 (100%)	0	100	100
All	All	685/776 (88%)	679 (99%)	6 (1%)	70	55

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	GLU
1	A	204	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	212	GLN
1	A	248	LYS
1	B	183	ASP
1	C	204	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	42	ASN
1	B	88	ASN
1	B	106	ASN
1	B	113	GLN
1	C	42	ASN
1	C	88	ASN
1	C	218	GLN
1	D	42	ASN
1	D	218	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	E	1	2	12,12,12	1.57	2 (16%)	17,17,17	2.01	5 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	E	2	2	11,11,12	0.61	0	15,15,17	2.12	7 (46%)
2	BGC	F	1	2	12,12,12	1.24	2 (16%)	17,17,17	1.95	7 (41%)
2	BGC	F	2	2	11,11,12	1.09	1 (9%)	15,15,17	2.56	4 (26%)
2	BGC	G	1	2	12,12,12	1.78	2 (16%)	17,17,17	3.17	11 (64%)
2	BGC	G	2	2	11,11,12	0.83	0	15,15,17	1.79	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	E	1	2	-	0/2/22/22	0/1/1/1
2	BGC	E	2	2	-	2/2/19/22	0/1/1/1
2	BGC	F	1	2	-	0/2/22/22	0/1/1/1
2	BGC	F	2	2	-	2/2/19/22	0/1/1/1
2	BGC	G	1	2	-	1/2/22/22	0/1/1/1
2	BGC	G	2	2	-	2/2/19/22	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	BGC	O1-C1	3.97	1.52	1.39
2	G	1	BGC	O2-C2	3.76	1.52	1.43
2	G	1	BGC	O1-C1	3.51	1.50	1.39
2	E	1	BGC	O2-C2	2.86	1.50	1.43
2	F	1	BGC	O2-C2	2.54	1.49	1.43
2	F	1	BGC	O1-C1	2.45	1.47	1.39
2	F	2	BGC	O5-C1	-2.39	1.39	1.43

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	BGC	O5-C1-C2	-6.56	98.77	110.30
2	F	2	BGC	O2-C2-C3	5.54	121.62	110.15
2	F	2	BGC	C1-O5-C5	5.48	119.53	112.19
2	G	2	BGC	C2-C3-C4	-4.37	103.17	110.86
2	G	1	BGC	C4-C3-C2	4.32	118.42	110.83
2	G	1	BGC	O1-C1-C2	4.24	121.28	108.98
2	E	1	BGC	O2-C2-C1	4.20	118.93	109.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	2	BGC	O4-C4-C3	3.83	119.41	110.38
2	E	2	BGC	O3-C3-C2	-3.76	102.37	110.05
2	G	1	BGC	O4-C4-C5	3.67	118.36	109.32
2	G	1	BGC	C6-C5-C4	3.56	121.75	113.02
2	F	1	BGC	O4-C4-C5	3.51	117.96	109.32
2	E	1	BGC	O1-C1-C2	3.42	118.90	108.98
2	G	1	BGC	O3-C3-C2	-3.35	102.47	110.38
2	E	1	BGC	O2-C2-C3	-3.32	102.55	110.38
2	G	1	BGC	C3-C4-C5	-3.31	104.23	110.23
2	E	1	BGC	C4-C3-C2	3.24	116.52	110.83
2	G	1	BGC	O3-C3-C4	-3.21	102.81	110.38
2	F	1	BGC	O3-C3-C4	-3.13	102.99	110.38
2	E	2	BGC	O5-C5-C4	-2.96	103.62	110.83
2	E	1	BGC	O3-C3-C4	-2.94	103.44	110.38
2	G	1	BGC	O2-C2-C1	2.93	116.01	109.25
2	E	2	BGC	O2-C2-C1	2.83	115.71	109.22
2	G	1	BGC	O1-C1-O5	2.81	118.76	110.41
2	F	1	BGC	O2-C2-C1	2.73	115.54	109.25
2	G	1	BGC	O4-C4-C3	-2.70	104.02	110.38
2	F	1	BGC	O4-C4-C3	-2.69	104.03	110.38
2	G	2	BGC	C1-C2-C3	-2.61	105.85	109.64
2	F	1	BGC	O1-C1-O5	-2.57	102.79	110.41
2	E	2	BGC	O2-C2-C3	2.53	115.40	110.15
2	E	2	BGC	O4-C4-C3	-2.53	104.41	110.38
2	E	2	BGC	C2-C3-C4	-2.46	106.53	110.86
2	F	2	BGC	C1-C2-C3	-2.40	106.14	109.64
2	F	1	BGC	O5-C1-C2	2.34	114.42	110.30
2	G	2	BGC	O5-C5-C4	-2.32	105.19	110.83
2	E	2	BGC	O4-C4-C5	2.31	115.00	109.32
2	F	1	BGC	O3-C3-C2	-2.15	105.30	110.38

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	2	BGC	O5-C5-C6-O6
2	F	2	BGC	C4-C5-C6-O6
2	G	2	BGC	O5-C5-C6-O6
2	G	2	BGC	C4-C5-C6-O6
2	E	2	BGC	O5-C5-C6-O6
2	G	1	BGC	C4-C5-C6-O6
2	E	2	BGC	C4-C5-C6-O6



There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2	BGC	1	0
2	F	2	BGC	1	0
2	E	2	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	B	301	-	12,12,12	0.99	0	17,17,17	2.35	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	B	301	-	-	2/2/22/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	BGC	O4-C4-C3	-5.05	98.47	110.38
3	B	301	BGC	O2-C2-C1	3.21	116.66	109.25
3	B	301	BGC	C1-C2-C3	3.16	116.80	110.36
3	B	301	BGC	O1-C1-O5	-3.01	101.47	110.41
3	B	301	BGC	O5-C1-C2	2.97	115.52	110.30
3	B	301	BGC	C1-O5-C5	-2.52	108.77	113.65
3	B	301	BGC	O4-C4-C5	2.43	115.31	109.32

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	BGC	O5-C5-C6-O6
3	B	301	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/261 (88%)	-0.07	2 (0%) 81 84	10, 16, 28, 45	1 (0%)
1	B	230/261 (88%)	-0.09	2 (0%) 81 84	8, 15, 26, 40	1 (0%)
1	C	230/261 (88%)	-0.15	3 (1%) 75 79	7, 16, 24, 38	1 (0%)
1	D	230/261 (88%)	-0.18	0 100 100	8, 16, 24, 32	2 (0%)
All	All	920/1044 (88%)	-0.12	7 (0%) 82 86	7, 16, 26, 45	5 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	33	GLY	3.7
1	C	33	GLY	3.1
1	A	34	SER	2.7
1	A	33	GLY	2.7
1	C	249	GLY	2.4
1	B	34	SER	2.4
1	C	181	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BGC	F	2	11/12	0.86	0.10	25,31,33,36	0
2	BGC	E	1	12/12	0.87	0.10	19,23,24,25	0
2	BGC	G	2	11/12	0.87	0.10	28,36,43,43	0
2	BGC	G	1	12/12	0.89	0.09	19,24,28,33	0
2	BGC	E	2	11/12	0.91	0.07	21,25,29,30	0
2	BGC	F	1	12/12	0.92	0.08	18,23,27,28	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	B	301	12/12	0.92	0.07	18,23,26,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.