



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:44 PM UTC

PDB ID : 6UJ6 / pdb_00006uj6
Title : X-ray Crystal Structure of Chromium-transferrin with Synergistic Anion Malonate
Authors : Petersen, C.M.; Edwards, K.C.; Gilbert, N.C.; Vincent, J.B.; Thompson, M.K.
Deposited on : 2019-10-02
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	698	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Transferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	652	5029	3168	874	944	43	0	0	0

- Molecule 2 is CHROMIUM ION (CCD ID: CR) (formula: Cr) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cr	0	0
			1	1		

- Molecule 3 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is BICARBONATE ION (CCD ID: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	34	Total 34	O 34	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transferrin



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	137.05Å 158.03Å 107.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.22 – 2.68 49.22 – 2.68	Depositor EDS
% Data completeness (in resolution range)	98.9 (49.22-2.68) 99.2 (49.22-2.68)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.220 , 0.270 0.221 , 0.271	Depositor DCC
R_{free} test set	1547 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	48.8	Xtrriage
Anisotropy	0.627	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5087	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CR, MLI, GOL, BCT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.12	1/5141 (0.0%)	1.50	18/6945 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	608	THR	C-O	6.87	1.30	1.24

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	431	TYR	CA-C-O	-6.78	112.09	122.32
1	A	349	THR	CA-CB-OG1	-6.23	100.25	109.60
1	A	527	PRO	CA-C-O	-6.14	116.76	121.31
1	A	297	LYS	N-CA-C	-6.00	106.28	113.97
1	A	665	THR	CA-CB-OG1	-5.56	101.25	109.60
1	A	671	GLY	CA-C-N	5.48	127.93	120.54
1	A	671	GLY	C-N-CA	5.48	127.93	120.54
1	A	625	HIS	N-CA-C	-5.47	103.93	111.81
1	A	641	PHE	CA-CB-CG	5.42	119.22	113.80
1	A	86	VAL	N-CA-C	-5.37	104.34	111.05
1	A	674	TYR	CA-C-N	5.29	127.33	120.56
1	A	674	TYR	C-N-CA	5.29	127.33	120.56
1	A	288	GLN	N-CA-C	-5.26	105.71	111.82
1	A	393	THR	N-CA-C	-5.18	105.55	111.14
1	A	102	GLU	CA-C-O	-5.10	116.00	121.56
1	A	111	GLN	CA-C-N	5.05	127.46	120.29
1	A	111	GLN	C-N-CA	5.05	127.46	120.29
1	A	604	HIS	CA-C-O	-5.01	116.36	121.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5029	0	4862	81	0
2	A	1	0	0	0	0
3	A	7	0	2	1	0
4	A	12	0	16	0	0
5	A	4	0	0	0	0
6	A	34	0	0	0	0
All	All	5087	0	4880	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:429:GLU:OE2	1:A:650:PHE:HB3	1.56	1.05
1:A:570:ALA:HA	1:A:573:LEU:HD12	1.55	0.87
1:A:207:TYR:CE1	1:A:225:LYS:HG3	2.17	0.79
1:A:136:SER:OG	1:A:176:SER:HB3	1.87	0.75
1:A:48:VAL:HG11	1:A:292:HIS:CG	2.22	0.74
1:A:182:ASP:OD2	1:A:185:ASP:OD2	2.10	0.69
1:A:570:ALA:HA	1:A:573:LEU:CD1	2.25	0.67
1:A:481:ILE:HB	1:A:482:PRO:HD3	1.74	0.67
1:A:502:GLY:HA2	1:A:516:LEU:HD12	1.78	0.64
1:A:393:THR:HG21	1:A:414:PHE:CD2	2.33	0.63
1:A:48:VAL:HG11	1:A:292:HIS:CD2	2.32	0.63
1:A:420:LYS:HE2	1:A:682:ARG:HH22	1.64	0.61
1:A:207:TYR:CE1	1:A:225:LYS:CG	2.83	0.61
1:A:88:ASP:OD1	1:A:346:ARG:NH2	2.34	0.60
1:A:107:LYS:N	1:A:107:LYS:HD2	2.17	0.60
1:A:373:LYS:HG2	1:A:649:LEU:HD22	1.83	0.59
1:A:128:MET:HE1	1:A:262:LEU:HD21	1.86	0.58
1:A:443:ALA:HB1	1:A:600:ARG:NH2	2.18	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:MET:HE2	1:A:293:PHE:CE2	2.40	0.56
1:A:273:ARG:NH1	1:A:276:GLY:O	2.37	0.56
1:A:153:LEU:HD11	1:A:262:LEU:HB3	1.87	0.55
1:A:604:HIS:CE1	3:A:702:MLI:O7	2.60	0.55
1:A:50:PRO:HB2	1:A:52:ASP:OD1	2.07	0.55
1:A:364:CYS:HA	1:A:388:VAL:O	2.06	0.55
1:A:429:GLU:OE2	1:A:650:PHE:CB	2.43	0.55
1:A:369:HIS:HB3	1:A:648:LEU:HD21	1.89	0.54
1:A:396:CYS:HB3	1:A:408:MET:SD	2.48	0.54
1:A:557:VAL:HB	1:A:558:PRO:HD3	1.92	0.52
1:A:446:PHE:HB2	1:A:554:HIS:ND1	2.24	0.52
1:A:42:ARG:HG3	1:A:56:VAL:O	2.10	0.52
1:A:665:THR:HG22	1:A:667:GLU:H	1.75	0.52
1:A:135:LYS:HA	1:A:175:GLY:O	2.11	0.51
1:A:48:VAL:CG1	1:A:292:HIS:CD2	2.94	0.51
1:A:624:GLN:NE2	1:A:657:LEU:HB2	2.26	0.51
1:A:127:GLN:HE22	1:A:251:ARG:HG3	1.76	0.50
1:A:400:ILE:HA	1:A:405:ALA:O	2.11	0.50
1:A:45:MET:CE	1:A:293:PHE:CD2	2.94	0.50
1:A:153:LEU:HD13	1:A:247:LEU:HD21	1.94	0.50
1:A:45:MET:HE2	1:A:293:PHE:HE2	1.78	0.49
1:A:444:GLY:O	1:A:600:ARG:HD2	2.12	0.49
1:A:619:ILE:O	1:A:623:GLN:HG2	2.13	0.49
1:A:621:ARG:HG2	1:A:624:GLN:HE21	1.78	0.48
1:A:483:MET:HB3	1:A:495:PHE:CE2	2.49	0.48
1:A:416:TYR:CE2	1:A:678:VAL:HG11	2.48	0.48
1:A:122:LYS:O	1:A:123:ASP:HB2	2.14	0.46
1:A:383:GLY:C	1:A:385:ILE:N	2.74	0.46
1:A:460:TRP:CD1	1:A:489:LYS:HZ1	2.32	0.46
1:A:531:GLU:O	1:A:531:GLU:HG3	2.16	0.45
1:A:23:LYS:O	1:A:54:PRO:HA	2.16	0.45
1:A:382:VAL:O	1:A:382:VAL:HG12	2.17	0.45
1:A:526:GLU:HB3	1:A:528:ASN:ND2	2.32	0.45
1:A:259:ASP:OD1	1:A:697:ARG:NH2	2.49	0.45
1:A:327:ARG:O	1:A:327:ARG:HG3	2.16	0.45
1:A:214:LEU:HB2	1:A:222:ALA:HB2	1.98	0.45
1:A:416:TYR:CD2	1:A:678:VAL:HG11	2.52	0.44
1:A:682:ARG:O	1:A:682:ARG:HG2	2.17	0.44
1:A:338:TYR:CZ	1:A:342:ILE:HG13	2.53	0.44
1:A:128:MET:HE2	1:A:245:LEU:HB3	1.99	0.43
1:A:159:PRO:HG3	1:A:171:ASN:HB3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:540:PHE:O	1:A:543:LEU:HB3	2.19	0.43
1:A:539:ALA:O	1:A:542:CYS:HB3	2.18	0.43
1:A:543:LEU:HB2	1:A:550:ALA:HB2	2.01	0.43
1:A:671:GLY:O	1:A:675:VAL:HG23	2.18	0.43
1:A:374:CYS:C	1:A:387:CYS:SG	3.02	0.42
1:A:45:MET:CE	1:A:293:PHE:HD2	2.31	0.42
1:A:165:LEU:O	1:A:169:VAL:HG23	2.19	0.42
1:A:481:ILE:CB	1:A:482:PRO:HD3	2.47	0.42
1:A:141:LEU:HD23	1:A:141:LEU:C	2.45	0.42
1:A:325:PRO:HA	1:A:326:PRO:HD3	1.96	0.42
1:A:682:ARG:NH1	1:A:694:THR:HG21	2.35	0.42
1:A:106:SER:O	1:A:110:PRO:HA	2.20	0.41
1:A:128:MET:HE2	1:A:128:MET:HB2	1.85	0.41
1:A:444:GLY:O	1:A:600:ARG:CD	2.68	0.41
1:A:128:MET:HE2	1:A:245:LEU:CB	2.50	0.41
1:A:402:ASN:C	1:A:402:ASN:OD1	2.63	0.41
1:A:420:LYS:HE2	1:A:682:ARG:NH2	2.32	0.41
1:A:411:ASP:HA	1:A:604:HIS:CD2	2.56	0.40
1:A:443:ALA:HB1	1:A:600:ARG:HH21	1.85	0.40
1:A:154:LEU:O	1:A:155:TYR:C	2.64	0.40
1:A:602:PRO:HG2	1:A:674:TYR:CD2	2.56	0.40
1:A:207:TYR:CD1	1:A:225:LYS:HG3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	644/698 (92%)	590 (92%)	52 (8%)	2 (0%)	36 57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	649	LEU
1	A	626	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	538/584 (92%)	524 (97%)	14 (3%)	40 68

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	TYR
1	A	107	LYS
1	A	176	SER
1	A	201	LEU
1	A	268	HIS
1	A	279	GLU
1	A	352	GLU
1	A	520	SER
1	A	602	PRO
1	A	615	CYS
1	A	617	HIS
1	A	618	LYS
1	A	643	SER
1	A	650	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	GLN
1	A	249	ASN
1	A	292	HIS
1	A	617	HIS
1	A	625	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	704	-	5,5,5	0.15	0	5,5,5	0.34	0
3	MLI	A	702	2	6,6,6	1.31	0	7,7,7	1.55	0
4	GOL	A	703	-	5,5,5	0.12	0	5,5,5	0.28	0
5	BCT	A	705	-	3,3,3	0.61	0	2,3,3	2.34	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	704	-	-	2/4/4/4	-
3	MLI	A	702	2	-	2/4/4/4	-
4	GOL	A	703	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	A	705	BCT	O2-C-O1	3.07	127.53	119.68

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	703	GOL	O1-C1-C2-O2
4	A	703	GOL	O1-C1-C2-C3
4	A	704	GOL	O1-C1-C2-O2
4	A	704	GOL	O1-C1-C2-C3
3	A	702	MLI	C3-C1-C2-O6
3	A	702	MLI	C3-C1-C2-O7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	MLI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	652/698 (93%)	0.41	51 (7%) 19 16	33, 54, 93, 114	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	382	VAL	4.4
1	A	431	TYR	4.1
1	A	641	PHE	3.9
1	A	359	LYS	3.8
1	A	489	LYS	3.7
1	A	275	ILE	3.6
1	A	630	ASN	3.6
1	A	473	VAL	3.5
1	A	443	ALA	3.5
1	A	455	ALA	3.5
1	A	384	LYS	3.4
1	A	358	CYS	3.4
1	A	383	GLY	3.2
1	A	686	THR	3.2
1	A	352	GLU	3.1
1	A	617	HIS	3.1
1	A	626	LEU	3.0
1	A	372	LEU	3.0
1	A	369	HIS	3.0
1	A	646	LYS	2.9
1	A	627	PHE	2.9
1	A	618	LYS	2.8
1	A	376	GLU	2.8
1	A	504	ALA	2.7
1	A	201	LEU	2.7
1	A	444	GLY	2.7
1	A	591	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	107	LYS	2.7
1	A	368	HIS	2.7
1	A	651	ARG	2.7
1	A	661	HIS	2.6
1	A	560	ASN	2.6
1	A	659	LYS	2.5
1	A	462	ASN	2.5
1	A	625	HIS	2.5
1	A	583	LEU	2.4
1	A	554	HIS	2.4
1	A	660	LEU	2.4
1	A	534	TYR	2.3
1	A	517	CYS	2.3
1	A	380	ASN	2.3
1	A	621	ARG	2.3
1	A	106	SER	2.2
1	A	619	ILE	2.1
1	A	576	LYS	2.1
1	A	308	HIS	2.1
1	A	474	GLY	2.1
1	A	446	PHE	2.1
1	A	653	ASP	2.0
1	A	472	ALA	2.0
1	A	610	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	704	6/6	0.85	0.21	69,73,77,83	0
5	BCT	A	705	4/4	0.89	0.15	62,68,77,77	0
4	GOL	A	703	6/6	0.95	0.10	39,45,47,51	0
3	MLI	A	702	7/7	0.96	0.07	42,51,54,58	0
2	CR	A	701	1/1	1.00	0.02	55,55,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.