



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:48 PM UTC

PDB ID : 6UNR / pdb\_00006unr  
Title : Kinase domain of ALK2-K492A/K493A with AMPPNP  
Authors : Agnew, C.; Jura, N.  
Deposited on : 2019-10-13  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Activin receptor type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2185	1394	369	408	14	0	0	0

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	170	MET	-	initiating methionine	UNP Q04771
A	171	SER	-	expression tag	UNP Q04771
A	172	TYR	-	expression tag	UNP Q04771
A	173	TYR	-	expression tag	UNP Q04771
A	174	HIS	-	expression tag	UNP Q04771
A	175	HIS	-	expression tag	UNP Q04771
A	176	HIS	-	expression tag	UNP Q04771
A	177	HIS	-	expression tag	UNP Q04771
A	178	HIS	-	expression tag	UNP Q04771
A	179	HIS	-	expression tag	UNP Q04771
A	180	HIS	-	expression tag	UNP Q04771
A	181	HIS	-	expression tag	UNP Q04771
A	182	HIS	-	expression tag	UNP Q04771
A	183	HIS	-	expression tag	UNP Q04771
A	184	ASP	-	expression tag	UNP Q04771
A	185	TYR	-	expression tag	UNP Q04771
A	186	ASP	-	expression tag	UNP Q04771
A	187	ILE	-	expression tag	UNP Q04771
A	188	PRO	-	expression tag	UNP Q04771
A	189	THR	-	expression tag	UNP Q04771
A	190	THR	-	expression tag	UNP Q04771
A	191	GLU	-	expression tag	UNP Q04771
A	192	ASN	-	expression tag	UNP Q04771
A	193	LEU	-	expression tag	UNP Q04771
A	194	TYR	-	expression tag	UNP Q04771
A	195	PHE	-	expression tag	UNP Q04771
A	196	GLN	-	expression tag	UNP Q04771

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	197	GLY	-	expression tag	UNP Q04771
A	198	ALA	-	expression tag	UNP Q04771
A	199	MET	-	expression tag	UNP Q04771
A	200	GLY	-	expression tag	UNP Q04771
A	492	ALA	LYS	conflict	UNP Q04771
A	493	ALA	LYS	conflict	UNP Q04771

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	31	10	6	12	3	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	8	8	8	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	12	12	12	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Activin receptor type-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.12Å 87.12Å 138.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.14 – 2.20 46.14 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.14-2.20) 86.4 (46.14-2.20)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.16_3549, PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.243 , 0.302 0.244 , 0.304	Depositor DCC
$R_{free}$ test set	889 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtrriage
Anisotropy	1.004	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2236	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.17	0/2232	0.46	2/3043 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	271	THR	CA-C-N	5.39	131.84	121.54
1	A	271	THR	C-N-CA	5.39	131.84	121.54

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2185	0	2084	26	0
2	A	31	0	13	0	0
3	A	8	0	0	0	0
4	A	12	0	0	0	0
All	All	2236	0	2097	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:VAL:HG13	1:A:330:PRO:HD2	1.80	0.63
1:A:469:SER:HA	1:A:472:LYS:HB2	1.81	0.62
1:A:260:GLU:O	1:A:345:LYS:NZ	2.25	0.62
1:A:473:LEU:HD21	1:A:494:THR:HG21	1.90	0.54
1:A:302:THR:HG22	1:A:418:MET:HE3	1.90	0.54
1:A:269:ASP:OD1	1:A:271:THR:HG23	2.08	0.52
1:A:261:ASN:HD22	1:A:314:SER:HB2	1.74	0.52
1:A:335:ARG:HD2	1:A:359:VAL:HG13	1.91	0.52
1:A:332:ILE:HG12	1:A:360:MET:HG2	1.92	0.51
1:A:288:MET:HE3	1:A:344:VAL:HG12	1.94	0.49
1:A:261:ASN:ND2	1:A:314:SER:HB2	2.29	0.48
1:A:272:SER:OG	1:A:273:ARG:N	2.47	0.48
1:A:273:ARG:O	1:A:276:SER:N	2.47	0.48
1:A:419:VAL:HG12	1:A:424:VAL:CG1	2.44	0.48
1:A:357:LEU:HD11	1:A:375:ARG:HG2	1.97	0.47
1:A:463:SER:O	1:A:463:SER:OG	2.31	0.47
1:A:335:ARG:HD3	1:A:357:LEU:O	2.15	0.47
1:A:461:TRP:HB3	1:A:467:LEU:HB3	1.98	0.46
1:A:255:VAL:HG11	1:A:332:ILE:HD11	1.98	0.45
1:A:431:PHE:O	1:A:435:VAL:HG22	2.18	0.44
1:A:412:TRP:CZ2	1:A:416:ARG:HD2	2.53	0.43
1:A:376:VAL:HG12	1:A:382:MET:HE2	2.00	0.42
1:A:476:GLU:HB3	1:A:486:LEU:HD13	2.01	0.42
1:A:434:VAL:HG13	1:A:435:VAL:HG13	2.01	0.42
1:A:288:MET:HE2	1:A:346:LYS:HA	2.01	0.42
1:A:459:ASN:OD1	1:A:459:ASN:N	2.54	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	280/330 (85%)	260 (93%)	18 (6%)	2 (1%)	18 19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	272	SER
1	A	434	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	229/294 (78%)	229 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	261	ASN
1	A	318	HIS
1	A	347	ASN
1	A	361	HIS
1	A	480	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ANP	A	501	-	33,33,33	0.91	3 (9%)	45,52,52	0.76	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	501	-	-	4/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ANP	PG-O1G	2.63	1.50	1.46
2	A	501	ANP	PG-N3B	2.44	1.69	1.63
2	A	501	ANP	PB-O1B	2.42	1.49	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ANP	O1B-PB-N3B	-3.56	106.52	111.77

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	ANP	PB-N3B-PG-O1G
2	A	501	ANP	O4'-C4'-C5'-O5'
2	A	501	ANP	C3'-C4'-C5'-O5'
2	A	501	ANP	PB-O3A-PA-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/330 (86%)	0.82	27 (9%) <b>14</b> <b>11</b>	41, 73, 108, 129	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	227	TRP	4.7
1	A	274	HIS	4.7
1	A	208	ILE	4.3
1	A	276	SER	3.9
1	A	211	LEU	3.8
1	A	217	GLY	3.1
1	A	219	TYR	3.1
1	A	277	THR	3.0
1	A	496	THR	2.9
1	A	441	PHE	2.8
1	A	272	SER	2.6
1	A	450	VAL	2.5
1	A	468	THR	2.5
1	A	462	PHE	2.4
1	A	231	ASN	2.4
1	A	421	ASN	2.3
1	A	350	CYS	2.2
1	A	230	GLU	2.2
1	A	465	PRO	2.2
1	A	326	THR	2.2
1	A	424	VAL	2.1
1	A	471	ALA	2.1
1	A	391	ILE	2.1
1	A	437	ASN	2.1
1	A	302	THR	2.1
1	A	310	LEU	2.0
1	A	270	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	504	1/1	0.65	0.22	112,112,112,112	0
3	MG	A	508	1/1	0.79	0.12	62,62,62,62	0
2	ANP	A	501	31/31	0.80	0.14	56,82,132,138	0
3	MG	A	505	1/1	0.83	0.14	84,84,84,84	0
3	MG	A	506	1/1	0.85	0.14	63,63,63,63	0
3	MG	A	507	1/1	0.89	0.13	70,70,70,70	0
3	MG	A	503	1/1	0.89	0.10	76,76,76,76	0
3	MG	A	502	1/1	0.91	0.16	65,65,65,65	0
3	MG	A	509	1/1	0.93	0.14	67,67,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.