



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:40 AM UTC

PDB ID : 6UXG / pdb\_00006uxg  
Title : Structure of *V. metoecus* NucC, trimer form  
Authors : Ye, Q.; Corbett, K.D.  
Deposited on : 2019-11-07  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	 12% 74% 18% • 6%
1	B	245	 10% 76% 18% • 6%
1	C	245	 9% 75% 18% • 6%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10916 atoms, of which 5331 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called *Vibrio metoecus* sp. RC341 NucC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	230	Total 3611	C 1172	H 1777	N 313	O 341	S 8	0	0	0
1	B	230	Total 3611	C 1172	H 1777	N 313	O 341	S 8	0	0	0
1	C	230	Total 3611	C 1172	H 1777	N 313	O 341	S 8	0	0	0

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	17	Total O 17 17	0	0
3	B	18	Total O 18 18	0	0
3	C	18	Total O 18 18	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: *Vibrio metoecus* sp. RC341 NucC



- Molecule 1: *Vibrio metoecus* sp. RC341 NucC



- Molecule 1: *Vibrio metoecus* sp. RC341 NucC



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.61Å 123.61Å 123.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	87.51 – 2.15 87.51 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.8 (87.51-2.15) 99.3 (87.51-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.17_3644	Depositor
R, $R_{free}$	0.225 , 0.239 0.235 , 0.248	Depositor DCC
$R_{free}$ test set	2615 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.4	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 60.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.487 for -h,l,k 0.487 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10916	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/1874	0.63	0/2544
1	B	0.37	0/1874	0.63	0/2544
1	C	0.38	0/1874	0.61	0/2544
All	All	0.37	0/5622	0.62	0/7632

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1834	1777	1776	32	0
1	B	1834	1777	1776	33	0
1	C	1834	1777	1776	32	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
3	A	17	0	0	0	0
3	B	18	0	0	0	0
3	C	18	0	0	0	0
All	All	5585	5331	5328	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:THR:HG22	1:B:188:ASP:OD2	1.74	0.86
1:A:23:THR:HG22	1:A:188:ASP:OD2	1.77	0.85
1:C:23:THR:HG22	1:C:188:ASP:OD2	1.80	0.80
1:B:20:HIS:O	1:B:24:THR:HG23	1.96	0.66
1:B:79:ASP:O	1:B:83:SER:HB2	1.96	0.66
1:A:20:HIS:O	1:A:24:THR:HG23	1.96	0.65
1:C:20:HIS:O	1:C:24:THR:HG23	1.98	0.63
1:C:50:TYR:CD2	1:C:214:LEU:HD13	2.32	0.63
1:A:50:TYR:CD2	1:A:214:LEU:HD13	2.34	0.62
1:C:79:ASP:O	1:C:83:SER:HB2	2.01	0.61
1:A:79:ASP:O	1:A:83:SER:HB2	2.02	0.60
1:B:167:LEU:HD22	1:B:203:VAL:HG21	1.82	0.60
1:B:50:TYR:CD2	1:B:214:LEU:HD13	2.36	0.59
1:A:167:LEU:HD22	1:A:203:VAL:HG21	1.86	0.57
1:C:56:ARG:HD3	1:C:78:TYR:OH	2.05	0.57
1:A:56:ARG:HD3	1:A:78:TYR:OH	2.05	0.56
1:C:50:TYR:CG	1:C:214:LEU:HD13	2.41	0.56
1:C:167:LEU:C	1:C:167:LEU:HD23	2.30	0.56
1:B:80:ARG:NH2	1:C:53:GLU:OE2	2.38	0.56
1:C:167:LEU:HD22	1:C:203:VAL:HG21	1.89	0.55
1:C:174:ASP:OD1	1:C:175:LYS:N	2.39	0.55
1:B:180:MET:HE1	1:B:194:TYR:CD1	2.41	0.55
1:A:167:LEU:C	1:A:167:LEU:HD23	2.32	0.55
1:A:80:ARG:HH11	1:A:80:ARG:HG3	1.73	0.54
1:B:173:HIS:NE2	1:B:177:ASP:OD2	2.41	0.54
1:B:56:ARG:HD3	1:B:78:TYR:OH	2.08	0.53
1:A:50:TYR:CG	1:A:214:LEU:HD13	2.44	0.53
1:A:167:LEU:HD21	1:A:171:LEU:HD12	1.91	0.53
1:C:180:MET:HE1	1:C:194:TYR:CD1	2.44	0.52
1:B:167:LEU:C	1:B:167:LEU:HD23	2.34	0.52
1:A:180:MET:HE1	1:A:194:TYR:CD1	2.44	0.52
1:C:161:LEU:O	1:C:163:ILE:N	2.44	0.51
1:A:173:HIS:NE2	1:A:177:ASP:OD2	2.44	0.51
1:C:167:LEU:HD21	1:C:171:LEU:HD12	1.92	0.50
1:B:80:ARG:HG3	1:B:80:ARG:HH11	1.75	0.50
1:B:80:ARG:HG3	1:B:80:ARG:NH1	2.27	0.50
1:B:107:LYS:O	1:B:156:THR:HA	2.12	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:ILE:H	1:B:163:ILE:HD12	1.77	0.49
1:B:229:MET:HE3	1:C:222:GLN:HG2	1.93	0.49
1:B:166:THR:O	1:B:170:HIS:ND1	2.45	0.49
1:B:167:LEU:HD21	1:B:171:LEU:HD12	1.95	0.49
1:B:229:MET:HE3	1:C:222:GLN:CG	2.43	0.49
1:A:87:PHE:HB2	1:A:94:ILE:HB	1.93	0.49
1:A:163:ILE:H	1:A:163:ILE:HD12	1.77	0.48
1:C:227:VAL:HG13	1:C:228:PRO:O	2.13	0.48
1:C:166:THR:O	1:C:170:HIS:ND1	2.43	0.48
1:C:192:PHE:CD1	1:C:201:MET:HE1	2.48	0.48
1:A:204:MET:HE2	1:C:240:THR:HB	1.95	0.48
1:B:50:TYR:CG	1:B:214:LEU:HD13	2.48	0.48
1:B:192:PHE:CD1	1:B:201:MET:HE1	2.49	0.48
1:A:80:ARG:HG3	1:A:80:ARG:NH1	2.28	0.47
1:A:192:PHE:CD1	1:A:201:MET:HE1	2.50	0.47
1:B:50:TYR:CE2	1:B:214:LEU:HD13	2.49	0.47
1:A:50:TYR:CE2	1:A:214:LEU:HD13	2.50	0.47
1:A:240:THR:HB	1:B:204:MET:HE2	1.96	0.47
1:B:240:THR:HB	1:C:204:MET:HE2	1.97	0.47
1:A:107:LYS:O	1:A:156:THR:HA	2.15	0.47
1:B:229:MET:HE3	1:C:222:GLN:CD	2.40	0.47
1:C:50:TYR:CE2	1:C:214:LEU:HD13	2.49	0.46
1:C:107:LYS:O	1:C:156:THR:HA	2.16	0.46
1:A:199:GLN:O	1:A:199:GLN:HG2	2.16	0.46
1:A:161:LEU:HD22	1:A:166:THR:HG22	1.98	0.46
1:A:166:THR:O	1:A:170:HIS:ND1	2.48	0.45
1:B:199:GLN:HG2	1:B:199:GLN:O	2.17	0.45
1:C:135:ILE:O	1:C:142:HIS:N	2.46	0.45
1:A:174:ASP:OD1	1:A:175:LYS:N	2.49	0.44
1:B:161:LEU:HD22	1:B:166:THR:HG22	1.98	0.44
1:C:163:ILE:HD12	1:C:163:ILE:H	1.82	0.44
1:C:199:GLN:HG2	1:C:199:GLN:O	2.17	0.44
1:A:167:LEU:CD2	1:A:171:LEU:HD12	2.48	0.43
1:A:52:PRO:HG2	1:A:55:TYR:CE2	2.53	0.43
1:A:227:VAL:HG13	1:A:228:PRO:O	2.19	0.43
1:B:6:GLN:O	1:B:9:GLU:N	2.51	0.43
1:A:135:ILE:O	1:A:142:HIS:N	2.51	0.42
1:B:94:ILE:N	1:B:94:ILE:HD12	2.34	0.42
1:C:87:PHE:HB2	1:C:94:ILE:HB	2.01	0.42
1:B:174:ASP:OD1	1:B:175:LYS:N	2.52	0.42
1:A:6:GLN:O	1:A:9:GLU:N	2.53	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:TYR:O	1:B:90:ALA:HB3	2.19	0.41
1:B:147:LEU:HD23	1:B:147:LEU:HA	1.92	0.41
1:A:4:ASP:HB3	1:C:89:TYR:HE2	1.85	0.41
1:A:89:TYR:CE2	1:B:4:ASP:HB3	2.55	0.41
1:C:89:TYR:CD1	1:C:89:TYR:C	2.98	0.41
1:B:25:VAL:HG21	1:B:42:VAL:HG21	2.02	0.41
1:A:25:VAL:HG21	1:A:42:VAL:HG21	2.02	0.40
1:C:93:LEU:HD12	1:C:93:LEU:HA	1.93	0.40
1:C:161:LEU:HD22	1:C:166:THR:HG22	2.03	0.40
1:C:110:LEU:HB3	1:C:159:ASN:HA	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/245 (91%)	216 (96%)	8 (4%)	0	100	100
1	B	224/245 (91%)	215 (96%)	9 (4%)	0	100	100
1	C	224/245 (91%)	216 (96%)	8 (4%)	0	100	100
All	All	672/735 (91%)	647 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/214 (92%)	190 (97%)	6 (3%)	35	37
1	B	196/214 (92%)	192 (98%)	4 (2%)	48	54
1	C	196/214 (92%)	188 (96%)	8 (4%)	27	26
All	All	588/642 (92%)	570 (97%)	18 (3%)	35	37

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	SER
1	A	80	ARG
1	A	173	HIS
1	A	226	THR
1	A	227	VAL
1	A	229	MET
1	B	8	SER
1	B	173	HIS
1	B	226	THR
1	B	227	VAL
1	C	8	SER
1	C	80	ARG
1	C	83	SER
1	C	86	ILE
1	C	173	HIS
1	C	226	THR
1	C	227	VAL
1	C	229	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	92	GLN
1	B	68	GLN
1	B	81	GLN
1	B	129	HIS
1	C	68	GLN
1	C	81	GLN
1	C	92	GLN
1	C	206	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	301	-	4,4,4	0.30	0	6,6,6	0.28	0
2	SO4	C	302	-	4,4,4	0.23	0	6,6,6	0.30	0
2	SO4	A	301	-	4,4,4	0.26	0	6,6,6	0.38	0
2	SO4	B	302	-	4,4,4	0.30	0	6,6,6	0.48	0
2	SO4	C	301	-	4,4,4	0.23	0	6,6,6	0.30	0
2	SO4	A	302	-	4,4,4	0.23	0	6,6,6	0.40	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/245 (93%)	0.60	29 (12%) <b>8</b> <b>8</b>	50, 76, 157, 200	0
1	B	230/245 (93%)	0.61	24 (10%) <b>11</b> <b>13</b>	50, 77, 155, 200	0
1	C	230/245 (93%)	0.53	23 (10%) <b>12</b> <b>14</b>	49, 77, 156, 206	0
All	All	690/735 (93%)	0.58	76 (11%) <b>10</b> <b>12</b>	49, 77, 156, 206	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	VAL	7.1
1	B	163	ILE	5.7
1	C	176	ALA	4.9
1	A	163	ILE	4.6
1	C	197	ASP	4.5
1	B	141	VAL	4.4
1	A	175	LYS	4.1
1	A	29	PHE	4.0
1	C	136	PRO	4.0
1	A	164	PRO	3.9
1	A	176	ALA	3.9
1	A	142	HIS	3.9
1	A	162	LYS	3.8
1	C	174	ASP	3.8
1	C	30	LYS	3.8
1	C	141	VAL	3.8
1	C	173	HIS	3.8
1	A	136	PRO	3.8
1	C	142	HIS	3.6
1	B	197	ASP	3.5
1	C	198	HIS	3.4
1	C	163	ILE	3.4
1	B	136	PRO	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	4	ASP	3.3
1	B	242	ARG	3.2
1	B	162	LYS	3.1
1	B	176	ALA	3.1
1	B	164	PRO	3.1
1	B	165	ASP	3.0
1	A	23	THR	3.0
1	A	167	LEU	3.0
1	C	135	ILE	2.9
1	B	226	THR	2.9
1	B	142	HIS	2.9
1	B	172	ASP	2.8
1	A	198	HIS	2.8
1	C	112	LYS	2.8
1	A	135	ILE	2.8
1	B	143	SER	2.7
1	A	197	ASP	2.7
1	B	168	MET	2.7
1	A	165	ASP	2.7
1	A	173	HIS	2.6
1	A	242	ARG	2.6
1	A	27	LYS	2.6
1	B	135	ILE	2.6
1	A	26	ARG	2.6
1	C	113	GLN	2.5
1	A	30	LYS	2.4
1	A	171	LEU	2.4
1	B	50	TYR	2.4
1	C	175	LYS	2.4
1	C	167	LEU	2.3
1	A	172	ASP	2.3
1	A	201	MET	2.3
1	C	162	LYS	2.3
1	A	5	TRP	2.3
1	C	143	SER	2.3
1	C	29	PHE	2.3
1	A	143	SER	2.2
1	C	196	ASN	2.2
1	A	226	THR	2.2
1	A	170	HIS	2.2
1	B	23	THR	2.2
1	B	3	GLN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	5	TRP	2.1
1	A	227	VAL	2.1
1	A	22	LEU	2.1
1	C	164	PRO	2.1
1	C	226	THR	2.1
1	B	173	HIS	2.1
1	B	230	ILE	2.1
1	B	4	ASP	2.1
1	B	89	TYR	2.1
1	C	166	THR	2.1
1	B	51	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	301	5/5	0.92	0.13	142,143,144,144	0
2	SO4	B	302	5/5	0.92	0.12	195,196,196,197	0
2	SO4	C	301	5/5	0.92	0.14	145,146,146,147	0
2	SO4	C	302	5/5	0.93	0.14	206,207,208,208	0
2	SO4	A	301	5/5	0.94	0.13	150,151,151,151	0
2	SO4	A	302	5/5	0.95	0.12	189,190,190,191	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.