



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:51 PM UTC

PDB ID : 6UY8 / pdb_00006uy8
Title : Crystal structure of the STAC3 tandem SH3 domains - P269R, K329N
Authors : Rufenach, B.; Van Petegem, F.
Deposited on : 2019-11-12
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	123	 80% 19% .
1	B	123	 67% 24% . 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3782 atoms, of which 1700 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SH3 and cysteine-rich domain-containing protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	123	Total	C	H	N	O	0	0	0
			1864	624	896	171	173			
1	B	115	Total	C	H	N	O	0	0	0
			1651	568	774	150	159			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	242	SER	-	expression tag	UNP Q96MF2
A	243	ASN	-	expression tag	UNP Q96MF2
A	244	ALA	-	expression tag	UNP Q96MF2
A	269	ARG	PRO	engineered mutation	UNP Q96MF2
A	329	ASN	LYS	engineered mutation	UNP Q96MF2
B	242	SER	-	expression tag	UNP Q96MF2
B	243	ASN	-	expression tag	UNP Q96MF2
B	244	ALA	-	expression tag	UNP Q96MF2
B	269	ARG	PRO	engineered mutation	UNP Q96MF2
B	329	ASN	LYS	engineered mutation	UNP Q96MF2

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	123	Total	O	0	4
			127	127		
4	B	81	Total	O	0	8
			89	89		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SH3 and cysteine-rich domain-containing protein 3

Chain A:  80% 19% .



- Molecule 1: SH3 and cysteine-rich domain-containing protein 3

Chain B:  67% 24% . 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	87.88Å 87.88Å 29.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.77 – 1.65 28.77 – 1.65	Depositor EDS
% Data completeness (in resolution range)	95.8 (28.77-1.65) 95.8 (28.77-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.65Å)	Xtriage
Refinement program	PHENIX 1.15.1_3469	Depositor
R, R_{free}	0.187 , 0.208 0.191 , 0.205	Depositor DCC
R_{free} test set	1437 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 36.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.174 for -h,-k,l 0.067 for h,-h-k,-l 0.059 for -k,-h,-l	Xtriage
Reported twinning fraction	0.200 for -h,-k,l	Depositor
Outliers	0 of 29433 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3782	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.68	17/991 (1.7%)	1.24	0/1339
1	B	1.66	20/899 (2.2%)	1.29	1/1222 (0.1%)
All	All	1.67	37/1890 (2.0%)	1.26	1/2561 (0.0%)

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	297	PRO	C-O	-8.94	1.14	1.24
1	B	357	PRO	C-O	-8.91	1.14	1.23
1	B	297	PRO	C-O	-8.54	1.15	1.24
1	A	347	VAL	C-O	-8.23	1.15	1.24
1	B	293	VAL	C-O	-7.93	1.16	1.24
1	A	298	PRO	C-O	-7.87	1.14	1.24
1	A	357	PRO	C-O	-7.74	1.15	1.23
1	A	293	VAL	C-O	-7.24	1.16	1.24
1	A	270	PRO	C-O	-7.23	1.15	1.23
1	B	281	ASN	C-O	-6.63	1.15	1.24
1	B	270	PRO	C-O	-6.62	1.16	1.23
1	B	304	VAL	C-O	-6.44	1.17	1.24
1	A	269	ARG	C-O	-6.44	1.17	1.24
1	B	286	ARG	C-O	-6.43	1.16	1.24
1	B	253	VAL	C-O	-6.42	1.17	1.24
1	B	295	PHE	C-O	-6.26	1.16	1.23
1	A	334	VAL	C-O	-6.16	1.17	1.23
1	B	266	LEU	C-O	-6.05	1.16	1.24
1	A	303	ARG	C-O	-6.01	1.16	1.23
1	B	298	PRO	C-O	-5.79	1.17	1.24
1	B	260	ALA	C-O	-5.78	1.17	1.23
1	B	284	TRP	C-O	-5.68	1.17	1.24
1	B	292	LYS	C-O	-5.64	1.17	1.23
1	A	284	TRP	C-O	-5.62	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	278	ASP	C-O	-5.52	1.17	1.23
1	B	300	PHE	C-O	-5.38	1.16	1.24
1	A	274	ILE	C-O	-5.36	1.18	1.24
1	B	358	THR	C-O	-5.35	1.17	1.24
1	A	272	GLU	C-O	-5.30	1.17	1.23
1	B	342	GLY	C-O	-5.30	1.16	1.23
1	B	335	VAL	C-O	-5.29	1.18	1.24
1	A	280	SER	C-O	-5.27	1.17	1.24
1	A	295	PHE	C-O	-5.25	1.17	1.23
1	B	314	THR	C-O	-5.23	1.17	1.24
1	A	313	VAL	C-O	-5.19	1.18	1.24
1	A	335	VAL	C-O	-5.11	1.18	1.24
1	B	346	LYS	C-O	-5.04	1.17	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	264	ASP	CA-CB-CG	5.31	117.91	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	968	896	923	4	0
1	B	877	774	795	10	0
2	A	12	18	17	0	0
2	B	8	12	12	0	0
3	B	1	0	0	0	0
4	A	127	0	0	0	0
4	B	89	0	0	1	0
All	All	2082	1700	1747	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:268:PHE:CD2	1:B:274:ILE:HD11	2.01	0.95
1:B:268:PHE:CD2	1:B:274:ILE:CD1	2.75	0.68
1:B:326:ILE:HG23	1:B:354:GLY:HA3	1.95	0.48
1:B:250:HIS:C	1:B:250:HIS:ND1	2.71	0.48
1:A:272:GLU:OE2	1:A:289:ILE:HD11	2.14	0.47
1:B:305:ARG:NH1	4:B:574[B]:HOH:O	2.47	0.47
1:A:330:LYS:O	1:A:331:ASP:HB2	2.17	0.44
1:A:326:ILE:HG23	1:A:354:GLY:HA3	2.00	0.43
1:B:345:VAL:HG13	1:B:358:THR:HG22	2.01	0.42
1:B:305:ARG:O	1:B:308:GLU:HB2	2.18	0.42
1:B:288:LYS:HA	1:B:292:LYS:O	2.20	0.42
1:A:305:ARG:O	1:A:308:GLU:HB2	2.20	0.42
1:B:262:GLU:HB2	1:B:265:ASP:CG	2.44	0.42
1:B:268:PHE:CE2	1:B:274:ILE:CD1	3.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/123 (98%)	119 (98%)	2 (2%)	0	100	100
1	B	113/123 (92%)	111 (98%)	2 (2%)	0	100	100
All	All	234/246 (95%)	230 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/105 (90%)	95 (100%)	0	100	100
1	B	80/105 (76%)	80 (100%)	0	100	100
All	All	175/210 (83%)	175 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	299	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	403	-	3,3,3	0.73	0	2,2,2	0.20	0
2	EDO	B	402	-	3,3,3	0.38	0	2,2,2	0.22	0
2	EDO	A	401	-	3,3,3	0.77	0	2,2,2	0.15	0
2	EDO	B	401	-	3,3,3	0.72	0	2,2,2	0.33	0
2	EDO	A	402	-	3,3,3	0.47	0	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	403	-	-	0/1/1/1	-
2	EDO	B	402	-	-	0/1/1/1	-
2	EDO	A	401	-	-	0/1/1/1	-
2	EDO	B	401	-	-	0/1/1/1	-
2	EDO	A	402	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	402	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	123/123 (100%)	-1.19	0 100 100	15, 24, 40, 52	0
1	B	115/123 (93%)	-1.11	0 100 100	18, 28, 42, 53	0
All	All	238/246 (96%)	-1.15	0 100 100	15, 26, 42, 53	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	402	4/4	0.98	0.05	25,30,31,32	0
2	EDO	B	402	4/4	0.98	0.06	36,43,44,46	0
2	EDO	A	403	4/4	0.99	0.04	23,27,28,29	0
3	CL	B	403	1/1	0.99	0.04	46,46,46,46	0
2	EDO	A	401	4/4	1.00	0.02	15,18,18,19	0
2	EDO	B	401	4/4	1.00	0.02	18,21,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.