



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 09:10 AM UTC

PDB ID : 6UYD / pdb_00006uyd
Title : Structure of Hepatitis C Virus Envelope Glycoprotein E2mc3-v1 redesigned core from genotype 1a bound to broadly neutralizing antibody AR3C
Authors : Tzarum, N.; Wilson, I.A.; Zhu, J.
Deposited on : 2019-11-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	177	
1	F	177	
2	A	233	
2	H	233	
3	B	214	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	L	214	 <p>% 94% 5%</p>
4	C	2	 <p>50% 50%</p>
4	D	2	 <p>50% 50%</p>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9525 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein E2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	143	Total	C	N	O	S	0	0	0
			1118	719	189	197	13			
1	F	158	Total	C	N	O	S	0	0	0
			1230	782	210	225	13			

- Molecule 2 is a protein called Fab AR3C heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	222	Total	C	N	O	S	0	0	0
			1659	1044	281	327	7			
2	A	222	Total	C	N	O	S	0	0	0
			1656	1042	280	326	8			

- Molecule 3 is a protein called Fab AR3C light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	211	Total	C	N	O	S	0	0	0
			1612	1005	277	326	4			
3	B	212	Total	C	N	O	S	0	0	0
			1618	1008	278	327	5			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	2	28	16	2	10	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	E	1	14	8	1	5	0	0
5	E	1	14	8	1	5	0	0
5	E	1	14	8	1	5	0	0
5	F	1	14	8	1	5	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	E	38	38	38	0	0
6	H	134	134	134	0	0
6	L	104	104	104	0	0
6	A	100	100	100	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	57	Total O 57 57	0	0
6	F	87	Total O 87 87	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Envelope glycoprotein E2




- Molecule 1: Envelope glycoprotein E2



- Molecule 2: Fab AR3C heavy chain





- Molecule 2: Fab AR3C heavy chain







- Molecule 3: Fab AR3C light chain



- Molecule 3: Fab AR3C light chain



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.49Å 93.66Å 96.80Å 90.00° 100.48° 90.00°	Depositor
Resolution (Å)	29.91 – 1.90 29.91 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (29.91-1.90) 98.7 (29.91-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.208 , 0.243 0.209 , 0.245	Depositor DCC
R_{free} test set	6108 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	35.2	Xtrriage
Anisotropy	0.517	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9525	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.40	0/1156	0.61	0/1580
1	F	0.39	0/1273	0.61	2/1744 (0.1%)
2	A	0.37	0/1695	0.54	0/2306
2	H	0.41	0/1698	0.58	0/2310
3	B	0.26	0/1652	0.48	0/2243
3	L	0.36	0/1646	0.60	0/2235
All	All	0.37	0/9120	0.57	2/12418 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	422	ILE	CA-C-N	-5.14	113.72	122.56
1	F	422	ILE	C-N-CA	-5.14	113.72	122.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1118	0	1037	15	0
1	F	1230	0	1124	15	0
2	A	1656	0	1610	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1659	0	1614	9	0
3	B	1618	0	1574	9	0
3	L	1612	0	1570	5	0
4	C	28	0	25	1	0
4	D	28	0	25	1	0
5	E	42	0	39	1	0
5	F	14	0	13	0	0
6	A	100	0	0	0	0
6	B	57	0	0	1	0
6	E	38	0	0	1	0
6	F	87	0	0	3	0
6	H	134	0	0	2	0
6	L	104	0	0	0	0
All	All	9525	0	8631	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:41:PRO:O	2:A:43:GLN:NE2	2.03	0.92
1:F:429:CYS:HB3	1:F:438:LEU:HD21	1.67	0.75
1:E:604:THR:HG22	1:E:606:ARG:H	1.52	0.74
1:F:450:SER:HB3	1:F:453:GLN:HB2	1.70	0.73
3:L:142:ARG:NH2	3:L:163:VAL:HG11	2.09	0.67
1:F:434:ASN:ND2	6:F:802:HOH:O	2.28	0.66
3:B:19:ALA:HB2	3:B:78:LEU:HD11	1.75	0.66
3:B:120:PRO:HD3	3:B:132:VAL:HG22	1.81	0.63
1:E:500:LYS:NZ	1:E:531:ALA:O	2.32	0.62
1:E:604:THR:HB	1:E:607:CYS:HB2	1.83	0.61
1:E:423:ASN:HB3	1:E:529:TRP:CE2	2.37	0.59
1:E:557:SER:HB2	4:C:1:NAG:H82	1.84	0.59
2:A:199:ASN:ND2	2:A:206:LYS:HG2	2.18	0.59
2:A:123:PRO:HD3	2:A:209:LYS:NZ	2.18	0.58
2:H:58:ARG:NH2	6:H:303:HOH:O	2.28	0.57
1:F:582:THR:N	6:F:804:HOH:O	2.38	0.57
1:F:582:THR:N	6:F:805:HOH:O	2.39	0.56
1:E:628:LYS:HE3	2:A:192:GLN:HA	1.87	0.55
1:E:540:ASN:ND2	1:E:540:ASN:O	2.39	0.55
2:H:40:ALA:HB3	2:H:43:GLN:HB2	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:52(A):PRO:HB2	2:A:73:LYS:HE2	1.91	0.53
1:E:510:THR:HG22	1:E:512:SER:O	2.09	0.52
1:F:519:THR:HG22	1:F:525:PRO:HA	1.91	0.52
3:B:210:ASN:O	3:B:213:GLU:HG2	2.10	0.52
1:E:423:ASN:HB3	1:E:529:TRP:CZ2	2.45	0.52
3:L:89:GLN:HG3	3:L:98:PHE:CE2	2.47	0.49
2:H:4:LEU:HB2	2:H:104:GLY:HA2	1.95	0.49
5:E:702:NAG:H83	5:E:702:NAG:H3	1.95	0.49
3:B:142:ARG:NH1	3:B:163:VAL:HG11	2.28	0.49
1:F:509:PHE:HE1	1:F:553:THR:HG22	1.78	0.49
1:F:557:SER:HB2	4:D:1:NAG:H82	1.95	0.48
1:E:433:LEU:HD23	1:E:433:LEU:HA	1.70	0.47
2:A:99:ARG:NH1	1:F:431:GLU:OE1	2.38	0.47
2:A:199:ASN:HD22	2:A:206:LYS:HG2	1.80	0.46
2:A:41:PRO:O	2:A:43:GLN:CD	2.58	0.46
1:F:423:ASN:HB3	1:F:529:TRP:CZ2	2.52	0.45
1:F:599:SER:O	1:F:602:TRP:NE1	2.49	0.45
1:E:636:VAL:HA	6:E:814:HOH:O	2.17	0.45
2:A:52(A):PRO:HG3	2:A:71:ASP:HB3	1.99	0.45
2:H:191:THR:HG22	1:F:628:LYS:HB3	1.98	0.45
2:A:20:VAL:HG12	2:A:80:MET:HE2	1.99	0.44
1:E:509:PHE:CD1	1:E:601:PRO:HB3	2.53	0.44
2:H:39:GLN:HG3	2:H:44:GLY:O	2.17	0.44
2:A:123:PRO:HD3	2:A:209:LYS:HZ1	1.82	0.44
1:F:500:LYS:HD2	1:F:533:ASP:OD1	2.17	0.44
3:B:213:GLU:O	3:B:213:GLU:HG3	2.17	0.43
2:H:100(C):GLY:HA2	2:H:100(D):GLY:HA2	1.54	0.43
2:A:87:THR:HG23	2:A:110:THR:HA	2.01	0.43
3:B:36:TYR:HB2	3:B:87:PHE:CE1	2.53	0.43
3:L:24:ARG:HD3	3:L:70:GLU:OE2	2.18	0.43
2:A:20:VAL:CG1	2:A:80:MET:HE2	2.48	0.43
1:E:432:SER:OG	1:E:433:LEU:N	2.52	0.43
3:L:125:LEU:O	3:L:183:LYS:HD2	2.19	0.43
3:L:48:ILE:HD13	3:L:54:ARG:HA	2.01	0.43
2:A:33:ALA:HB2	2:A:52:VAL:HG22	2.00	0.43
2:A:100(C):GLY:HA3	1:F:529:TRP:HB2	2.00	0.43
1:F:510:THR:HG22	1:F:549:TRP:CD2	2.53	0.42
2:H:87:THR:HA	2:H:109:VAL:O	2.18	0.42
2:A:47:TRP:CE2	3:B:96:LEU:HD13	2.54	0.42
1:E:526:THR:HG21	1:E:535:ASP:OD1	2.20	0.42
3:B:96:LEU:HD12	3:B:96:LEU:H	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:86:ASP:OD1	6:H:301:HOH:O	2.21	0.41
2:H:178:LEU:C	2:H:178:LEU:HD12	2.45	0.41
2:A:100(C):GLY:HA2	2:A:100(D):GLY:HA2	1.52	0.41
1:E:631:MET:HE3	1:E:631:MET:HB2	1.91	0.41
2:A:119:PRO:HD2	2:A:205:THR:HG21	2.03	0.41
2:A:143:LYS:HE2	6:B:319:HOH:O	2.21	0.41
2:A:60:ALA:HB3	2:A:63:PHE:HD2	1.86	0.41
3:B:54:ARG:HD2	3:B:58:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	133/177 (75%)	129 (97%)	3 (2%)	1 (1%)	16	8
1	F	152/177 (86%)	148 (97%)	4 (3%)	0	100	100
2	A	218/233 (94%)	212 (97%)	6 (3%)	0	100	100
2	H	218/233 (94%)	209 (96%)	9 (4%)	0	100	100
3	B	210/214 (98%)	203 (97%)	7 (3%)	0	100	100
3	L	209/214 (98%)	203 (97%)	6 (3%)	0	100	100
All	All	1140/1248 (91%)	1104 (97%)	35 (3%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	510	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	122/149 (82%)	120 (98%)	2 (2%)	55	54
1	F	134/149 (90%)	132 (98%)	2 (2%)	57	56
2	A	186/195 (95%)	186 (100%)	0	100	100
2	H	186/195 (95%)	186 (100%)	0	100	100
3	B	182/184 (99%)	182 (100%)	0	100	100
3	L	181/184 (98%)	181 (100%)	0	100	100
All	All	991/1056 (94%)	987 (100%)	4 (0%)	84	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	431	GLU
1	E	540	ASN
1	F	443	TYR
1	F	522	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	444	GLN
2	H	39	GLN
2	H	59	ASN
3	L	38	GLN
3	L	147	GLN
3	L	160	GLN
3	B	210	ASN
1	F	428	ASN
1	F	430	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	1	1,4	14,14,15	0.18	0	17,19,21	0.63	1 (5%)
4	NAG	C	2	4	14,14,15	0.27	0	17,19,21	0.67	1 (5%)
4	NAG	D	1	1,4	14,14,15	0.49	0	17,19,21	0.38	0
4	NAG	D	2	4	14,14,15	0.32	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	2	NAG	C1-O5-C5	2.30	115.27	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1	NAG	C1-O5-C5	2.07	114.97	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	2	NAG	C4-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1	NAG	1	0
4	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	E	703	1	14,14,15	0.35	0	17,19,21	0.58	0
5	NAG	F	701	1	14,14,15	0.47	0	17,19,21	0.57	0
5	NAG	E	702	1	14,14,15	0.45	0	17,19,21	1.34	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	E	701	1	14,14,15	0.63	0	17,19,21	0.67	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	703	1	-	2/6/23/26	0/1/1/1
5	NAG	F	701	1	-	0/6/23/26	0/1/1/1
5	NAG	E	702	1	-	4/6/23/26	0/1/1/1
5	NAG	E	701	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	702	NAG	C2-N2-C7	4.73	129.24	122.90
5	E	701	NAG	C1-O5-C5	2.10	115.00	112.19
5	E	702	NAG	C1-C2-N2	2.05	113.66	110.43

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	701	NAG	O5-C5-C6-O6
5	E	703	NAG	O5-C5-C6-O6
5	E	703	NAG	C4-C5-C6-O6
5	E	701	NAG	C4-C5-C6-O6
5	E	702	NAG	C8-C7-N2-C2
5	E	702	NAG	O7-C7-N2-C2
5	E	702	NAG	C1-C2-N2-C7
5	E	702	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	702	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	143/177 (80%)	0.68	16 (11%) 10 10	27, 44, 78, 98	0
1	F	158/177 (89%)	0.41	8 (5%) 33 36	26, 42, 72, 99	0
2	A	222/233 (95%)	0.63	16 (7%) 21 23	30, 45, 75, 104	0
2	H	222/233 (95%)	0.37	14 (6%) 26 27	25, 40, 70, 82	0
3	B	212/214 (99%)	0.81	16 (7%) 20 21	38, 54, 75, 97	0
3	L	211/214 (98%)	0.34	2 (0%) 81 83	27, 42, 61, 76	0
All	All	1168/1248 (93%)	0.54	72 (6%) 26 28	25, 44, 73, 104	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	100(C)	GLY	5.0
2	A	43	GLN	5.0
1	E	549	TRP	4.9
2	H	100(C)	GLY	4.7
2	A	41	PRO	4.6
2	A	3	LEU	4.4
2	A	44	GLY	4.1
1	F	582	THR	4.0
2	A	100	TYR	3.9
2	H	92	CYS	3.9
1	E	632	TYR	3.7
2	A	42	GLY	3.7
2	H	41	PRO	3.6
1	E	550	PHE	3.5
1	F	450	SER	3.4
3	B	192	TYR	3.2
1	E	633	VAL	3.2
3	B	214	CYS	3.2
1	E	600	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	190	LYS	3.1
1	F	445	HIS	3.1
3	B	154	LEU	3.0
2	H	3	LEU	3.0
1	E	458	PRO	3.0
1	E	421	HIS	2.9
2	H	42	GLY	2.9
1	F	433	LEU	2.9
1	E	451	GLY	2.9
2	A	216	CYS	2.8
2	H	27	GLY	2.8
3	B	87	PHE	2.8
1	F	598	GLY	2.8
2	A	134	GLY	2.8
1	E	452	CYS	2.7
2	H	127	SER	2.7
1	E	603	ILE	2.6
1	E	510	THR	2.6
3	B	193	ALA	2.6
3	B	114	SER	2.6
3	B	129	THR	2.6
1	E	434	ASN	2.6
1	F	444	GLN	2.5
2	A	127	SER	2.5
3	B	150	VAL	2.5
2	H	40	ALA	2.5
1	E	645	ASN	2.4
1	E	605	PRO	2.4
2	H	100	TYR	2.4
3	B	209	PHE	2.4
1	F	434	ASN	2.4
3	B	212	GLY	2.3
1	F	452	CYS	2.3
2	A	92	CYS	2.3
1	E	635	GLY	2.2
2	H	43	GLN	2.2
2	H	214	LYS	2.2
3	B	81	GLU	2.2
3	B	186	TYR	2.2
1	E	628	LYS	2.2
2	H	28	THR	2.2
2	A	28	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	84	SER	2.2
2	H	215	SER	2.2
2	A	100(G)	TYR	2.2
2	A	25	SER	2.2
3	B	189	HIS	2.1
3	B	191	VAL	2.1
2	A	215	SER	2.1
3	L	114	SER	2.1
3	B	3	GLU	2.1
2	A	8	GLY	2.1
3	L	125	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	1	14/15	-	-	45,54,60,71	0
4	NAG	C	2	14/15	-	-	64,80,91,96	0
4	NAG	D	1	14/15	-	-	45,57,69,71	0
4	NAG	D	2	14/15	-	-	62,74,96,98	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	E	701	14/15	0.51	0.19	82,93,101,117	0
5	NAG	E	702	14/15	0.58	0.17	67,88,97,100	0
5	NAG	E	703	14/15	0.62	0.18	91,100,105,106	0
5	NAG	F	701	14/15	0.68	0.15	82,90,101,108	0

6.5 Other polymers [i](#)

There are no such residues in this entry.