



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:56 AM UTC

PDB ID : 6V95 / pdb\_00006v95  
Title : Peanut lectin complexed with divalent N-beta-D-galactopyranosyl-L-tartaram idoyl derivative (diNGT)  
Authors : Otero, L.H.; Primo, E.D.; Cagnoni, A.J.; Klinke, S.; Goldbaum, F.A.; Uhrig, M.L.  
Deposited on : 2019-12-13  
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	 6% 91% 7%
1	B	236	 6% 88% 10%
1	C	236	 45% 88% 8%
1	D	236	 19% 89% 8%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8369 atoms, of which 192 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galactose-binding lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	Total 1743	C 1102	N 287	O 352	S 2	0	0	0
1	B	232	Total 1743	C 1102	N 287	O 352	S 2	0	0	0
1	C	232	Total 1743	C 1102	N 287	O 352	S 2	0	0	0
1	D	232	Total 1743	C 1102	N 287	O 352	S 2	0	0	0

- Molecule 2 is (2R,3R)-N-[(1-[(3S,3aR,6S,6aR)-6-[4-[(2R,3R)-2,3-dihydroxy-4-oxo-4-[(2R,3R,4R,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl]amino}butanoyl]amino}methyl)-1H-1,2,3-triazol-1-yl]hexahydrofuro[3,2-b]furan-3-yl}-1H-1,2,3-triazol-4-yl)methyl]-2,3-dihydroxy-N'-[(2R,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl]butanediamide (non-preferred name) (CCD ID: QSG) (formula: C<sub>32</sub>H<sub>48</sub>N<sub>10</sub>O<sub>20</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	74	0
			110	32	48	10	20		
2	B	1	Total	C	H	N	O	76	0
			110	32	48	10	20		
2	C	1	Total	C	H	N	O	76	0
			110	32	48	10	20		
2	D	1	Total	C	H	N	O	80	0
			110	32	48	10	20		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	B	1	Total	Mn	0	0
			1	1		
3	C	1	Total	Mn	0	0
			1	1		
3	D	1	Total	Mn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	241	Total	O	0	0
			241	241		
5	B	266	Total	O	0	0
			266	266		
5	C	211	Total	O	0	0
			211	211		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	D	231	Total 231	O 231	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Galactose-binding lectin




- Molecule 1: Galactose-binding lectin




- Molecule 1: Galactose-binding lectin



- Molecule 1: Galactose-binding lectin





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.94Å 124.55Å 127.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.06 – 1.78 26.06 – 1.78	Depositor EDS
% Data completeness (in resolution range)	100.0 (26.06-1.78) 99.9 (26.06-1.78)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 1.78Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.222 , 0.243 0.225 , 0.247	Depositor DCC
$R_{free}$ test set	5835 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtrriage
Anisotropy	0.646	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 39.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.007 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8369	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MN, QSG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.83	0/1779	1.13	5/2426 (0.2%)
1	B	0.89	0/1779	1.15	8/2426 (0.3%)
1	C	0.81	0/1779	1.15	5/2426 (0.2%)
1	D	0.85	0/1779	1.15	4/2426 (0.2%)
All	All	0.85	0/7116	1.14	22/9704 (0.2%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	181	ASP	CA-CB-CG	7.50	120.10	112.60
1	C	38	ASN	CA-CB-CG	7.17	119.78	112.60
1	C	31	ASN	CA-CB-CG	6.83	119.43	112.60
1	D	31	ASN	CA-CB-CG	6.79	119.39	112.60
1	A	31	ASN	CA-CB-CG	6.65	119.25	112.60
1	B	31	ASN	CA-CB-CG	6.37	118.97	112.60
1	D	182	ASN	CA-CB-CG	6.08	118.69	112.60
1	B	179	THR	CA-C-N	5.87	131.64	120.97
1	B	179	THR	C-N-CA	5.87	131.64	120.97
1	B	19	ASN	CA-CB-CG	5.74	118.34	112.60
1	D	123	ASP	CA-CB-CG	5.67	118.27	112.60
1	B	182	ASN	CA-CB-CG	5.64	118.24	112.60
1	C	182	ASN	CA-CB-CG	5.64	118.24	112.60
1	A	182	ASN	CA-CB-CG	5.59	118.19	112.60
1	A	19	ASN	CA-CB-CG	5.55	118.15	112.60
1	D	80	ASP	CA-CB-CG	5.50	118.10	112.60
1	C	29	ASN	CA-CB-CG	5.43	118.03	112.60
1	C	184	ASP	CA-CB-CG	5.42	118.02	112.60
1	B	184	ASP	CA-CB-CG	5.41	118.00	112.60
1	B	110	ASP	CA-CB-CG	5.25	117.85	112.60

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	PHE	CA-C-N	5.03	129.97	122.99
1	A	88	PHE	C-N-CA	5.03	129.97	122.99

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1743	0	1699	8	0
1	B	1743	0	1699	11	0
1	C	1743	0	1699	15	0
1	D	1743	0	1699	10	0
2	A	62	48	0	0	0
2	B	62	48	0	0	0
2	C	62	48	0	0	0
2	D	62	48	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	241	0	0	1	0
5	B	266	0	0	2	0
5	C	211	0	0	1	0
5	D	231	0	0	0	0
All	All	8177	192	6796	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:77:LYS:HE3	1:C:77:LYS:H	1.32	0.93
1:D:32:ILE:HD13	1:D:46:VAL:HG11	1.77	0.67
1:A:32:ILE:HD13	1:A:46:VAL:HG11	1.77	0.66
1:B:32:ILE:HD13	1:B:46:VAL:HG11	1.80	0.64
1:C:77:LYS:H	1:C:77:LYS:CE	2.12	0.59
1:C:29:ASN:HB2	1:D:219:LEU:HD11	1.87	0.56
1:C:32:ILE:HD13	1:C:46:VAL:HG11	1.89	0.55
1:A:34:LEU:O	1:A:44:GLY:HA3	2.08	0.54
1:D:37:LEU:HG	1:D:76:ILE:HD13	1.91	0.53
1:A:50:MET:SD	1:B:17:ALA:HA	2.48	0.52
1:A:41:ASN:HD22	1:A:211:SER:HA	1.74	0.52
1:D:41:ASN:HD22	1:D:211:SER:HA	1.74	0.52
1:B:173:THR:HG21	1:C:173:THR:HG21	1.92	0.51
1:C:77:LYS:HE3	1:C:77:LYS:N	2.14	0.51
1:D:34:LEU:O	1:D:44:GLY:HA3	2.10	0.51
1:B:41:ASN:HD22	1:B:211:SER:HA	1.76	0.51
1:D:19:ASN:ND2	1:D:49:ALA:HA	2.28	0.49
1:C:10:SER:HA	5:C:425:HOH:O	2.13	0.48
1:D:55:TRP:HB2	1:D:195:LYS:HZ2	1.79	0.48
1:A:173:THR:HG21	1:D:173:THR:HG21	1.96	0.47
1:C:41:ASN:HD22	1:C:211:SER:HA	1.81	0.46
1:A:50:MET:HE1	5:B:451:HOH:O	2.15	0.46
1:C:74:LYS:CE	1:D:10:SER:HB3	2.46	0.46
1:C:24:VAL:HG22	1:C:34:LEU:HA	1.98	0.46
1:B:128:SER:HA	5:B:471:HOH:O	2.18	0.43
1:B:34:LEU:O	1:B:44:GLY:HA3	2.19	0.43
1:B:164:THR:HG21	1:C:164:THR:HG21	2.00	0.42
5:A:401:HOH:O	1:B:5:SER:HB3	2.19	0.42
1:A:50:MET:HB2	1:B:50:MET:HE3	2.01	0.42
1:A:133:PRO:HD2	1:A:137:HIS:CE1	2.54	0.42
1:B:133:PRO:HD2	1:B:137:HIS:CE1	2.54	0.42
1:C:10:SER:HB3	1:D:74:LYS:CE	2.50	0.42
1:C:12:SER:HA	1:C:26:VAL:HG11	2.00	0.42
1:C:133:PRO:HD2	1:C:137:HIS:CE1	2.55	0.42
1:B:177:ALA:HB1	1:B:185:ILE:HD11	2.03	0.41
1:C:34:LEU:O	1:C:44:GLY:HA3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/236 (98%)	225 (98%)	5 (2%)	0	100	100
1	B	230/236 (98%)	226 (98%)	4 (2%)	0	100	100
1	C	230/236 (98%)	226 (98%)	4 (2%)	0	100	100
1	D	230/236 (98%)	227 (99%)	3 (1%)	0	100	100
All	All	920/944 (98%)	904 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/201 (98%)	193 (98%)	4 (2%)	48	29
1	B	197/201 (98%)	193 (98%)	4 (2%)	48	29
1	C	197/201 (98%)	191 (97%)	6 (3%)	36	15
1	D	197/201 (98%)	193 (98%)	4 (2%)	48	29
All	All	788/804 (98%)	770 (98%)	18 (2%)	44	25

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	LEU
1	A	80	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	182	ASN
1	A	185	ILE
1	B	37	LEU
1	B	80	ASP
1	B	181	ASP
1	B	182	ASN
1	C	37	LEU
1	C	38	ASN
1	C	77	LYS
1	C	80	ASP
1	C	83	ASP
1	C	182	ASN
1	D	37	LEU
1	D	80	ASP
1	D	112	LYS
1	D	185	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	15	ASN
1	A	41	ASN
1	A	190	GLN
1	B	7	ASN
1	B	15	ASN
1	B	41	ASN
1	B	61	ASN
1	B	190	GLN
1	C	7	ASN
1	C	33	GLN
1	C	41	ASN
1	C	61	ASN
1	C	190	GLN
1	D	7	ASN
1	D	15	ASN
1	D	19	ASN
1	D	41	ASN
1	D	61	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	QSG	C	301	-	67,67,67	0.17	0	98,98,98	0.69	3 (3%)
2	QSG	B	301	-	67,67,67	0.27	0	98,98,98	1.03	7 (7%)
2	QSG	A	301	-	67,67,67	0.17	0	98,98,98	0.70	1 (1%)
2	QSG	D	301	-	67,67,67	0.14	0	98,98,98	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QSG	C	301	-	-	22/54/116/116	0/6/6/6
2	QSG	B	301	-	-	17/54/116/116	0/6/6/6
2	QSG	A	301	-	-	18/54/116/116	0/6/6/6
2	QSG	D	301	-	-	20/54/116/116	0/6/6/6

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	QSG	C21-C22-N9	4.33	122.20	112.34
2	B	301	QSG	C19-C14-N3	2.79	118.19	112.95
2	B	301	QSG	C2-C3-C4	2.56	115.33	110.83
2	C	301	QSG	C2-C3-C4	2.51	115.24	110.83
2	A	301	QSG	C2-C3-C4	2.47	115.16	110.83
2	B	301	QSG	C8-C7-N1	2.47	119.57	116.22
2	C	301	QSG	C15-C14-C19	2.39	105.45	102.03
2	B	301	QSG	C15-C14-N3	-2.20	110.09	113.08
2	B	301	QSG	C12-C11-N2	2.09	117.10	112.34
2	C	301	QSG	O3-C3-C4	2.09	115.30	110.38
2	B	301	QSG	O15-C15-C14	2.09	108.11	104.86

There are no chirality outliers.

All (77) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	QSG	C16-C17-N6-N7
2	A	301	QSG	C16-C17-N6-C20
2	A	301	QSG	C28-C27-N10-C26
2	A	301	QSG	O31-C27-N10-C26
2	B	301	QSG	C16-C17-N6-N7
2	B	301	QSG	C16-C17-N6-C20
2	B	301	QSG	O31-C27-N10-C26
2	B	301	QSG	C21-C22-N9-C23
2	B	301	QSG	N8-C21-C22-N9
2	B	301	QSG	C7-C8-C9-C10
2	B	301	QSG	C7-C8-C9-O9
2	B	301	QSG	O8-C8-C9-C10
2	C	301	QSG	C16-C17-N6-N7
2	C	301	QSG	C16-C17-N6-C20
2	C	301	QSG	C28-C27-N10-C26
2	C	301	QSG	O31-C27-N10-C26
2	C	301	QSG	C25-C26-N10-C27
2	C	301	QSG	O26-C26-N10-C27
2	C	301	QSG	C15-C14-N3-N4
2	C	301	QSG	C20-C21-C22-N9
2	D	301	QSG	C28-C27-N10-C26
2	D	301	QSG	O31-C27-N10-C26
2	D	301	QSG	C25-C26-N10-C27
2	D	301	QSG	O26-C26-N10-C27

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	D	301	QSG	C24-C23-N9-C22
2	D	301	QSG	C8-C7-N1-C1
2	D	301	QSG	O7-C7-N1-C1
2	D	301	QSG	O23-C23-C24-C25
2	D	301	QSG	C7-C8-C9-C10
2	D	301	QSG	C7-C8-C9-O9
2	D	301	QSG	O8-C8-C9-C10
2	D	301	QSG	O8-C8-C9-O9
2	D	301	QSG	O23-C23-N9-C22
2	B	301	QSG	O8-C8-C9-O9
2	D	301	QSG	N9-C23-C24-C25
2	A	301	QSG	O31-C31-C32-O32
2	D	301	QSG	O31-C31-C32-O32
2	B	301	QSG	O31-C31-C32-O32
2	C	301	QSG	O31-C31-C32-O32
2	A	301	QSG	C20-C21-C22-N9
2	C	301	QSG	N2-C11-C12-C13
2	A	301	QSG	N8-C21-C22-N9
2	C	301	QSG	N2-C11-C12-N5
2	A	301	QSG	C24-C23-N9-C22
2	C	301	QSG	O8-C8-C9-O9
2	A	301	QSG	C7-C8-C9-O9
2	A	301	QSG	O8-C8-C9-C10
2	C	301	QSG	C7-C8-C9-O9
2	C	301	QSG	O8-C8-C9-C10
2	C	301	QSG	O23-C23-C24-C25
2	A	301	QSG	O8-C8-C9-O9
2	C	301	QSG	N9-C23-C24-C25
2	A	301	QSG	N2-C11-C12-C13
2	B	301	QSG	C20-C21-C22-N9
2	B	301	QSG	N2-C11-C12-C13
2	D	301	QSG	C15-C14-N3-N4
2	A	301	QSG	C7-C8-C9-C10
2	C	301	QSG	C7-C8-C9-C10
2	D	301	QSG	C19-C14-N3-N4
2	A	301	QSG	N2-C11-C12-N5
2	B	301	QSG	N2-C11-C12-N5
2	C	301	QSG	N8-C21-C22-N9
2	C	301	QSG	O24-C24-C25-O25
2	D	301	QSG	C12-C11-N2-C10
2	A	301	QSG	O23-C23-N9-C22
2	A	301	QSG	O23-C23-C24-O24

*Continued on next page...*


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	301	QSG	O23-C23-C24-O24
2	C	301	QSG	O23-C23-C24-O24
2	B	301	QSG	O23-C23-C24-C25
2	A	301	QSG	N9-C23-C24-O24
2	B	301	QSG	N9-C23-C24-O24
2	C	301	QSG	N9-C23-C24-O24
2	D	301	QSG	N9-C23-C24-O24
2	C	301	QSG	C12-C11-N2-C10
2	A	301	QSG	N9-C23-C24-C25
2	B	301	QSG	N9-C23-C24-C25
2	D	301	QSG	O23-C23-C24-O24

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/236 (98%)	0.46	14 (6%) 27 33	18, 26, 44, 96	0
1	B	232/236 (98%)	0.37	15 (6%) 25 30	18, 24, 41, 98	0
1	C	232/236 (98%)	2.00	106 (45%) 0 0	18, 38, 67, 89	0
1	D	232/236 (98%)	0.87	46 (19%) 3 3	18, 28, 54, 82	0
All	All	928/944 (98%)	0.93	181 (19%) 3 3	18, 27, 59, 98	0

All (181) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	232	THR	8.0
1	B	232	THR	6.9
1	A	232	THR	6.4
1	C	181	ASP	6.2
1	C	213	GLY	6.0
1	C	101	ILE	5.8
1	C	37	LEU	5.8
1	C	82	ALA	5.7
1	C	212	LEU	5.5
1	B	179	THR	5.5
1	C	38	ASN	5.4
1	C	79	TYR	5.4
1	C	29	ASN	5.4
1	C	36	ASN	5.4
1	C	108	VAL	5.4
1	C	114	ALA	5.4
1	C	129	GLU	5.3
1	D	232	THR	5.2
1	C	40	VAL	5.2
1	A	181	ASP	5.2
1	C	145	VAL	5.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	103	GLY	5.1
1	C	102	GLY	4.9
1	D	58	ALA	4.9
1	B	181	ASP	4.9
1	C	76	ILE	4.8
1	C	214	GLY	4.8
1	C	125	TYR	4.7
1	C	39	LYS	4.7
1	C	77	LYS	4.6
1	C	111	THR	4.6
1	C	80	ASP	4.6
1	C	98	ALA	4.6
1	C	25	THR	4.5
1	D	26	VAL	4.5
1	C	100	SER	4.4
1	C	78	ASP	4.4
1	C	81	PRO	4.4
1	C	24	VAL	4.4
1	C	26	VAL	4.3
1	C	110	ASP	4.3
1	C	28	SER	4.3
1	C	130	TYR	4.3
1	C	128	SER	4.3
1	C	113	GLY	4.3
1	C	96	ILE	4.2
1	C	27	LEU	4.2
1	C	11	PHE	4.2
1	C	115	GLY	4.2
1	C	97	PRO	4.1
1	C	43	VAL	4.0
1	D	59	THR	3.9
1	C	106	LEU	3.9
1	C	107	GLY	3.9
1	D	181	ASP	3.9
1	A	183	GLY	3.9
1	C	35	THR	3.8
1	C	182	ASN	3.8
1	D	38	ASN	3.7
1	C	211	SER	3.7
1	D	182	ASN	3.7
1	A	182	ASN	3.7
1	C	180	ASN	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	99	GLY	3.6
1	C	179	THR	3.5
1	C	42	SER	3.5
1	C	12	SER	3.5
1	C	210	GLY	3.5
1	C	208	ALA	3.4
1	D	8	PHE	3.4
1	C	109	SER	3.4
1	C	41	ASN	3.4
1	D	76	ILE	3.4
1	C	14	GLY	3.4
1	C	95	GLN	3.4
1	D	217	ILE	3.4
1	C	13	GLU	3.3
1	D	11	PHE	3.3
1	C	30	GLY	3.2
1	C	104	GLY	3.2
1	B	180	ASN	3.1
1	C	127	ASN	3.1
1	C	34	LEU	3.1
1	C	22	GLY	3.1
1	A	50	MET	3.1
1	C	209	SER	3.0
1	C	8	PHE	3.0
1	B	182	ASN	3.0
1	D	29	ASN	3.0
1	C	215	ARG	3.0
1	D	213	GLY	3.0
1	C	148	VAL	3.0
1	C	10	SER	2.9
1	D	10	SER	2.9
1	D	28	SER	2.9
1	C	124	THR	2.9
1	D	78	ASP	2.9
1	A	13	GLU	2.9
1	C	112	LYS	2.9
1	C	151	VAL	2.8
1	D	62	VAL	2.8
1	A	180	ASN	2.8
1	D	13	GLU	2.8
1	A	231	THR	2.8
1	C	126	SER	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	83	ASP	2.8
1	C	15	ASN	2.7
1	A	20	PHE	2.7
1	C	142	VAL	2.7
1	D	40	VAL	2.7
1	D	111	THR	2.7
1	D	50	MET	2.7
1	C	217	ILE	2.7
1	C	105	THR	2.6
1	C	131	ASN	2.6
1	C	50	MET	2.6
1	C	32	ILE	2.6
1	C	17	ALA	2.6
1	D	14	GLY	2.6
1	C	20	PHE	2.6
1	D	37	LEU	2.6
1	D	9	ASN	2.6
1	C	23	ASP	2.5
1	D	60	GLY	2.5
1	C	74	LYS	2.5
1	D	24	VAL	2.5
1	B	58	ALA	2.5
1	D	12	SER	2.5
1	D	7	ASN	2.5
1	D	17	ALA	2.5
1	C	7	ASN	2.5
1	D	212	LEU	2.5
1	D	53	ARG	2.5
1	D	80	ASP	2.4
1	C	16	PRO	2.4
1	D	201	ARG	2.4
1	D	112	LYS	2.4
1	B	231	THR	2.4
1	B	1	ALA	2.4
1	B	20	PHE	2.4
1	C	75	ASP	2.4
1	B	185	ILE	2.4
1	B	230	ILE	2.4
1	C	18	ILE	2.4
1	B	50	MET	2.4
1	C	216	GLN	2.3
1	B	13	GLU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	185	ILE	2.3
1	C	146	ASP	2.3
1	A	230	ILE	2.3
1	D	39	LYS	2.3
1	D	18	ILE	2.3
1	D	1	ALA	2.2
1	D	183	GLY	2.2
1	A	53	ARG	2.2
1	C	231	THR	2.2
1	C	93	ASP	2.2
1	C	185	ILE	2.2
1	C	21	GLN	2.2
1	D	16	PRO	2.2
1	C	44	GLY	2.1
1	C	84	GLY	2.1
1	D	27	LEU	2.1
1	D	79	TYR	2.1
1	D	77	LYS	2.1
1	C	46	VAL	2.1
1	A	9	ASN	2.1
1	C	31	ASN	2.1
1	B	57	SER	2.1
1	C	133	PRO	2.1
1	C	86	ILE	2.1
1	B	38	ASN	2.0
1	C	9	ASN	2.0
1	A	221	ARG	2.0
1	C	156	VAL	2.0
1	C	154	ASN	2.0
1	D	30	GLY	2.0
1	D	36	ASN	2.0
1	D	61	ASN	2.0
1	D	55	TRP	2.0
1	C	207	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	QSG	C	301	62/62	0.66	0.28	96,132,134,134	76
2	QSG	D	301	62/62	0.76	0.18	32,116,125,125	80
4	CA	C	303	1/1	0.86	0.16	44,44,44,44	0
2	QSG	A	301	62/62	0.89	0.12	26,108,117,117	74
2	QSG	B	301	62/62	0.90	0.11	23,114,134,134	76
3	MN	A	302	1/1	0.96	0.13	33,33,33,33	0
3	MN	C	302	1/1	0.97	0.08	53,53,53,53	0
4	CA	A	303	1/1	0.98	0.03	25,25,25,25	0
4	CA	D	303	1/1	0.98	0.03	26,26,26,26	0
4	CA	B	303	1/1	0.99	0.02	22,22,22,22	0
3	MN	D	302	1/1	0.99	0.12	35,35,35,35	0
3	MN	B	302	1/1	0.99	0.10	32,32,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around QSG C 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around QSG D 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around QSG A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around QSG B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.