



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:18 PM UTC

PDB ID : 6VHU / pdb_00006vhu
Title : Klebsiella oxytoca NpsA N-terminal subdomain in space group P21
Authors : Kreitler, D.F.; Gulick, A.M.
Deposited on : 2020-01-10
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	
1	B	406	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6421 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NpsA Adenylation Domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	2960	1890	495	556	19	0	7	0
1	B	381	2890	1852	480	538	20	0	6	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A2U4DY99
A	0	HIS	-	expression tag	UNP A0A2U4DY99
A	258	SER	LEU	variant	UNP A0A2U4DY99
A	312	ALA	GLU	engineered mutation	UNP A0A2U4DY99
A	313	ALA	GLU	engineered mutation	UNP A0A2U4DY99
A	314	ALA	GLN	engineered mutation	UNP A0A2U4DY99
B	-1	GLY	-	expression tag	UNP A0A2U4DY99
B	0	HIS	-	expression tag	UNP A0A2U4DY99
B	258	SER	LEU	variant	UNP A0A2U4DY99
B	312	ALA	GLU	engineered mutation	UNP A0A2U4DY99
B	313	ALA	GLU	engineered mutation	UNP A0A2U4DY99
B	314	ALA	GLN	engineered mutation	UNP A0A2U4DY99

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cl	0	0
			4	4		
2	B	3	Total	Cl	0	0
			3	3		


- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0

- Molecule 4 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Br 1 1	0	0

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	287	Total O 287 287	0	0
6	B	259	Total O 260 260	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NpsA Adenylation Domain



- Molecule 1: NpsA Adenylation Domain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.95Å 59.11Å 80.41Å 90.00° 101.12° 90.00°	Depositor
Resolution (Å)	70.60 – 1.60 70.60 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.5 (70.60-1.60) 89.7 (70.60-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.86 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.15_3459	Depositor
R, R_{free}	0.188 , 0.218 0.188 , 0.219	Depositor DCC
R_{free} test set	4200 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtrriage
Anisotropy	0.280	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6421	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, BR, EPE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	1/3017 (0.0%)	0.55	0/4111
1	B	0.56	0/2944	0.54	0/4014
All	All	0.59	1/5961 (0.0%)	0.54	0/8125

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	196	GLN	C-O	-5.16	1.17	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2960	0	2965	14	0
1	B	2890	0	2890	28	0
2	A	4	0	0	0	0
2	B	3	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	B	15	0	18	1	0
6	A	287	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	260	0	0	2	0
All	All	6421	0	5873	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:MET:HE3	1:B:88:MET:HE3	1.38	1.04
1:A:199:ASP:OD1	1:A:201:SER:OG	2.02	0.78
1:B:56[B]:ASN:OD1	1:B:57:CYS:N	2.21	0.74
1:A:46[B]:CYS:SG	6:A:825:HOH:O	2.46	0.72
1:B:131:ARG:NH2	1:B:135:GLN:OE1	2.29	0.65
1:A:200:ILE:HD11	6:A:617:HOH:O	1.98	0.62
1:B:274:PHE:HE2	1:B:277:PRO:HG3	1.66	0.61
1:B:82:MET:HE1	1:B:154:ALA:O	2.05	0.57
1:B:274:PHE:HZ	1:B:315:LEU:HD13	1.72	0.55
1:B:338:ARG:NH1	1:B:340:GLN:HG3	2.23	0.53
1:A:290:LYS:HE2	1:A:292:PHE:CZ	2.45	0.52
1:A:60:LYS:HG3	1:A:85:GLY:HA3	1.92	0.50
1:B:97:GLU:O	1:B:121:PRO:HB3	2.13	0.48
1:B:274:PHE:CZ	1:B:315:LEU:HD13	2.47	0.48
1:B:297:CYS:HB2	5:B:501:EPE:H81	1.95	0.48
1:A:320:PRO:HB2	1:A:399:VAL:HB	1.95	0.48
1:B:67:LYS:NZ	1:B:118:THR:OG1	2.38	0.47
1:B:315:LEU:HD23	1:B:315:LEU:HA	1.65	0.45
1:B:38:THR:OG1	1:B:41:GLU:HG3	2.16	0.45
1:B:97:GLU:OE2	6:B:601:HOH:O	2.21	0.45
1:A:11:LYS:NZ	1:A:19:GLU:OE2	2.32	0.44
1:B:10:LEU:HD23	1:B:10:LEU:HA	1.78	0.44
1:A:274:PHE:HE2	1:A:277:PRO:HG3	1.83	0.44
1:B:16:ILE:HD12	1:B:177:ALA:HB1	2.00	0.44
1:A:74:LEU:HD21	1:A:200:ILE:HG21	2.00	0.44
1:A:200:ILE:CD1	6:A:617:HOH:O	2.63	0.44
1:B:104:LEU:HD12	1:B:125:TRP:CE2	2.52	0.43
1:B:154:ALA:HB2	1:B:173:HIS:CD2	2.53	0.43
1:B:196:GLN:HB2	1:B:219:LEU:HD11	2.00	0.43
1:B:57:CYS:O	1:B:59[A]:ILE:HD12	2.18	0.43
1:B:159:THR:HG22	1:B:167:LYS:HB3	2.01	0.43
1:B:83:LYS:HE3	1:B:145:TYR:CD2	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:394:GLN:H	1:B:394:GLN:CD	2.24	0.42
1:B:57:CYS:SG	1:B:111:TYR:CE1	3.11	0.41
1:B:59[B]:ILE:HG22	1:B:62:ASP:CG	2.45	0.41
1:A:83:LYS:HE3	1:A:145:TYR:CD2	2.55	0.41
1:B:207:ASN:O	1:B:211:MET:HB2	2.20	0.41
1:A:92:ASN:OD1	1:A:94:ASN:N	2.49	0.40
1:A:253:LEU:O	1:A:257:VAL:HG22	2.21	0.40
1:A:230:MET:HG2	1:B:226:LEU:HB2	2.04	0.40
1:B:338:ARG:NH1	6:B:608:HOH:O	2.27	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/406 (96%)	378 (97%)	10 (3%)	2 (0%)	24	10
1	B	379/406 (93%)	367 (97%)	12 (3%)	0	100	100
All	All	769/812 (95%)	745 (97%)	22 (3%)	2 (0%)	36	20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	ASP
1	A	143	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/336 (96%)	319 (99%)	2 (1%)	78	66
1	B	310/336 (92%)	309 (100%)	1 (0%)	86	78
All	All	631/672 (94%)	628 (100%)	3 (0%)	81	70

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	142	SER
1	A	196	GLN
1	B	155	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	300	ASN
1	B	300	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 9 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EPE	B	501	-	15,15,15	1.02	0	19,20,20	1.94	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	B	501	-	-	1/9/19/19	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	501	EPE	O3S-S-O1S	-3.83	101.82	111.40
5	B	501	EPE	O1S-S-C10	3.66	112.26	106.73
5	B	501	EPE	O3S-S-C10	2.96	111.79	106.00
5	B	501	EPE	C5-N4-C3	2.63	114.50	108.84
5	B	501	EPE	C7-N4-C5	-2.55	104.43	111.24
5	B	501	EPE	C6-N1-C2	-2.46	103.55	108.84

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	501	EPE	N4-C7-C8-O8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	501	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/406 (95%)	0.30	22 (5%) 29 30	8, 23, 45, 74	7 (1%)
1	B	381/406 (93%)	0.72	45 (11%) 9 9	15, 26, 52, 74	6 (1%)
All	All	768/812 (94%)	0.51	67 (8%) 16 16	8, 24, 50, 74	13 (1%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	TYR	6.5
1	B	117	VAL	5.3
1	A	140	GLN	5.0
1	B	164	GLY	4.5
1	B	120	VAL	4.4
1	B	57	CYS	4.3
1	B	321	VAL	4.0
1	B	143	GLY	3.8
1	A	312	ALA	3.8
1	B	138	THR	3.8
1	B	124	ALA	3.6
1	B	116	ASP	3.5
1	B	56[A]	ASN	3.4
1	B	286	ALA	3.3
1	B	59[A]	ILE	3.2
1	B	119	SER	3.2
1	B	118	THR	3.2
1	B	159	THR	3.2
1	B	122	GLY	3.1
1	B	46	CYS	3.1
1	A	141	ARG	3.1
1	A	124	ALA	3.1
1	A	143	GLY	3.0
1	B	347	TYR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	285	GLN	3.0
1	A	55	ASN	2.9
1	B	10	LEU	2.8
1	B	315	LEU	2.8
1	B	125	TRP	2.7
1	B	282	VAL	2.7
1	B	344[A]	CYS	2.7
1	B	316	SER	2.7
1	A	162	SER	2.7
1	B	343	THR	2.7
1	A	139	GLN	2.6
1	A	362	SER	2.6
1	A	401	VAL	2.6
1	B	401	VAL	2.6
1	B	379	THR	2.6
1	A	142	SER	2.6
1	B	55	ASN	2.6
1	B	121	PRO	2.5
1	B	320	PRO	2.5
1	B	137	CYS	2.5
1	B	145	TYR	2.4
1	A	97	GLU	2.4
1	A	57	CYS	2.3
1	A	94	ASN	2.3
1	B	94	ASN	2.3
1	B	322	PRO	2.3
1	B	58	LEU	2.3
1	A	321	VAL	2.3
1	A	96	PRO	2.2
1	B	96	PRO	2.2
1	A	111	TYR	2.2
1	A	138	THR	2.2
1	B	144	ASN	2.2
1	B	274	PHE	2.2
1	A	93	SER	2.2
1	B	134	ILE	2.1
1	A	161	GLY	2.1
1	B	287	GLY	2.1
1	B	52	VAL	2.1
1	B	394	GLN	2.1
1	B	230[A]	MET	2.0
1	A	311	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	123	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EPE	B	501	15/15	0.76	0.17	60,62,64,66	0
2	CL	A	503	1/1	0.95	0.09	28,28,28,28	0
3	K	A	505	1/1	0.96	0.08	37,37,37,37	0
2	CL	A	501	1/1	0.97	0.09	19,19,19,19	0
2	CL	B	503	1/1	0.98	0.07	24,24,24,24	0
2	CL	B	504	1/1	0.99	0.04	22,22,22,22	0
2	CL	B	502	1/1	0.99	0.03	20,20,20,20	0
2	CL	A	504	1/1	0.99	0.04	21,21,21,21	0
4	BR	A	506	1/1	1.00	0.02	24,24,24,24	0
2	CL	A	502	1/1	1.00	0.06	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.