



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:44 PM UTC

PDB ID : 6VUA / pdb_00006vua
Title : X-ray structure of human CD38 catalytic domain with 2'-Cl-araNAD+
Authors : Dai, Z.; Zhang, X.N.; Nasertorabi, F.; Han, G.W.; Stevens, R.C.; Zhang, Y.
Deposited on : 2020-02-14
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	 7% 84% 14% .
1	B	257	 11% 85% 12% ..

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 4973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	255	2216	1405	373	419	19	0	26	0
1	B	251	2094	1332	356	388	18	0	16	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	SER	-	expression tag	UNP P28907
A	100	ASP	ASN	engineered mutation	UNP P28907
A	164	ALA	ASN	engineered mutation	UNP P28907
A	209	ASP	ASN	engineered mutation	UNP P28907
A	219	ASP	ASN	engineered mutation	UNP P28907
B	44	SER	-	expression tag	UNP P28907
B	100	ASP	ASN	engineered mutation	UNP P28907
B	164	ALA	ASN	engineered mutation	UNP P28907
B	209	ASP	ASN	engineered mutation	UNP P28907
B	219	ASP	ASN	engineered mutation	UNP P28907

- Molecule 2 is [(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl [(2 {R},3 {R},4 {R})-4-chloranyl-3-oxidanyl-oxolan-2-yl]methyl hydrogen phosphate (CCD ID: ROJ) (formula: C₁₅H₂₂ClN₅O₁₂P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	C O	0	0
			10	6 4		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	282	Total	O	0	4
			286	286		
8	B	220	Total	O	0	6
			226	226		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1



- Molecule 1: ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	114.76Å 114.76Å 97.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.71 – 1.50 28.71 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.71-1.50) 100.0 (28.71-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.166 , 0.190 0.172 , 0.194	Depositor DCC
R_{free} test set	5330 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtrriage
Anisotropy	0.427	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4973	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ROJ, PGE, K, PO4, PEG, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.06	2/2311 (0.1%)	1.12	4/3128 (0.1%)
1	B	1.07	2/2174 (0.1%)	1.23	6/2950 (0.2%)
All	All	1.07	4/4485 (0.1%)	1.18	10/6078 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	HIS	CE1-NE2	8.27	1.40	1.32
1	B	133	HIS	CE1-NE2	6.97	1.39	1.32
1	A	224	SER	C-O	5.93	1.31	1.24
1	B	81	ASP	CG-OD1	-5.13	1.15	1.25

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	100	ASP	CB-CA-C	-6.15	100.76	110.85
1	B	140	ARG	CA-C-N	6.14	129.01	120.29
1	B	140	ARG	C-N-CA	6.14	129.01	120.29
1	B	72	GLU	CB-CA-C	6.13	122.89	110.38
1	B	97	HIS	CA-CB-CG	-5.68	108.12	113.80
1	A	81	ASP	CA-CB-CG	-5.67	106.93	112.60
1	B	229	ASN	CA-CB-CG	-5.14	107.46	112.60
1	A	72	GLU	CA-C-N	5.08	128.62	120.30
1	A	72	GLU	C-N-CA	5.08	128.62	120.30
1	A	293	ASP	CA-CB-CG	5.04	117.64	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2216	0	2145	25	0
1	B	2094	0	2003	27	0
2	A	35	0	0	0	0
2	B	35	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	14	0	20	1	0
5	B	21	0	30	2	0
6	A	8	0	14	0	0
6	B	16	0	28	5	0
7	A	10	0	14	1	0
8	A	286	0	0	6	0
8	B	226	0	0	3	0
All	All	4973	0	4254	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:LYS:HD3	1:A:261[B]:GLU:OE2	1.67	0.94
1:B:110[B]:MET:HE1	1:B:192[B]:VAL:HG22	1.52	0.90
1:B:110[B]:MET:CE	1:B:192[B]:VAL:HG22	2.13	0.77
5:B:408:PEG:H31	5:B:408:PEG:O1	1.88	0.74
1:B:110[A]:MET:HE3	6:B:404:MPD:H52	1.70	0.72
1:A:131:LEU:HB3	1:A:207[B]:MET:HE1	1.71	0.72
1:A:251:ARG:NH2	8:A:502:HOH:O	2.23	0.71
1:B:110[A]:MET:HE2	1:B:195:ARG:CD	2.24	0.68
1:A:138[B]:VAL:HG21	1:A:289:LYS:HB2	1.76	0.67
5:B:408:PEG:H32	8:B:556:HOH:O	1.94	0.67
1:B:110[B]:MET:HE1	1:B:192[B]:VAL:CG2	2.24	0.67
1:A:107:GLN:OE1	1:A:111[B]:LYS:HE3	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69[B]:LYS:NZ	8:B:501:HOH:O	2.31	0.64
1:A:130[B]:ASP:HB3	8:A:656:HOH:O	1.97	0.64
1:B:72:GLU:O	1:B:78:ARG:NH2	2.29	0.63
1:A:49:GLN:HG2	1:A:50:TRP:O	1.99	0.63
1:B:107[B]:GLN:HB3	1:B:108:PRO:HD3	1.83	0.60
1:B:110[A]:MET:CE	6:B:404:MPD:H52	2.33	0.58
1:A:110[B]:MET:CE	1:A:195:ARG:HG3	2.34	0.57
1:A:138[A]:VAL:HG12	1:A:288:VAL:HG12	1.85	0.56
1:A:75:PRO:HA	1:A:78:ARG:HG2	1.87	0.56
1:A:248[A]:GLU:HG2	1:A:253:LEU:HD11	1.88	0.55
1:B:110[A]:MET:SD	1:B:192[A]:VAL:HG12	2.47	0.55
1:B:100:ASP:HB3	8:B:684[B]:HOH:O	2.08	0.54
1:A:140:ARG:O	7:A:406:PGE:H42	2.08	0.53
1:A:75:PRO:HA	1:A:78:ARG:CG	2.39	0.52
1:B:110[A]:MET:HE2	1:B:195:ARG:HD3	1.91	0.52
1:B:118:PRO:HD3	6:B:405:MPD:HM3	1.91	0.52
1:B:284:PHE:O	1:B:288:VAL:HG23	2.09	0.52
1:B:48:GLN:C	1:B:163[B]:PHE:HE2	2.17	0.51
1:B:68:VAL:O	1:B:72:GLU:HG3	2.11	0.51
1:B:110[A]:MET:HE2	1:B:195:ARG:HD2	1.92	0.51
1:A:218:LYS:NZ	8:A:720[B]:HOH:O	2.44	0.51
1:A:234[A]:LYS:NZ	8:A:510:HOH:O	2.46	0.48
1:B:110[A]:MET:HE3	1:B:110[A]:MET:HA	1.97	0.46
1:A:145:LEU:HD11	1:A:192[B]:VAL:HG12	1.96	0.46
1:B:114:THR:HA	6:B:404:MPD:H51	1.98	0.46
1:A:110[B]:MET:HE2	1:A:195:ARG:NH2	2.31	0.46
1:B:188:PHE:O	1:B:192[B]:VAL:HG23	2.16	0.45
1:B:110[B]:MET:SD	1:B:192[B]:VAL:HG22	2.56	0.45
1:A:278[A]:ILE:CD1	1:A:300:ILE:HG12	2.47	0.45
1:A:49:GLN:HB2	1:A:171:GLN:OE1	2.17	0.45
1:B:48:GLN:O	1:B:163[B]:PHE:HE2	2.01	0.44
1:A:56:THR:HG23	5:A:407:PEG:H31	1.99	0.44
1:A:177[A]:ARG:NH2	8:A:512[A]:HOH:O	2.48	0.43
1:B:218:LYS:HD3	1:B:261:GLU:OE2	2.18	0.43
1:B:107[B]:GLN:HE21	1:B:111:LYS:CG	2.31	0.43
1:A:107:GLN:HE22	1:A:195:ARG:HH12	1.66	0.43
1:A:232:PRO:HB3	1:A:269:ARG:O	2.19	0.42
1:B:90:LYS:CG	1:B:94:ILE:HG13	2.50	0.42
6:B:404:MPD:O4	6:B:404:MPD:HM1	2.20	0.41
1:B:98:PRO:O	1:B:183:ASN:HA	2.20	0.41
1:A:107:GLN:NE2	8:A:519:HOH:O	2.53	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:PHE:HB3	1:A:60:PRO:HD3	2.03	0.40
1:B:48:GLN:C	1:B:163[B]:PHE:CE2	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/257 (109%)	271 (97%)	7 (2%)	1 (0%)	30	12
1	B	265/257 (103%)	254 (96%)	10 (4%)	1 (0%)	30	12
All	All	544/514 (106%)	525 (96%)	17 (3%)	2 (0%)	30	12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	128	ILE
1	A	128	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/237 (106%)	251 (100%)	1 (0%)	84	71
1	B	231/237 (98%)	228 (99%)	3 (1%)	61	35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	483/474 (102%)	479 (99%)	4 (1%)	70 54

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	GLN
1	B	95	SER
1	B	250	SER
1	B	276	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	229	ASN
1	B	74	HIS
1	B	134	GLN
1	B	182	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ROJ	A	401	1	37,38,38	1.16	4 (10%)	49,58,58	1.12	2 (4%)
6	MPD	A	405	-	7,7,7	0.14	0	9,10,10	0.28	0
6	MPD	B	404	-	7,7,7	0.18	0	9,10,10	0.45	0
4	PO4	A	403	-	4,4,4	0.96	0	6,6,6	0.61	0
2	ROJ	B	401	1	37,38,38	1.56	5 (13%)	49,58,58	0.93	2 (4%)
5	PEG	A	407	-	6,6,6	0.25	0	5,5,5	0.17	0
5	PEG	A	404	-	6,6,6	0.15	0	5,5,5	0.08	0
6	MPD	B	405	-	7,7,7	0.16	0	9,10,10	0.42	0
4	PO4	B	403	-	4,4,4	1.78	1 (25%)	6,6,6	0.83	0
5	PEG	B	406	-	6,6,6	0.24	0	5,5,5	0.19	0
7	PGE	A	406	-	9,9,9	0.22	0	8,8,8	0.20	0
5	PEG	B	408	-	6,6,6	0.19	0	5,5,5	0.13	0
5	PEG	B	407	-	6,6,6	0.17	0	5,5,5	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ROJ	A	401	1	-	2/22/51/51	0/4/4/4
6	MPD	A	405	-	-	0/5/5/5	-
6	MPD	B	404	-	-	3/5/5/5	-
5	PEG	A	407	-	-	2/4/4/4	-
2	ROJ	B	401	1	-	2/22/51/51	0/4/4/4
5	PEG	A	404	-	-	2/4/4/4	-
6	MPD	B	405	-	-	2/5/5/5	-
5	PEG	B	406	-	-	2/4/4/4	-
7	PGE	A	406	-	-	1/7/7/7	-
5	PEG	B	408	-	-	1/4/4/4	-
5	PEG	B	407	-	-	3/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	ROJ	P1-O6	5.40	1.65	1.59
2	B	401	ROJ	C10-C13	3.23	1.44	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	ROJ	C11-N4	3.19	1.44	1.35
2	A	401	ROJ	C9-N1	3.15	1.42	1.37
2	B	401	ROJ	C12-N5	2.56	1.38	1.33
4	B	403	PO4	P-O1	2.52	1.56	1.50
2	A	401	ROJ	C1-C4	-2.50	1.46	1.53
2	B	401	ROJ	O11-C14	-2.25	1.37	1.43
2	A	401	ROJ	C12-N5	2.07	1.37	1.33
2	A	401	ROJ	C11-N3	2.02	1.39	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ROJ	O2-C4-C1	5.13	109.37	104.63
2	B	401	ROJ	O7-P2-O8	2.05	121.98	112.44
2	B	401	ROJ	O12-C15-C7	2.03	116.90	111.08
2	A	401	ROJ	N5-C12-N4	-2.00	125.55	128.58

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	404	MPD	O2-C2-C3-C4
6	B	404	MPD	CM-C2-C3-C4
5	A	407	PEG	O2-C3-C4-O4
5	B	407	PEG	O2-C3-C4-O4
5	A	404	PEG	O1-C1-C2-O2
5	A	407	PEG	O1-C1-C2-O2
5	B	407	PEG	O1-C1-C2-O2
5	A	404	PEG	C1-C2-O2-C3
5	B	408	PEG	O2-C3-C4-O4
2	B	401	ROJ	C14-C8-N1-C9
5	B	406	PEG	C4-C3-O2-C2
6	B	405	MPD	C2-C3-C4-C5
2	A	401	ROJ	C14-C8-N1-C9
7	A	406	PGE	C4-C3-O2-C2
5	B	407	PEG	C4-C3-O2-C2
2	A	401	ROJ	O10-C8-N1-C9
2	B	401	ROJ	O10-C8-N1-C9
6	B	405	MPD	C2-C3-C4-O4
6	B	404	MPD	C1-C2-C3-C4
5	B	406	PEG	O1-C1-C2-O2


There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	404	MPD	4	0
5	A	407	PEG	1	0
6	B	405	MPD	1	0
7	A	406	PGE	1	0
5	B	408	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/257 (99%)	0.20	19 (7%) 20 22	10, 24, 48, 79	26 (10%)
1	B	251/257 (97%)	0.58	28 (11%) 10 10	13, 31, 78, 125	16 (6%)
All	All	506/514 (98%)	0.39	47 (9%) 14 15	10, 27, 64, 125	42 (8%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163[A]	PHE	7.5
1	A	46	TRP	7.1
1	B	46	TRP	7.0
1	A	247[A]	ARG	6.5
1	A	300	ILE	4.8
1	A	164	ALA	4.3
1	B	164[A]	ALA	4.3
1	A	246[A]	GLY	4.0
1	B	296	CYS	3.7
1	B	176[A]	TRP	3.7
1	B	295	SER	3.3
1	B	287	CYS	3.2
1	A	296	CYS	3.1
1	B	137	GLN	3.1
1	A	299	GLU	3.0
1	B	293	ASP	3.0
1	B	100	ASP	2.9
1	B	294	SER	2.9
1	A	48	GLN	2.8
1	B	143	PHE	2.8
1	B	291	PRO	2.8
1	B	162[A]	GLU	2.8
1	A	49	GLN	2.7
1	A	294	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	298	SER	2.6
1	A	297	THR	2.5
1	B	73	ILE	2.5
1	A	249[A]	ASP	2.5
1	B	138	VAL	2.5
1	B	279	TYR	2.4
1	B	141	ASP	2.4
1	B	248	GLU	2.3
1	B	247	ARG	2.3
1	A	290[A]	ASN	2.3
1	B	288	VAL	2.2
1	B	286	GLN	2.2
1	A	195	ARG	2.2
1	A	73	ILE	2.2
1	A	47	ARG	2.2
1	B	292	GLU	2.2
1	B	246	GLY	2.1
1	B	119	CYS	2.1
1	B	47	ARG	2.1
1	A	248[A]	GLU	2.1
1	B	117	VAL	2.0
1	B	290	ASN	2.0
1	A	176[A]	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	B	406	7/7	0.57	0.25	71,75,85,85	0
5	PEG	A	404	7/7	0.68	0.19	53,57,61,64	0
5	PEG	A	407	7/7	0.76	0.17	37,49,55,59	0
5	PEG	B	407	7/7	0.76	0.19	70,77,84,86	0
6	MPD	B	405	8/8	0.76	0.20	73,74,75,77	0
7	PGE	A	406	10/10	0.77	0.15	45,49,54,55	0
6	MPD	B	404	8/8	0.78	0.19	58,60,62,63	0
6	MPD	A	405	8/8	0.82	0.17	39,49,56,57	0
5	PEG	B	408	7/7	0.82	0.14	35,50,63,66	0
4	PO4	B	403	5/5	0.96	0.07	28,33,38,40	0
4	PO4	A	403	5/5	0.96	0.07	22,24,29,35	0
3	K	B	402	1/1	0.98	0.11	33,33,33,33	0
2	ROJ	B	401	35/35	0.98	0.05	19,21,26,27	0
2	ROJ	A	401	35/35	0.99	0.04	16,18,20,22	0
3	K	A	402	1/1	0.99	0.10	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ROJ B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.