



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:48 PM UTC

PDB ID : 6VVD / pdb_00006vvd
Title : Legionella pneumophila Lpg2603 kinase bound to IP6
Authors : Tomchick, D.R.; Tagliabracci, V.S.; Sreelatha, A.
Deposited on : 2020-02-17
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	
1	B	314	
1	C	314	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 14090 atoms, of which 6875 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dot/Icm T4SS effector.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	279	Total 4654	C 1528	H 2282	N 399	O 440	S 5	0	0	0
1	B	282	Total 4719	C 1549	H 2317	N 402	O 446	S 5	0	0	0
1	C	275	Total 4591	C 1510	H 2258	N 388	O 430	S 5	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	SER	-	expression tag	UNP A0A2S6F2W2
B	9	SER	-	expression tag	UNP A0A2S6F2W2
C	9	SER	-	expression tag	UNP A0A2S6F2W2

- Molecule 2 is INOSITOL HEXAKISPHOSPHATE (CCD ID: IHP) (formula: $C_6H_{18}O_{24}P_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 42	C 6	H 6	O 24	P 6	0	0
2	B	1	Total 42	C 6	H 6	O 24	P 6	0	0
2	C	1	Total 42	C 6	H 6	O 24	P 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dot/Icm T4SS effector




- Molecule 1: Dot/Icm T4SS effector



- Molecule 1: Dot/Icm T4SS effector





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.93Å 73.21Å 261.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.54 – 2.65 43.54 – 2.65	Depositor EDS
% Data completeness (in resolution range)	87.0 (43.54-2.65) 87.0 (43.54-2.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.17.1	Depositor
R, R_{free}	0.256 , 0.281 0.258 , 0.281	Depositor DCC
R_{free} test set	2056 reflections (6.77%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	14090	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/2441	0.30	0/3304
1	B	0.17	0/2470	0.34	0/3341
1	C	0.15	0/2399	0.30	0/3245
All	All	0.16	0/7310	0.32	0/9890

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2372	2282	2282	6	1
1	B	2402	2317	2317	16	2
1	C	2333	2258	2258	10	1
2	A	36	6	6	0	0
2	B	36	6	6	0	0
2	C	36	6	6	0	0
All	All	7215	6875	6875	32	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:54:GLU:OE1	1:C:70:LEU:HD11	1.79	0.82
1:B:39:LEU:HD22	1:B:130:LEU:HD12	1.75	0.67
1:C:125:TYR:CD2	1:C:161:VAL:HG11	2.35	0.61
1:B:209:ILE:HG21	1:B:258:PHE:CE1	2.36	0.61
1:B:69:VAL:HG12	1:B:140:LYS:HD2	1.83	0.60
1:C:25:ILE:HD11	1:C:58:VAL:HG13	1.85	0.58
1:C:25:ILE:HD13	1:C:68:TYR:HB3	1.88	0.54
1:A:61:GLU:HB3	1:A:66:MET:HG3	1.92	0.52
1:B:39:LEU:HD13	1:B:43:ASN:HD22	1.75	0.51
1:B:173:ILE:HG12	1:B:216:ALA:HB2	1.93	0.50
1:B:96:LEU:HD12	1:B:106:GLU:HB2	1.94	0.49
1:C:25:ILE:HD11	1:C:58:VAL:CG1	2.44	0.48
1:B:39:LEU:HD13	1:B:43:ASN:ND2	2.29	0.47
1:A:67:ASP:OD1	1:A:68:TYR:N	2.47	0.46
1:B:107:LYS:HG3	1:B:108:THR:HG23	1.97	0.45
1:B:192:LEU:HG	1:B:197:TYR:HB2	1.98	0.45
1:B:192:LEU:HD21	1:B:230:PHE:CE1	2.52	0.45
1:B:132:ILE:HG21	1:B:151:GLY:O	2.17	0.44
1:B:179:GLN:O	1:B:183:GLU:HG3	2.17	0.44
1:A:61:GLU:HB3	1:A:66:MET:CG	2.48	0.43
1:A:68:TYR:O	1:A:140:LYS:NZ	2.45	0.43
1:C:209:ILE:HG21	1:C:258:PHE:CE1	2.54	0.43
1:C:201:ASP:O	1:C:213:ASN:ND2	2.51	0.42
1:C:288:LEU:HB3	1:C:295:LEU:HD21	2.00	0.42
1:C:56:TYR:CE1	1:C:70:LEU:HD13	2.55	0.42
1:B:171:ILE:O	1:B:216:ALA:HB3	2.19	0.41
1:A:71:MET:HE2	1:A:152:LEU:HG	2.03	0.41
1:B:193:ASN:HA	1:B:197:TYR:O	2.21	0.40
1:C:63:ASP:OD1	1:C:64:GLU:N	2.55	0.40
1:A:193:ASN:HA	1:A:197:TYR:O	2.22	0.40
1:B:69:VAL:HG12	1:B:140:LYS:CD	2.48	0.40
1:B:46:PHE:CG	1:B:133:ILE:HG21	2.56	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:ARG:HH11	1:B:172:PRO:O[3_654]	1.41	0.19
1:B:119:ARG:NH1	1:B:172:PRO:O[3_654]	2.17	0.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:PRO:O	1:C:119:ARG:HH11[1_455]	1.59	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/314 (88%)	261 (95%)	14 (5%)	0	100	100
1	B	276/314 (88%)	265 (96%)	11 (4%)	0	100	100
1	C	267/314 (85%)	254 (95%)	13 (5%)	0	100	100
All	All	818/942 (87%)	780 (95%)	38 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/292 (88%)	257 (100%)	1 (0%)	84	93
1	B	262/292 (90%)	259 (99%)	3 (1%)	65	80
1	C	254/292 (87%)	251 (99%)	3 (1%)	63	79
All	All	774/876 (88%)	767 (99%)	7 (1%)	70	82

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	227	ASP
1	B	117	ARG
1	B	130	LEU
1	B	175	LYS
1	C	127	GLU
1	C	130	LEU
1	C	164	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	204	ASN
1	A	268	HIS
1	A	291	ASN
1	C	62	ASN
1	C	204	ASN
1	C	291	ASN
1	C	293	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IHP	A	401	-	36,36,36	1.48	6 (16%)	60,60,60	0.93	3 (5%)
2	IHP	B	401	-	36,36,36	1.53	6 (16%)	60,60,60	0.93	4 (6%)
2	IHP	C	401	-	36,36,36	1.63	6 (16%)	60,60,60	0.92	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IHP	A	401	-	-	2/30/54/54	0/1/1/1
2	IHP	B	401	-	-	8/30/54/54	0/1/1/1
2	IHP	C	401	-	-	5/30/54/54	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	IHP	P2-O12	3.84	1.66	1.59
2	C	401	IHP	P1-O11	3.72	1.66	1.59
2	B	401	IHP	P4-O14	3.44	1.65	1.59
2	C	401	IHP	P6-O16	3.40	1.65	1.59
2	B	401	IHP	P2-O12	3.37	1.65	1.59
2	B	401	IHP	P1-O11	3.37	1.65	1.59
2	A	401	IHP	P4-O14	3.33	1.65	1.59
2	C	401	IHP	P5-O15	3.33	1.65	1.59
2	B	401	IHP	P5-O15	3.28	1.65	1.59
2	C	401	IHP	P4-O14	3.27	1.65	1.59
2	A	401	IHP	P3-O13	3.26	1.65	1.59
2	C	401	IHP	P3-O13	3.18	1.65	1.59
2	A	401	IHP	P2-O12	3.16	1.65	1.59
2	A	401	IHP	P1-O11	3.15	1.65	1.59
2	B	401	IHP	P3-O13	3.09	1.64	1.59
2	A	401	IHP	P5-O15	3.05	1.64	1.59
2	B	401	IHP	P6-O16	2.88	1.64	1.59
2	A	401	IHP	P6-O16	2.75	1.64	1.59

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	IHP	P4-O14-C4	-2.68	116.26	123.43
2	A	401	IHP	P2-O12-C2	-2.57	116.56	123.43
2	C	401	IHP	P5-O15-C5	-2.56	116.60	123.43
2	A	401	IHP	P5-O15-C5	-2.48	116.81	123.43
2	B	401	IHP	P6-O16-C6	-2.39	117.04	123.43
2	B	401	IHP	P5-O15-C5	-2.37	117.10	123.43
2	C	401	IHP	P6-O16-C6	-2.32	117.25	123.43
2	B	401	IHP	P4-O14-C4	-2.23	117.48	123.43
2	C	401	IHP	P2-O12-C2	-2.22	117.51	123.43
2	B	401	IHP	P1-O11-C1	-2.18	117.61	123.43

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	401	IHP	C2-C1-O11-P1
2	B	401	IHP	C4-O14-P4-O24
2	B	401	IHP	C6-O16-P6-O26
2	B	401	IHP	C5-O15-P5-O45
2	C	401	IHP	C1-O11-P1-O41
2	C	401	IHP	C5-O15-P5-O45
2	B	401	IHP	C6-C1-O11-P1
2	B	401	IHP	C3-C2-O12-P2
2	A	401	IHP	C5-O15-P5-O25
2	B	401	IHP	C5-O15-P5-O25
2	C	401	IHP	C5-O15-P5-O25
2	C	401	IHP	C2-C1-O11-P1
2	A	401	IHP	C5-O15-P5-O35
2	C	401	IHP	C3-O13-P3-O43
2	B	401	IHP	C1-O11-P1-O21


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/314 (88%)	0.60	34 (12%) 8 7	12, 47, 101, 151	0
1	B	282/314 (89%)	0.63	24 (8%) 16 12	12, 52, 94, 107	0
1	C	275/314 (87%)	2.02	130 (47%) 0 0	42, 92, 136, 215	0
All	All	836/942 (88%)	1.08	188 (22%) 2 1	12, 63, 122, 215	0

All (188) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	173	ILE	8.1
1	C	295	LEU	6.8
1	C	172	PRO	6.3
1	C	296	LEU	6.1
1	C	25	ILE	5.6
1	C	237	VAL	5.6
1	C	263	THR	5.6
1	C	288	LEU	5.4
1	A	317	THR	5.1
1	C	99	GLY	5.0
1	C	293	HIS	4.8
1	C	171	ILE	4.8
1	C	303	LEU	4.7
1	C	302	GLY	4.7
1	C	300	HIS	4.7
1	C	276	ILE	4.7
1	A	158	GLU	4.6
1	C	245	LEU	4.6
1	A	62	ASN	4.3
1	A	314	LEU	4.2
1	C	243	LYS	4.2
1	C	233	ASP	4.1
1	C	279	TRP	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	270	ILE	4.0
1	C	304	ILE	4.0
1	C	297	ARG	4.0
1	C	259	ILE	3.9
1	A	157	ASN	3.9
1	C	282	ASN	3.9
1	C	277	GLU	3.8
1	C	278	LYS	3.8
1	C	292	PRO	3.7
1	A	61	GLU	3.7
1	C	246	VAL	3.7
1	C	244	SER	3.6
1	A	63	ASP	3.6
1	C	112	ALA	3.5
1	C	274	VAL	3.5
1	C	249	LYS	3.5
1	C	110	ASP	3.5
1	C	64	GLU	3.5
1	C	109	GLU	3.5
1	C	286	GLU	3.4
1	A	315	HIS	3.4
1	A	237	VAL	3.4
1	C	298	PHE	3.4
1	C	105	TYR	3.4
1	A	64	GLU	3.4
1	C	113	MET	3.4
1	C	240	LEU	3.3
1	C	40	SER	3.3
1	C	253	LEU	3.3
1	C	287	SER	3.3
1	C	180	PRO	3.2
1	B	63	ASP	3.2
1	C	182	ILE	3.1
1	A	235	GLY	3.1
1	C	63	ASP	3.0
1	C	238	ASP	3.0
1	B	173	ILE	3.0
1	A	159	GLU	3.0
1	A	25	ILE	3.0
1	C	236	ARG	3.0
1	C	214	LEU	3.0
1	C	219	GLU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	101	GLN	3.0
1	B	108	THR	3.0
1	C	235	GLY	3.0
1	C	314	LEU	2.9
1	B	265	GLU	2.9
1	C	92	ASP	2.9
1	C	177	ILE	2.9
1	C	256	TYR	2.9
1	C	299	TYR	2.9
1	A	313	ASP	2.9
1	A	294	THR	2.9
1	C	283	ASN	2.8
1	C	285	ASP	2.8
1	A	308	LYS	2.8
1	A	240	LEU	2.8
1	C	215	ILE	2.8
1	C	108	THR	2.8
1	C	55	VAL	2.8
1	C	221	VAL	2.8
1	B	96	LEU	2.8
1	C	61	GLU	2.8
1	C	24	LYS	2.8
1	C	318	ILE	2.8
1	A	30	ASN	2.7
1	C	257	ASN	2.7
1	C	209	ILE	2.7
1	C	53	ASN	2.7
1	C	72	TRP	2.7
1	C	290	ASP	2.7
1	C	258	PHE	2.7
1	C	56	TYR	2.6
1	C	247	TYR	2.6
1	A	236	ARG	2.6
1	C	176	LYS	2.6
1	A	241	SER	2.6
1	C	275	PRO	2.6
1	C	294	THR	2.6
1	C	178	ASP	2.6
1	C	168	ILE	2.6
1	C	306	LEU	2.6
1	C	41	GLU	2.6
1	C	181	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	211	PRO	2.5
1	C	291	ASN	2.5
1	B	240	LEU	2.5
1	C	315	HIS	2.5
1	B	102	GLY	2.5
1	C	65	PRO	2.5
1	C	307	PRO	2.5
1	C	66	MET	2.5
1	A	284	ARG	2.5
1	C	231	ARG	2.5
1	C	262	PRO	2.5
1	C	42	PRO	2.4
1	B	61	GLU	2.4
1	C	301	GLU	2.4
1	C	174	ASN	2.4
1	A	65	PRO	2.4
1	C	94	LYS	2.4
1	C	98	LYS	2.4
1	C	280	TYR	2.4
1	B	268	HIS	2.4
1	C	289	SER	2.4
1	C	271	ASP	2.4
1	B	98	LYS	2.4
1	C	170	ALA	2.4
1	A	267	ASP	2.3
1	C	68	TYR	2.3
1	A	312	HIS	2.3
1	B	30	ASN	2.3
1	A	297	ARG	2.3
1	C	28	ILE	2.3
1	A	293	HIS	2.3
1	C	308	LYS	2.3
1	C	242	ARG	2.3
1	B	104	VAL	2.3
1	C	186	VAL	2.3
1	B	112	ALA	2.3
1	C	111	LYS	2.3
1	C	241	SER	2.3
1	C	104	VAL	2.3
1	B	66	MET	2.2
1	C	93	TYR	2.2
1	B	42	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	166	PRO	2.2
1	C	309	LYS	2.2
1	B	177	ILE	2.2
1	C	45	GLU	2.2
1	C	106	GLU	2.2
1	A	296	LEU	2.2
1	C	118	GLY	2.2
1	C	216	ALA	2.2
1	C	187	LEU	2.2
1	C	305	SER	2.2
1	C	75	HIS	2.2
1	C	60	PHE	2.2
1	A	311	ALA	2.1
1	B	93	TYR	2.1
1	A	310	LEU	2.1
1	B	267	ASP	2.1
1	A	41	GLU	2.1
1	C	183	GLU	2.1
1	C	284	ARG	2.1
1	A	141	TYR	2.1
1	C	96	LEU	2.1
1	C	239	GLU	2.1
1	A	121	PRO	2.1
1	B	270	ILE	2.1
1	C	51	TYR	2.1
1	C	141	TYR	2.1
1	C	207	TYR	2.1
1	A	119	ARG	2.0
1	C	273	ARG	2.0
1	B	40	SER	2.0
1	A	295	LEU	2.0
1	B	319	LEU	2.0
1	C	179	GLN	2.0
1	C	95	GLU	2.0
1	B	24	LYS	2.0
1	C	73	LYS	2.0
1	C	74	SER	2.0
1	C	248	GLY	2.0
1	B	269	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	IHP	C	401	36/36	0.81	0.12	66,87,108,116	0
2	IHP	B	401	36/36	0.86	0.11	59,69,86,88	0
2	IHP	A	401	36/36	0.90	0.10	41,52,63,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IHP C 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IHP B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.