



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:47 AM UTC

PDB ID : 6VVI / pdb_00006vvi
Title : Arabidopsis thaliana dihydrodipicolinate synthase isoform 1 (DHDPS1)
Authors : Lee, M.; Hall, C.J.
Deposited on : 2020-02-18
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	321	 2% 91% 5% .
1	BBB	321	 % 93% . . .
1	CCC	321	 % 88% 7% . .
1	DDD	321	 % 92% 5% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	DDD	404	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxy-tetrahydrodipicolinate synthase 1, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	308	2401	1514	425	448	14	0	1	0
1	BBB	312	2455	1545	436	460	14	0	4	0
1	CCC	309	2413	1522	425	452	14	0	2	0
1	DDD	311	2419	1524	426	455	14	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	45	GLY	-	expression tag	UNP Q9LZX6
AAA	46	PRO	-	expression tag	UNP Q9LZX6
AAA	47	GLY	-	expression tag	UNP Q9LZX6
AAA	48	SER	-	expression tag	UNP Q9LZX6
BBB	45	GLY	-	expression tag	UNP Q9LZX6
BBB	46	PRO	-	expression tag	UNP Q9LZX6
BBB	47	GLY	-	expression tag	UNP Q9LZX6
BBB	48	SER	-	expression tag	UNP Q9LZX6
CCC	45	GLY	-	expression tag	UNP Q9LZX6
CCC	46	PRO	-	expression tag	UNP Q9LZX6
CCC	47	GLY	-	expression tag	UNP Q9LZX6
CCC	48	SER	-	expression tag	UNP Q9LZX6
DDD	45	GLY	-	expression tag	UNP Q9LZX6
DDD	46	PRO	-	expression tag	UNP Q9LZX6
DDD	47	GLY	-	expression tag	UNP Q9LZX6
DDD	48	SER	-	expression tag	UNP Q9LZX6

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total C O 6 3 3	0	0
2	CCC	1	Total C O 6 3 3	0	0
2	DDD	1	Total C O 6 3 3	0	0
2	DDD	1	Total C O 6 3 3	0	0

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total Na 1 1	0	0
3	BBB	1	Total Na 1 1	0	0
3	CCC	1	Total Na 1 1	0	0
3	DDD	1	Total Na 1 1	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	BBB	1	Total	O	S	0	0
			5	4	1		
4	CCC	1	Total	O	S	0	0
			5	4	1		
4	CCC	1	Total	O	S	0	0
			5	4	1		
4	DDD	1	Total	O	S	0	0
			5	4	1		
4	DDD	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	85	Total	O	0	0
			85	85		
5	BBB	130	Total	O	0	0
			130	130		
5	CCC	107	Total	O	0	0
			107	107		
5	DDD	143	Total	O	0	0
			143	143		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase 1, chloroplastic




- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase 1, chloroplastic



- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase 1, chloroplastic



- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase 1, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	95.47Å 98.10Å 176.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.10 – 2.15 49.10 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.10-2.15) 99.3 (49.10-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 2.14Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.172 , 0.209 0.180 , 0.214	Depositor DCC
R_{free} test set	4371 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtrriage
Anisotropy	0.505	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.032 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10206	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.98	0/2456	1.39	1/3332 (0.0%)
1	BBB	0.98	0/2519	1.33	0/3416
1	CCC	0.99	0/2472	1.37	2/3355 (0.1%)
1	DDD	1.01	0/2474	1.35	1/3358 (0.0%)
All	All	0.99	0/9921	1.36	4/13461 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	CCC	347	GLY	CA-C-O	-6.33	116.75	122.57
1	AAA	297	GLN	CB-CA-C	-6.18	100.92	110.81
1	DDD	347	GLY	CA-C-O	-5.41	117.91	122.29
1	CCC	120	ILE	N-CA-CB	5.24	117.66	110.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2401	0	2368	6	0
1	BBB	2455	0	2423	8	0
1	CCC	2413	0	2375	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	DDD	2419	0	2376	8	0
2	AAA	6	0	8	1	0
2	CCC	6	0	8	3	0
2	DDD	12	0	16	0	0
3	AAA	1	0	0	0	0
3	BBB	1	0	0	0	0
3	CCC	1	0	0	0	0
3	DDD	1	0	0	0	0
4	BBB	5	0	0	0	0
4	CCC	10	0	0	1	0
4	DDD	10	0	0	2	0
5	AAA	85	0	0	0	0
5	BBB	130	0	0	2	0
5	CCC	107	0	0	1	0
5	DDD	143	0	0	1	0
All	All	10206	0	9574	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:CCC:116:TRP:CZ2	1:CCC:120:ILE:HG23	2.13	0.83
1:BBB:227:ASN:ND2	1:BBB:251:SER:OG	2.20	0.75
1:DDD:227:ASN:ND2	1:DDD:251:SER:OG	2.22	0.73
1:DDD:213:SER:HA	1:DDD:218:MET:HG2	1.74	0.68
1:CCC:218:MET:O	1:CCC:218:MET:HG3	1.94	0.68
1:CCC:213:SER:HA	1:CCC:218:MET:HG2	1.76	0.67
1:CCC:92:ASN:HD21	1:CCC:133:ARG:HH12	1.47	0.62
1:BBB:119:HIS:ND1	5:BBB:501:HOH:O	2.32	0.58
1:DDD:351:VAL:HG12	4:DDD:404:SO4:O2	2.07	0.55
1:AAA:117:ASP:HB3	5:CCC:601:HOH:O	2.07	0.54
1:CCC:143:ASN:HD21	2:CCC:401:GOL:C3	2.20	0.54
1:CCC:116:TRP:CZ2	1:CCC:151:ALA:HA	2.43	0.54
1:BBB:280[B]:ARG:NH2	1:BBB:282:SER:OG	2.42	0.53
1:BBB:98:GLY:HA3	1:BBB:273:ARG:HD3	1.91	0.52
1:AAA:143:ASN:HD21	2:AAA:401:GOL:C3	2.23	0.51
1:DDD:150:HIS:ND1	5:DDD:504:HOH:O	2.34	0.51
1:CCC:116:TRP:O	1:CCC:120:ILE:HG13	2.11	0.50
1:DDD:178:MET:HE2	1:DDD:202:GLN:OE1	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:120:ILE:HG22	1:CCC:120:ILE:HD13	1.94	0.50
1:CCC:120:ILE:HB	1:CCC:158:MET:HE1	1.94	0.49
1:BBB:124:GLY:HA3	1:BBB:158:MET:HE3	1.95	0.49
1:AAA:267:LEU:HD11	1:AAA:295:LEU:HD21	1.96	0.47
1:CCC:78:PRO:O	1:CCC:362:ILE:HD11	2.16	0.46
1:AAA:294:TRP:CZ2	1:AAA:333:PHE:HB2	2.52	0.44
1:DDD:120:ILE:O	1:DDD:158:MET:HE1	2.18	0.44
1:CCC:212:LEU:HB2	1:CCC:218:MET:HE2	1.99	0.43
1:BBB:120:ILE:O	1:BBB:158:MET:HE1	2.19	0.43
1:CCC:294:TRP:CZ2	1:CCC:333:PHE:HB2	2.53	0.43
1:CCC:143:ASN:ND2	2:CCC:401:GOL:H31	2.34	0.42
1:CCC:160:MET:HE3	1:CCC:163:ALA:HB2	2.00	0.42
1:DDD:351:VAL:N	4:DDD:404:SO4:O2	2.38	0.42
1:AAA:300:ASN:OD1	1:AAA:301:PRO:HA	2.18	0.42
1:CCC:120:ILE:O	1:CCC:158:MET:HE1	2.19	0.42
1:CCC:143:ASN:HD21	2:CCC:401:GOL:H31	1.83	0.41
1:CCC:60:THR:O	1:CCC:64:ARG:HG3	2.19	0.41
1:BBB:68:VAL:HG21	1:BBB:273:ARG:HA	2.02	0.41
1:CCC:67:ARG:NE	4:CCC:404:SO4:O2	2.42	0.40
1:BBB:71:ALA:HB1	5:BBB:599:HOH:O	2.20	0.40
1:DDD:289:LEU:N	1:DDD:290:PRO:CD	2.84	0.40
1:CCC:68:VAL:HG21	1:CCC:273:ARG:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	307/321 (96%)	301 (98%)	6 (2%)	0	100	100
1	BBB	314/321 (98%)	309 (98%)	5 (2%)	0	100	100
1	CCC	309/321 (96%)	303 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	DDD	310/321 (97%)	306 (99%)	4 (1%)	0	100	100
All	All	1240/1284 (97%)	1219 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	261/272 (96%)	255 (98%)	6 (2%)	44	47
1	BBB	268/272 (98%)	263 (98%)	5 (2%)	50	55
1	CCC	263/272 (97%)	257 (98%)	6 (2%)	44	47
1	DDD	263/272 (97%)	261 (99%)	2 (1%)	73	79
All	All	1055/1088 (97%)	1036 (98%)	19 (2%)	51	57

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	102	VAL
1	AAA	201	CYS
1	AAA	207	GLN
1	AAA	218	MET
1	AAA	251	SER
1	AAA	342	ARG
1	BBB	54	ASP
1	BBB	201	CYS
1	BBB	251	SER
1	BBB	273	ARG
1	BBB	365	TYR
1	CCC	92	ASN
1	CCC	102	VAL
1	CCC	120	ILE
1	CCC	201	CYS
1	CCC	218	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	CCC	251	SER
1	DDD	201	CYS
1	DDD	301	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	DDD	401	-	5,5,5	0.27	0	5,5,5	0.62	0
4	SO4	BBB	402	-	4,4,4	0.32	0	6,6,6	0.07	0
4	SO4	CCC	404	-	4,4,4	0.39	0	6,6,6	0.13	0
4	SO4	CCC	403	-	4,4,4	0.36	0	6,6,6	0.11	0
4	SO4	DDD	404	-	4,4,4	0.35	0	6,6,6	0.16	0
2	GOL	CCC	401	-	5,5,5	0.16	0	5,5,5	0.35	0
2	GOL	DDD	402	-	5,5,5	0.13	0	5,5,5	0.40	0
2	GOL	AAA	401	-	5,5,5	0.12	0	5,5,5	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	DDD	405	-	4,4,4	0.40	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	DDD	402	-	-	4/4/4/4	-
2	GOL	DDD	401	-	-	3/4/4/4	-
2	GOL	CCC	401	-	-	2/4/4/4	-
2	GOL	AAA	401	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	CCC	401	GOL	O1-C1-C2-C3
2	DDD	401	GOL	C1-C2-C3-O3
2	DDD	402	GOL	O1-C1-C2-C3
2	DDD	402	GOL	C1-C2-C3-O3
2	DDD	401	GOL	O2-C2-C3-O3
2	AAA	401	GOL	O1-C1-C2-C3
2	AAA	401	GOL	O1-C1-C2-O2
2	CCC	401	GOL	O1-C1-C2-O2
2	DDD	402	GOL	O1-C1-C2-O2
2	DDD	402	GOL	O2-C2-C3-O3
2	DDD	401	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	CCC	404	SO4	1	0
4	DDD	404	SO4	2	0
2	CCC	401	GOL	3	0
2	AAA	401	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	308/321 (95%)	0.14	5 (1%) 70 74	26, 45, 64, 88	1 (0%)
1	BBB	312/321 (97%)	-0.17	3 (0%) 79 83	27, 38, 54, 98	4 (1%)
1	CCC	309/321 (96%)	0.02	3 (0%) 79 83	25, 43, 59, 85	2 (0%)
1	DDD	311/321 (96%)	-0.06	4 (1%) 75 79	30, 40, 58, 82	1 (0%)
All	All	1240/1284 (96%)	-0.02	15 (1%) 76 80	25, 41, 61, 98	8 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	58	THR	5.7
1	DDD	55	LYS	5.6
1	CCC	57	ARG	3.5
1	CCC	116	TRP	3.4
1	AAA	116	TRP	3.3
1	BBB	116	TRP	3.1
1	BBB	54	ASP	2.8
1	CCC	58	THR	2.8
1	DDD	338	LYS	2.6
1	DDD	365	TYR	2.6
1	AAA	59	ASN	2.5
1	AAA	365	TYR	2.4
1	AAA	339	GLU	2.3
1	DDD	287	LYS	2.1
1	BBB	365	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	CCC	401	6/6	0.81	0.16	51,58,65,65	0
4	SO4	DDD	404	5/5	0.82	0.15	48,49,50,54	5
2	GOL	AAA	401	6/6	0.83	0.13	48,54,57,62	0
4	SO4	CCC	403	5/5	0.87	0.12	49,52,54,54	5
4	SO4	BBB	402	5/5	0.88	0.16	51,51,54,56	5
4	SO4	DDD	405	5/5	0.91	0.15	41,42,45,53	5
2	GOL	DDD	401	6/6	0.92	0.12	48,55,59,61	0
2	GOL	DDD	402	6/6	0.92	0.12	56,62,63,64	0
4	SO4	CCC	404	5/5	0.94	0.11	39,41,42,43	5
3	NA	BBB	401	1/1	0.97	0.05	37,37,37,37	0
3	NA	AAA	402	1/1	0.97	0.11	40,40,40,40	0
3	NA	DDD	403	1/1	0.98	0.04	33,33,33,33	0
3	NA	CCC	402	1/1	0.98	0.10	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.