



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:59 PM UTC

PDB ID : 6VWF / pdb_00006vwf
Title : Structure of ALDH9A1 complexed with NAD⁺ in space group C222
Authors : Wyatt, J.W.; Tanner, J.J.
Deposited on : 2020-02-19
Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2053 (2.66-2.62)
Clashscore	190562	2097 (2.66-2.62)
Ramachandran outliers	187476	2066 (2.66-2.62)
Sidechain outliers	187428	2066 (2.66-2.62)
RSRZ outliers	180081	2052 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	494	 4% 84% 7% • 9%
1	B	494	 3% 81% 9% • 9%

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total 10	O 10	0	0
3	B	21	Total 21	O 21	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	158.15Å 163.47Å 84.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.13 – 2.64 47.13 – 2.64	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.13-2.64) 99.9 (47.13-2.64)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.14	Depositor
R, R_{free}	0.218 , 0.273 0.225 , 0.277	Depositor DCC
R_{free} test set	1596 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtrriage
Anisotropy	0.388	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.038 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6806	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/3429	0.30	0/4647
1	B	0.11	0/3428	0.30	0/4650
All	All	0.11	0/6857	0.30	0/9297

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3365	0	3245	20	0
1	B	3364	0	3248	23	0
2	A	23	0	12	0	0
2	B	23	0	12	0	0
3	A	10	0	0	0	0
3	B	21	0	0	0	0
All	All	6806	0	6517	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:GLY:HA2	1:A:374:ARG:HB2	1.67	0.76
1:B:365:PRO:HA	1:B:368:LYS:HD2	1.70	0.74
1:A:409:LEU:HD22	1:A:437:GLU:HG2	1.73	0.70
1:A:1:MET:HE2	1:A:6:PHE:HB2	1.75	0.69
1:B:266:ASP:OD2	1:B:425:THR:OG1	2.18	0.62
1:B:468:GLU:OE2	1:B:471:ARG:NH2	2.34	0.60
1:A:12:LEU:HD12	1:A:43:PHE:HB3	1.83	0.60
1:A:101:LYS:HE2	1:A:283:THR:HG23	1.88	0.56
1:B:427:ASP:HB3	1:B:430:ARG:HB3	1.90	0.54
1:A:258:LYS:HD3	1:A:292:THR:HG23	1.89	0.54
1:B:284:GLN:HA	1:B:327:MET:HE2	1.90	0.53
1:B:101:LYS:HE2	1:B:283:THR:HG23	1.89	0.53
1:A:284:GLN:HA	1:A:327:MET:HE2	1.93	0.51
1:B:12:LEU:HD12	1:B:43:PHE:HB3	1.93	0.49
1:B:168:ALA:HB3	1:B:169:PRO:HD3	1.94	0.49
1:A:356:GLY:CA	1:A:374:ARG:HB2	2.40	0.48
1:A:297:GLN:HG3	1:A:402:PHE:CE2	2.48	0.48
1:A:409:LEU:HD21	1:A:434:VAL:HG22	1.96	0.48
1:A:111:ASP:HA	1:A:114:TRP:NE1	2.31	0.46
1:B:142:THR:HG22	1:B:482:THR:HG22	1.96	0.46
1:B:356:GLY:HA2	1:B:374:ARG:HB2	1.98	0.46
1:A:107:ARG:NH1	1:A:323:GLU:OE1	2.41	0.45
1:A:359:ILE:HD12	1:A:359:ILE:H	1.80	0.45
1:A:284:GLN:HG3	1:A:327:MET:HG3	1.99	0.45
1:B:81:ALA:HB2	1:B:117:LEU:HB3	1.98	0.45
1:B:319:ASP:HB3	1:B:322:LEU:HG	1.98	0.44
1:B:35:ALA:HA	1:B:329:PRO:HG3	1.99	0.44
1:B:147:LEU:N	1:B:175:ASN:OD1	2.49	0.43
1:A:51:VAL:HG13	1:A:219:LEU:HD23	2.00	0.43
1:A:84:ILE:HG13	1:A:195:ILE:HG21	1.99	0.43
1:B:297:GLN:HG3	1:B:402:PHE:CE2	2.53	0.43
1:B:111:ASP:HA	1:B:114:TRP:NE1	2.34	0.42
1:A:319:ASP:HB3	1:A:322:LEU:HG	2.01	0.42
1:A:35:ALA:HA	1:A:329:PRO:HG3	2.02	0.42
1:A:480:LEU:HD12	1:A:480:LEU:HA	1.89	0.41
1:B:84:ILE:HG13	1:B:195:ILE:HG21	2.02	0.41
1:B:145:GLU:OE1	1:B:481:LYS:HD2	2.20	0.41
1:B:280:ASN:OD1	1:B:292:THR:HA	2.21	0.41
1:A:145:GLU:OE1	1:A:481:LYS:HD2	2.20	0.41
1:B:326:ARG:HD3	1:B:326:ARG:HA	1.74	0.41
1:B:173:CYS:SG	1:B:474:ILE:HG23	2.61	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:ASN:HB3	1:B:292:THR:HG23	2.02	0.40
1:B:265:SER:HA	1:B:300:ILE:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/494 (90%)	434 (98%)	8 (2%)	0	100	100
1	B	442/494 (90%)	434 (98%)	8 (2%)	0	100	100
All	All	884/988 (90%)	868 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/403 (85%)	336 (98%)	7 (2%)	48	68
1	B	345/403 (86%)	335 (97%)	10 (3%)	37	57
All	All	688/806 (85%)	671 (98%)	17 (2%)	42	63

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	LEU
1	A	292	THR
1	A	374	ARG
1	A	386	THR
1	A	393	PHE
1	A	437	GLU
1	A	467	ARG
1	B	117	LEU
1	B	258	LYS
1	B	300	ILE
1	B	348	GLU
1	B	359	ILE
1	B	368	LYS
1	B	383	ASP
1	B	415	THR
1	B	425	THR
1	B	433	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	GLN
1	A	133	GLN
1	A	469	ASN
1	B	10	GLN
1	B	67	GLN
1	B	439	GLN
1	B	469	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	B	501	-	25,25,48	3.26	13 (52%)	37,38,73	2.52	10 (27%)
2	NAD	A	501	-	25,25,48	3.37	13 (52%)	37,38,73	2.51	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	501	-	-	5/10/26/62	0/3/3/5
2	NAD	A	501	-	-	7/10/26/62	0/3/3/5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NAD	C3B-C4B	-9.29	1.29	1.53
2	B	501	NAD	C3B-C4B	-9.09	1.29	1.53
2	B	501	NAD	O4B-C4B	7.57	1.61	1.45
2	A	501	NAD	O4B-C4B	7.52	1.61	1.45
2	B	501	NAD	C6A-N6A	6.00	1.49	1.34
2	A	501	NAD	C6A-N6A	5.97	1.49	1.34
2	A	501	NAD	PA-O1A	4.73	1.65	1.50
2	B	501	NAD	O4B-C1B	-4.22	1.32	1.42
2	A	501	NAD	O4B-C1B	-4.15	1.32	1.42
2	B	501	NAD	O3B-C3B	3.15	1.50	1.43
2	A	501	NAD	O3B-C3B	3.12	1.50	1.43
2	A	501	NAD	C2A-N1A	2.90	1.39	1.33
2	B	501	NAD	C2A-N1A	2.89	1.39	1.33
2	B	501	NAD	PA-O3	2.83	1.65	1.54
2	A	501	NAD	C5A-C4A	-2.79	1.34	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAD	C5A-C4A	-2.76	1.34	1.39
2	A	501	NAD	PA-O5B	2.45	1.68	1.60
2	B	501	NAD	PA-O5B	2.44	1.68	1.60
2	B	501	NAD	O2B-C2B	-2.44	1.36	1.43
2	B	501	NAD	C8A-N9A	-2.42	1.33	1.37
2	A	501	NAD	O2B-C2B	-2.41	1.37	1.43
2	A	501	NAD	C8A-N9A	-2.41	1.33	1.37
2	A	501	NAD	C8A-N7A	2.19	1.35	1.31
2	B	501	NAD	C5A-N7A	-2.14	1.35	1.39
2	B	501	NAD	C8A-N7A	2.11	1.35	1.31
2	A	501	NAD	C5A-N7A	-2.04	1.35	1.39

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAD	C4A-N9A-C1B	-6.86	110.58	126.63
2	B	501	NAD	C4A-N9A-C1B	-6.59	111.23	126.63
2	A	501	NAD	N3A-C2A-N1A	-5.63	120.06	128.58
2	B	501	NAD	N3A-C2A-N1A	-5.53	120.21	128.58
2	A	501	NAD	C1B-N9A-C8A	5.42	139.11	127.09
2	A	501	NAD	N9A-C8A-N7A	-5.26	106.48	113.94
2	B	501	NAD	N9A-C8A-N7A	-5.25	106.49	113.94
2	B	501	NAD	C1B-N9A-C8A	5.19	138.61	127.09
2	B	501	NAD	C5A-C4A-N3A	-4.73	120.20	126.72
2	A	501	NAD	C5A-C4A-N3A	-4.49	120.53	126.72
2	A	501	NAD	C4A-N9A-C8A	4.35	110.30	105.74
2	B	501	NAD	C4A-N9A-C8A	4.21	110.16	105.74
2	B	501	NAD	N3A-C4A-N9A	3.73	133.50	127.17
2	A	501	NAD	N3A-C4A-N9A	3.57	133.24	127.17
2	B	501	NAD	C2A-N3A-C4A	3.18	119.60	111.83
2	A	501	NAD	C2A-N3A-C4A	3.15	119.53	111.83
2	B	501	NAD	C5A-N7A-C8A	3.13	108.37	103.45
2	A	501	NAD	C5A-N7A-C8A	3.00	108.16	103.45
2	B	501	NAD	C6A-C5A-C4A	2.07	120.00	117.18

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NAD	C5B-O5B-PA-O1A
2	A	501	NAD	C5B-O5B-PA-O2A
2	A	501	NAD	C5B-O5B-PA-O3

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	501	NAD	C5B-O5B-PA-O1A
2	B	501	NAD	C5B-O5B-PA-O2A
2	B	501	NAD	C5B-O5B-PA-O3
2	A	501	NAD	C2B-C1B-N9A-C8A
2	B	501	NAD	C2B-C1B-N9A-C8A
2	A	501	NAD	C2B-C1B-N9A-C4A
2	A	501	NAD	O4B-C1B-N9A-C8A
2	A	501	NAD	O4B-C4B-C5B-O5B
2	B	501	NAD	O4B-C1B-N9A-C8A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/494 (91%)	0.26	21 (4%) 36 30	26, 50, 87, 112	0
1	B	450/494 (91%)	0.08	17 (3%) 44 38	25, 43, 85, 115	0
All	All	900/988 (91%)	0.17	38 (4%) 40 34	25, 46, 86, 115	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	136	GLY	4.0
1	A	412	ALA	3.7
1	A	413	ASN	3.6
1	A	408	VAL	3.6
1	A	410	GLU	3.6
1	A	231	THR	3.6
1	A	299	GLU	3.0
1	A	403	ASP	3.0
1	B	423	VAL	2.9
1	B	137	GLY	2.9
1	B	415	THR	2.9
1	B	413	ASN	2.9
1	B	428	ILE	2.8
1	B	429	GLN	2.7
1	B	434	VAL	2.7
1	B	24	ASP	2.7
1	A	409	LEU	2.6
1	A	363	GLU	2.6
1	A	437	GLU	2.6
1	B	289	CYS	2.6
1	B	264	PHE	2.6
1	B	487	MET	2.5
1	B	133	GLN	2.5
1	B	300	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	67	GLN	2.3
1	A	351	ALA	2.3
1	A	290	ASN	2.3
1	B	431	ALA	2.3
1	A	487	MET	2.2
1	A	381	CYS	2.2
1	A	415	THR	2.2
1	B	136	GLY	2.2
1	A	383	ASP	2.2
1	A	24	ASP	2.1
1	A	402	PHE	2.1
1	B	138	SER	2.0
1	A	230	PHE	2.0
1	A	428	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAD	A	501	23/44	0.82	0.12	37,54,82,90	0
2	NAD	B	501	23/44	0.84	0.10	45,53,77,84	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAD A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.