



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 08:54 PM EDT

PDB ID : 6YA2
Title : Crystal structure of TSWV glycoprotein N ectodomain (Trypsin treated)
Authors : Dessau, M.; Bahat, Y.
Deposited on : 2020-03-11
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	 10% 73% 21% • 6%
1	B	199	 9% 81% 13% • 5%
1	C	199	 19% 69% 17% • 13%
2	D	2	 50% 50%
2	E	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	D	2	-	-	-	X
2	NAG	E	2	-	-	-	X
3	NAG	A	303	-	-	-	X
3	NAG	B	301	-	-	-	X
3	NAG	C	301	-	-	-	X
3	NAG	C	302	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	188	1454	917	238	290	9	0	0	0
1	B	190	1467	925	240	293	9	0	0	0
1	C	173	1326	837	219	261	9	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	214	CYS	SER	engineered mutation	UNP A0A3G1GK10
B	214	CYS	SER	engineered mutation	UNP A0A3G1GK10
C	214	CYS	SER	engineered mutation	UNP A0A3G1GK10

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	2	28	16	2	10	0	0	0
2	E	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0
3	B	1	14	8	1	5	0	0
3	C	1	14	8	1	5	0	0
3	C	1	14	8	1	5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	21	21	21	0	0
4	B	47	47	47	0	0
4	C	11	11	11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycoprotein



- Molecule 1: Glycoprotein



- Molecule 1: Glycoprotein



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  50% 50%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.66Å 76.01Å 71.18Å 90.00° 106.56° 90.00°	Depositor
Resolution (Å)	18.85 – 2.50 18.99 – 2.50	Depositor EDS
% Data completeness (in resolution range)	82.7 (18.85-2.50) 82.7 (18.99-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.49Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.213 , 0.244 0.213 , 0.244	Depositor DCC
R_{free} test set	1026 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	50.2	Xtrriage
Anisotropy	0.039	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 62.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.010 for l,-k,h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4438	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1482	0.49	0/2007
1	B	0.34	0/1495	0.52	0/2025
1	C	0.36	0/1349	0.50	0/1820
All	All	0.35	0/4326	0.50	0/5852

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1454	0	1439	23	0
1	B	1467	0	1453	14	0
1	C	1326	0	1331	20	0
2	D	28	0	25	0	0
2	E	28	0	25	2	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
3	C	28	0	26	3	0
4	A	21	0	0	0	0
4	B	47	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	11	0	0	0	0
All	All	4438	0	4325	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:TYR:HB3	1:A:272:PHE:HB2	1.67	0.76
1:A:185:VAL:HG11	1:A:220:LEU:HD13	1.74	0.70
1:C:250:ILE:HG21	1:C:276:MET:HG2	1.79	0.63
1:C:110:LEU:HD12	1:C:111:PRO:HD2	1.81	0.62
1:A:272:PHE:HE2	1:A:292:ILE:HD12	1.66	0.60
3:C:301:NAG:C8	3:C:301:NAG:H3	2.30	0.60
1:C:124:ILE:HB	1:C:158:ASN:HB2	1.83	0.60
1:B:111:PRO:HB3	1:B:208:PRO:HG3	1.82	0.60
1:C:226:ILE:HG22	1:C:239:ILE:HD11	1.84	0.59
1:C:278:ASP:HA	1:C:294:LYS:HG3	1.84	0.59
1:B:250:ILE:HG21	1:B:276:MET:HE3	1.85	0.58
1:B:114:CYS:SG	1:B:203:THR:OG1	2.63	0.57
1:C:222:GLY:O	1:C:287:ASP:HB2	2.06	0.56
1:A:250:ILE:HG21	1:A:276:MET:HE3	1.85	0.56
1:C:179:ASN:HB3	1:C:182:HIS:HB3	1.87	0.55
1:C:174:ILE:HD13	1:C:180:LYS:HG2	1.88	0.55
1:A:228:LYS:NZ	1:A:234:PRO:O	2.36	0.54
1:C:171:VAL:HG22	1:C:218:VAL:HG22	1.91	0.53
1:B:185:VAL:HG11	1:B:220:LEU:HD13	1.91	0.53
2:E:1:NAG:H62	2:E:2:NAG:O5	2.10	0.52
1:B:255:GLU:HB3	1:B:257:VAL:HG23	1.91	0.52
1:B:166:PHE:O	1:B:168:LYS:NZ	2.39	0.52
1:A:224:CYS:O	1:A:288:LYS:HG3	2.11	0.51
1:C:250:ILE:HG12	1:C:276:MET:HE3	1.93	0.51
1:B:158:ASN:HA	1:C:158:ASN:HA	1.93	0.51
1:C:145:CYS:SG	1:C:208:PRO:HA	2.51	0.51
1:A:110:LEU:HB2	1:A:111:PRO:HD3	1.93	0.50
1:A:239:ILE:HD13	1:A:270:ILE:HD11	1.93	0.50
1:C:273:THR:OG1	1:C:276:MET:SD	2.64	0.50
1:B:212:THR:OG1	1:B:213:VAL:N	2.45	0.49
1:A:225:LYS:HB2	1:A:240:THR:HG22	1.94	0.49
1:A:281:HIS:O	1:A:292:ILE:HG12	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:ASN:HA	4:B:411:HOH:O	2.14	0.48
1:A:228:LYS:HA	1:A:237:VAL:HG23	1.96	0.48
1:B:257:VAL:HG12	1:B:260:LYS:HB3	1.95	0.48
1:A:205:ASN:O	1:A:206:SER:OG	2.29	0.48
1:A:209:THR:HG22	1:A:210:ASN:N	2.30	0.47
1:A:274:GLU:H	1:A:274:GLU:CD	2.18	0.46
1:C:276:MET:HA	1:C:281:HIS:CE1	2.50	0.46
1:C:116:ASN:OD1	1:C:119:SER:OG	2.34	0.46
1:A:154:GLU:H	1:A:154:GLU:CD	2.20	0.45
1:B:124:ILE:HG21	1:B:127:ILE:HD11	1.99	0.45
2:E:1:NAG:H4	2:E:2:NAG:H2	1.63	0.45
3:C:301:NAG:H3	3:C:301:NAG:H83	1.97	0.45
1:A:153:LEU:HB2	1:A:162:LEU:HD21	1.98	0.45
1:A:156:CYS:O	1:A:159:SER:HB2	2.16	0.44
1:A:122:CYS:SG	1:A:123:GLU:N	2.89	0.44
1:A:109:ASP:HB3	1:A:112:ASN:HB2	2.00	0.44
1:B:292:ILE:HA	1:B:293:PRO:HD3	1.87	0.44
1:A:248:TYR:CE1	1:A:262:ILE:HB	2.53	0.43
1:B:157:ASP:OD1	1:B:158:ASN:N	2.51	0.43
1:C:257:VAL:CG1	1:C:260:LYS:HB3	2.48	0.43
3:C:301:NAG:C8	3:C:301:NAG:C3	2.95	0.43
1:A:225:LYS:HB2	1:A:240:THR:CG2	2.48	0.42
1:C:130:TYR:HB2	1:C:188:LYS:HG3	2.00	0.42
1:C:225:LYS:HB2	1:C:240:THR:OG1	2.19	0.42
1:C:124:ILE:HD13	1:C:134:TYR:HB3	2.01	0.42
1:B:229:SER:HB2	1:B:231:PHE:CE2	2.55	0.41
1:A:174:ILE:HG12	1:A:219:LYS:HB3	2.03	0.41
1:A:283:LEU:HB2	1:A:292:ILE:HD11	2.03	0.40
1:C:133:TYR:CE1	1:C:213:VAL:HG21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/199 (94%)	182 (98%)	4 (2%)	0	100	100
1	B	188/199 (94%)	184 (98%)	4 (2%)	0	100	100
1	C	169/199 (85%)	161 (95%)	8 (5%)	0	100	100
All	All	543/597 (91%)	527 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/183 (94%)	167 (97%)	5 (3%)	42	69
1	B	174/183 (95%)	166 (95%)	8 (5%)	27	50
1	C	157/183 (86%)	153 (98%)	4 (2%)	47	73
All	All	503/549 (92%)	486 (97%)	17 (3%)	37	63

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198	ASP
1	A	214	CYS
1	A	218	VAL
1	A	220	LEU
1	A	231	PHE
1	B	116	ASN
1	B	124	ILE
1	B	165	ARG
1	B	220	LEU
1	B	259	HIS
1	B	260	LYS
1	B	261	VAL
1	B	296	ASN
1	C	174	ILE
1	C	212	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	238	SER
1	C	239	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	1	1,2	14,14,15	0.60	0	17,19,21	0.50	0
2	NAG	D	2	2	14,14,15	0.54	0	17,19,21	0.92	1 (5%)
2	NAG	E	1	1,2	14,14,15	0.32	0	17,19,21	0.78	1 (5%)
2	NAG	E	2	2	14,14,15	1.21	1 (7%)	17,19,21	0.97	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	2	2	-	3/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	NAG	C1-C2	4.06	1.58	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	NAG	C2-N2-C7	2.18	126.00	122.90
2	E	1	NAG	O4-C4-C3	-2.17	105.33	110.35
2	E	2	NAG	C4-C3-C2	2.14	114.15	111.02
2	E	2	NAG	O5-C5-C4	-2.02	105.92	110.83

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	NAG	C4-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C1-C2-N2-C7
2	D	2	NAG	O5-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
2	E	1	NAG	C3-C2-N2-C7
2	D	2	NAG	C4-C5-C6-O6
2	D	1	NAG	C3-C2-N2-C7


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	NAG	2	0
2	E	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	301	1	14,14,15	0.67	0	17,19,21	0.81	1 (5%)
3	NAG	C	302	1	14,14,15	0.75	1 (7%)	17,19,21	0.63	0
3	NAG	A	303	1	14,14,15	1.30	2 (14%)	17,19,21	1.03	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	301	1	14,14,15	0.59	0	17,19,21	1.99	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	301	1	-	1/6/23/26	0/1/1/1
3	NAG	C	302	1	-	2/6/23/26	0/1/1/1
3	NAG	A	303	1	-	1/6/23/26	0/1/1/1
3	NAG	C	301	1	-	5/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	NAG	C1-C2	3.59	1.57	1.52
3	C	302	NAG	C1-C2	2.49	1.56	1.52
3	A	303	NAG	O5-C1	2.48	1.47	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	NAG	C2-N2-C7	7.60	133.73	122.90
3	A	303	NAG	C1-O5-C5	2.80	115.99	112.19
3	B	301	NAG	C3-C4-C5	2.10	113.99	110.24
3	A	303	NAG	O5-C5-C4	-2.02	105.92	110.83

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	301	NAG	C3-C2-N2-C7
3	C	302	NAG	C4-C5-C6-O6
3	C	301	NAG	C8-C7-N2-C2
3	C	301	NAG	O7-C7-N2-C2
3	C	301	NAG	O5-C5-C6-O6
3	C	302	NAG	O5-C5-C6-O6
3	C	301	NAG	C4-C5-C6-O6
3	A	303	NAG	O5-C5-C6-O6
3	B	301	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	301	NAG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	188/199 (94%)	0.66	19 (10%) 7 6	31, 64, 134, 172	0
1	B	190/199 (95%)	0.48	17 (8%) 9 9	26, 52, 119, 183	0
1	C	173/199 (86%)	1.08	37 (21%) 0 0	38, 82, 145, 174	0
All	All	551/597 (92%)	0.73	73 (13%) 3 3	26, 66, 135, 183	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	234	PRO	9.2
1	A	231	PHE	8.6
1	B	110	LEU	7.9
1	B	296	ASN	7.2
1	C	232	ALA	6.8
1	A	296	ASN	6.7
1	B	109	ASP	6.4
1	A	232	ALA	6.3
1	A	110	LEU	5.9
1	C	175	THR	5.9
1	A	295	THR	5.6
1	C	256	ASN	5.3
1	A	230	ASN	5.2
1	C	288	LYS	4.7
1	C	258	GLU	4.6
1	A	294	LYS	4.6
1	A	109	ASP	4.5
1	C	235	TYR	4.5
1	B	107	VAL	4.4
1	A	259	HIS	4.2
1	B	108	SER	4.0
1	C	110	LEU	3.9
1	C	220	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	ASN	3.7
1	C	274	GLU	3.6
1	C	259	HIS	3.6
1	A	235	TYR	3.6
1	C	257	VAL	3.6
1	B	257	VAL	3.6
1	B	232	ALA	3.4
1	C	285	CYS	3.2
1	C	225	LYS	3.2
1	C	233	ASN	3.1
1	B	112	ASN	3.0
1	C	245	ILE	3.0
1	B	115	LEU	3.0
1	A	233	ASN	3.0
1	A	111	PRO	2.9
1	A	257	VAL	2.9
1	C	255	GLU	2.8
1	B	111	PRO	2.8
1	B	295	THR	2.7
1	A	112	ASN	2.7
1	C	210	ASN	2.7
1	B	134	TYR	2.6
1	C	242	PRO	2.6
1	C	252	LYS	2.6
1	B	197	GLN	2.6
1	B	259	HIS	2.6
1	A	274	GLU	2.6
1	C	164	LYS	2.5
1	B	231	PHE	2.5
1	C	265	SER	2.4
1	C	230	ASN	2.4
1	C	273	THR	2.4
1	C	177	LEU	2.4
1	A	140	GLY	2.3
1	C	275	GLU	2.3
1	C	294	LYS	2.3
1	C	221	SER	2.3
1	C	228	LYS	2.3
1	B	145	CYS	2.3
1	A	113	ASN	2.2
1	C	254	GLY	2.2
1	C	116	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	145	CYS	2.2
1	C	176	LYS	2.1
1	C	295	THR	2.1
1	C	111	PRO	2.0
1	A	139	ASN	2.0
1	B	230	ASN	2.0
1	C	286	GLY	2.0
1	C	293	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	2	14/15	0.57	0.61	131,139,146,146	0
2	NAG	D	1	14/15	0.65	0.38	120,129,139,146	0
2	NAG	E	1	14/15	0.70	0.35	108,117,122,127	0
2	NAG	E	2	14/15	0.76	0.59	130,132,136,139	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	303	14/15	0.18	1.14	183,194,203,203	0
3	NAG	C	301	14/15	0.54	0.69	105,126,137,138	0
3	NAG	C	302	14/15	0.55	0.66	132,160,172,177	0
3	NAG	B	301	14/15	0.56	0.51	144,148,154,154	0

6.5 Other polymers [i](#)

There are no such residues in this entry.