



Full wwPDB X-ray Structure Validation Report i

Jun 24, 2024 – 10:56 AM EDT

PDB ID : 6YCZ
Title : Plasmodium falciparum Myosin A delta-Nter, Post-Rigor state
Authors : Moussaoui, D.; Robblee, J.P.; Auguin, D.; Krementsova, E.B.; Robert-Paganin, J.; Trybus, K.M.; Houdusse, A.
Deposited on : 2020-03-19
Resolution : 3.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1177 (3.32-3.24)
Clashscore	141614	1044 (3.30-3.26)
Ramachandran outliers	138981	1026 (3.30-3.26)
Sidechain outliers	138945	1025 (3.30-3.26)
RSRZ outliers	127900	1141 (3.32-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 8 unique types of molecules in this entry. The entry contains 8556 atoms, of which 12 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	795	Total	C 6361	N 4053	O 1080	S 1195	33	0	6	0


- Molecule 2 is a protein called Myosin A tail domain interacting protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	B	131	Total	C 1051	N 661	O 167	S 218	5	0	0	0

- Molecule 3 is a protein called Uncharacterized protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	C	131	Total	C 1077	N 687	O 175	S 209	6	0	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H O 10 2 6 2	0	0
5	C	1	Total C H O 10 2 6 2	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mg 1 1	0	0

- Molecule 7 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
7	A	1	27	10	5	10	2	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	4	Total 4 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Myosin-A



- Molecule 2: Myosin A tail domain interacting protein



- Molecule 3: Uncharacterized protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.08 Å 114.43 Å 170.71 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.30 – 3.27 48.30 – 3.27	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.30-3.27) 100.0 (48.30-3.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.72 (at 3.25 Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R , R_{free}	0.179 , 0.227 0.193 , 0.246	Depositor DCC
R_{free} test set	1399 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	84.4	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 75.9	EDS
L-test for twinning ²	$< L > = 0.46$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8556	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ADP, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/6474	0.72	0/8729
2	B	0.46	0/1068	0.65	0/1440
3	C	0.51	0/1096	0.71	0/1476
All	All	0.51	0/8638	0.71	0/11645

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6361	0	6457	53	0
2	B	1051	0	1005	8	0
3	C	1077	0	1057	12	0
4	A	10	0	0	0	0
4	B	5	0	0	0	0
5	A	4	6	6	0	0
5	C	4	6	6	0	0
6	A	1	0	0	0	0
7	A	27	0	12	2	0
8	A	4	0	0	0	0
All	All	8544	12	8543	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:68:GLN:HE22	3:C:76:VAL:H	1.41	0.68
2:B:168:LEU:HB3	2:B:175:LEU:HD12	1.77	0.66
1:A:233:ALA:HB1	1:A:289[B]:ILE:HG23	1.79	0.64
1:A:766:LEU:HA	1:A:769:ILE:HD12	1.81	0.62
1:A:414:ALA:HB3	1:A:419:ILE:HD13	1.81	0.62
3:C:91:LEU:HD21	3:C:110:LEU:HD11	1.82	0.61
2:B:113:LYS:HD2	2:B:133:ILE:HG21	1.84	0.60
1:A:798:ASN:HA	1:A:801:ILE:HD12	1.85	0.59
1:A:197:LYS:HE3	7:A:905:ADP:O2B	2.04	0.58
1:A:785:GLU:HA	3:C:101:LEU:HD11	1.86	0.57
1:A:159:ASP:O	1:A:162:LYS:HG2	2.05	0.57
1:A:729:VAL:HG13	1:A:734:LYS:HG3	1.87	0.57
1:A:718:PHE:H	1:A:770:GLN:HE21	1.54	0.56
1:A:61[A]:SER:OG	1:A:66:LEU:HA	2.06	0.55
1:A:212:LYS:HD2	1:A:259:GLU:HG2	1.90	0.54
1:A:238:LYS:HB3	1:A:285:ARG:HB2	1.90	0.53
1:A:488:ILE:HD13	1:A:539:LEU:HD11	1.91	0.53
1:A:221:GLN:HG3	1:A:262:ILE:HB	1.91	0.52
1:A:54:LYS:HD2	1:A:90:CYS:SG	2.50	0.51
3:C:70:PHE:HD2	3:C:131:THR:HG23	1.74	0.51
1:A:39[B]:ASP:OD1	1:A:86:HIS:HD2	1.94	0.50
1:A:41:SER:HB3	1:A:44:ILE:HD12	1.94	0.50
1:A:397:TYR:HB2	1:A:621:VAL:HG21	1.93	0.50
1:A:289[B]:ILE:HD11	1:A:440:TYR:OH	2.13	0.49
1:A:358:GLY:HA3	1:A:396:MET:HG3	1.94	0.48
1:A:582:ILE:HD12	1:A:591:GLN:HB2	1.95	0.48
2:B:167:ILE:HG13	3:C:19:CYS:SG	2.54	0.48
1:A:603:ASP:O	1:A:641:ILE:HD11	2.14	0.48
1:A:32:LYS:HD3	3:C:45:ALA:HA	1.96	0.47
1:A:166:HIS:O	1:A:169:THR:HB	2.14	0.47
1:A:693:LEU:HA	1:A:696:LEU:HD12	1.95	0.47
1:A:582:ILE:HB	1:A:591:GLN:HG3	1.97	0.47
1:A:197:LYS:HB3	1:A:469:ASP:OD1	2.16	0.45
1:A:185:GLN:HG3	1:A:662:HIS:CE1	2.51	0.45
1:A:424:ASN:ND2	1:A:424:ASN:H	2.15	0.45
1:A:185:GLN:HE21	1:A:662:HIS:HE1	1.64	0.45
3:C:14:ILE:HB	3:C:19:CYS:HA	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:106:PRO:HB3	2:B:134:CYS:SG	2.57	0.44
3:C:70:PHE:HB3	3:C:73:ILE:HD12	1.99	0.44
1:A:396:MET:HB3	1:A:398:LEU:HD12	1.99	0.44
1:A:398:LEU:HD21	1:A:613:ILE:HG12	2.00	0.44
2:B:169:THR:HG21	2:B:177:ASP:HA	1.99	0.44
1:A:184:SER:HB3	1:A:659:THR:HG22	1.98	0.44
1:A:302:LYS:HA	1:A:307:LEU:HD12	2.00	0.44
1:A:635:ILE:HD13	1:A:640:LEU:HD21	2.00	0.43
1:A:512:GLU:HG3	1:A:753:LYS:H	1.84	0.43
1:A:253:GLN:O	1:A:265:GLY:HA2	2.19	0.43
1:A:39[B]:ASP:OD1	1:A:86:HIS:CD2	2.72	0.43
1:A:196:GLY:HA2	7:A:905:ADP:O2A	2.19	0.43
2:B:195:TYR:HA	2:B:198:PHE:HB3	2.01	0.42
3:C:122:ASN:OD1	3:C:124:LYS:HB2	2.18	0.42
1:A:99:PHE:CE1	1:A:104:LEU:HB2	2.54	0.42
1:A:208:PHE:HB3	1:A:256:ILE:HG12	2.01	0.42
2:B:84:LYS:HB2	2:B:90:ILE:HD11	2.01	0.42
1:A:292:GLN:HA	1:A:328:ASP:HB3	2.01	0.42
1:A:99:PHE:CD1	1:A:104:LEU:HB2	2.54	0.42
1:A:802:PRO:HA	1:A:805:LEU:HD12	2.01	0.42
2:B:132:SER:O	2:B:135:VAL:HG12	2.20	0.42
1:A:682:LYS:HA	1:A:685:ILE:HD12	2.01	0.42
1:A:135:VAL:HG22	1:A:692:ILE:HG12	2.02	0.41
1:A:198:THR:O	1:A:202:LYS:HG3	2.20	0.41
1:A:224:ILE:O	1:A:227:ALA:HB3	2.21	0.41
1:A:442:LYS:HG3	1:A:625:PHE:CE2	2.55	0.41
1:A:587:ILE:HD11	1:A:590:ILE:HD11	2.03	0.41
3:C:131:THR:HA	3:C:134:ILE:HD12	2.03	0.40
1:A:582:ILE:HD11	1:A:589:PRO:HB2	2.02	0.40
3:C:86:LYS:HB2	3:C:89:ASN:HB2	2.03	0.40
1:A:233:ALA:O	1:A:289[B]:ILE:HG13	2.22	0.40
1:A:480:SER:H	1:A:483:GLN:HG2	1.86	0.40
3:C:113:THR:HA	3:C:129:LYS:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	797/818 (97%)	741 (93%)	49 (6%)	7 (1%)	17 50
2	B	129/204 (63%)	122 (95%)	7 (5%)	0	100 100
3	C	127/134 (95%)	118 (93%)	6 (5%)	3 (2%)	6 29
All	All	1053/1156 (91%)	981 (93%)	62 (6%)	10 (1%)	15 50

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	607	GLY
1	A	321	THR
1	A	635	ILE
1	A	817	ALA
3	C	83	VAL
3	C	119	SER
1	A	21	VAL
3	C	113	THR
1	A	34	TYR
1	A	550	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	713/726 (98%)	657 (92%)	56 (8%)	12 37
2	B	117/186 (63%)	110 (94%)	7 (6%)	19 49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	C	123/126 (98%)	109 (89%)	14 (11%)	5 23
All	All	953/1038 (92%)	876 (92%)	77 (8%)	11 36

All (77) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LYS
1	A	54	LYS
1	A	56	VAL
1	A	83	ASP
1	A	85	THR
1	A	91	ASN
1	A	157	THR
1	A	173	GLU
1	A	222	THR
1	A	228	ASN
1	A	241	ARG
1	A	282	ASP
1	A	304	LYS
1	A	347	SER
1	A	371	GLN
1	A	377	ASP
1	A	412	THR
1	A	417	THR
1	A	418	LYS
1	A	429	GLU
1	A	458	GLU
1	A	471	PHE
1	A	480	SER
1	A	483	GLN
1	A	508	LEU
1	A	512	GLU
1	A	519	LEU
1	A	522	THR
1	A	525	LYS
1	A	530	VAL
1	A	566	GLU
1	A	574	LYS
1	A	583	ILE
1	A	586	THR
1	A	597	PHE
1	A	598	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	606	ARG
1	A	615	ASP
1	A	629	VAL
1	A	630	ILE
1	A	635	ILE
1	A	644	GLN
1	A	646	LEU
1	A	680	GLU
1	A	699	ARG
1	A	706	ARG
1	A	728	SER
1	A	746	GLU
1	A	747	SER
1	A	754	SER
1	A	774	LEU
1	A	801	ILE
1	A	803	SER
1	A	805	LEU
1	A	806	ARG
1	A	814	LYS
2	B	76	ASP
2	B	93	ASP
2	B	133	ILE
2	B	135	VAL
2	B	138	LYS
2	B	153	ASN
2	B	155	CYS
3	C	6	GLU
3	C	18	SER
3	C	27	LYS
3	C	52	ASN
3	C	58	ASN
3	C	74	ASN
3	C	88	ASP
3	C	89	ASN
3	C	107	ASN
3	C	108	ILE
3	C	109	LEU
3	C	119	SER
3	C	125	THR
3	C	134	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	86	HIS
1	A	182	ASN
1	A	185	GLN
1	A	283	ASN
1	A	424	ASN
1	A	585	HIS
1	A	647	ASN
1	A	672	ASN
1	A	686	GLN
1	A	737	ASN
1	A	770	GLN
1	A	810	HIS
2	B	94	ASN
3	C	68	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	901	-	4,4,4	0.26	0	6,6,6	0.30	0
4	SO4	B	301	-	4,4,4	0.36	0	6,6,6	0.22	0
4	SO4	A	902	-	4,4,4	0.49	0	6,6,6	0.45	0
5	EDO	C	201	-	3,3,3	0.48	0	2,2,2	0.52	0
7	ADP	A	905	6	24,29,29	0.89	1 (4%)	29,45,45	0.84	1 (3%)
5	EDO	A	903	-	3,3,3	0.75	0	2,2,2	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	201	-	-	0/1/1/1	-
7	ADP	A	905	6	-	0/12/32/32	0/3/3/3
5	EDO	A	903	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	905	ADP	PB-O3B	-2.63	1.45	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	905	ADP	C5-C6-N6	2.11	123.53	120.31

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	905	ADP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	795/818 (97%)	0.29	54 (6%) 17 17	49, 82, 143, 174	0
2	B	131/204 (64%)	0.48	13 (9%) 7 7	69, 106, 154, 164	0
3	C	131/134 (97%)	0.34	4 (3%) 49 48	60, 93, 145, 174	0
All	All	1057/1156 (91%)	0.32	71 (6%) 17 17	49, 86, 145, 174	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	575	VAL	5.9
1	A	577	SER	5.6
1	A	541	TYR	5.2
1	A	752	GLY	5.0
1	A	576	ALA	4.6
1	A	751	ILE	4.4
3	C	117	ASN	3.6
1	A	735	CYS	3.6
1	A	566	GLU	3.6
1	A	371	GLN	3.5
1	A	376	SER	3.5
1	A	757	PHE	3.4
1	A	519	LEU	3.4
1	A	545	GLN	3.3
3	C	119	SER	3.3
1	A	573	ALA	3.3
2	B	104	LEU	3.2
2	B	134	CYS	3.2
2	B	103	GLY	3.1
1	A	568	ASN	3.1
1	A	516	THR	3.0
1	A	756	VAL	3.0
1	A	554	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	477	LYS	2.9
2	B	98	ASN	2.9
1	A	578	ASN	2.9
1	A	581	PHE	2.8
2	B	102	LEU	2.8
2	B	74	GLU	2.8
2	B	101	LYS	2.7
1	A	530	VAL	2.7
1	A	544	ASP	2.6
1	A	561	ALA	2.6
1	A	572	PRO	2.6
2	B	88	GLY	2.6
1	A	720	ASP	2.6
1	A	560	CYS	2.5
1	A	534	LYS	2.5
1	A	738	ILE	2.5
1	A	574	LYS	2.5
1	A	567	ASN	2.5
1	A	755	MET	2.5
1	A	731	ASN	2.4
1	A	724	ALA	2.4
2	B	135	VAL	2.4
2	B	75	SER	2.4
1	A	548	ALA	2.4
1	A	514	ILE	2.4
3	C	121	LEU	2.4
1	A	645	PHE	2.4
1	A	549	PRO	2.4
1	A	540	SER	2.3
1	A	511	ASP	2.3
1	A	758	LEU	2.3
1	A	580	ASN	2.3
3	C	120	THR	2.3
1	A	590	ILE	2.2
1	A	557	VAL	2.2
1	A	385	ASP	2.2
1	A	766	LEU	2.2
1	A	591	GLN	2.2
2	B	84	LYS	2.1
1	A	816	VAL	2.1
1	A	556	PHE	2.1
1	A	571	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	562	THR	2.1
1	A	538	VAL	2.1
1	A	749	TYR	2.1
2	B	80	TYR	2.0
1	A	535	GLY	2.0
2	B	97	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	903	4/4	0.78	0.38	85,87,91,91	0
5	EDO	C	201	4/4	0.88	0.13	90,91,93,93	0
4	SO4	B	301	5/5	0.94	0.24	103,104,105,105	0
4	SO4	A	902	5/5	0.96	0.18	88,89,91,94	0
4	SO4	A	901	5/5	0.96	0.40	96,97,98,98	0
7	ADP	A	905	27/27	0.97	0.24	58,64,69,71	0
6	MG	A	904	1/1	0.98	0.24	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.