



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2024 – 04:22 pm GMT

PDB ID : 6YQB  
Title : Taka-amylase in complex with alpha-glucosyl epi-cyclophellitol cyclosulfate inhibitor  
Authors : Armstrong, Z.; Chen, Y.; Artola, M.; Overkleeft, H.; Davies, G.  
Deposited on : 2020-04-16  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	499	 91% 5%
1	BBB	499	 92% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	93Z	AAA	506	-	-	X	-

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 15507 atoms, of which 7150 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	AAA	476	Total 7272	C 2365	H 3531	N 605	O 753	S 18	217	8	0
1	BBB	476	Total 7234	C 2355	H 3511	N 600	O 750	S 18	216	6	0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	AAA	1	Total 10	C 2	H 6	O 2	1	0
2	AAA	1	Total 10	C 2	H 6	O 2	1	0
2	BBB	1	Total 10	C 2	H 6	O 2	1	0
2	BBB	1	Total 10	C 2	H 6	O 2	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
2	BBB	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	AAA	1	Total	C	H	N	O	3	0
			28	8	14	1	5		
3	BBB	1	Total	C	H	N	O	3	0
			28	8	14	1	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	Ca	0	0
			1	1		
4	BBB	1	Total	Ca	0	0
			1	1		

- Molecule 5 is alpha-D-glucopyranose (three-letter code: GLC) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	AAA	1	Total	C	H	O	4	0
			22	6	11	5		
5	BBB	1	Total	C	H	O	4	0
			22	6	11	5		

- Molecule 6 is [(1 {S},2 {R},3 {R},4 {S},5 {R})-2-(hydroxymethyl)-3,4,5,6-tetrakis(oxidanyl)cyclohexyl] hydrogen sulfate (three-letter code: 93Z) (formula: C<sub>7</sub>H<sub>14</sub>O<sub>9</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	S		
6	AAA	1	Total	C	H	O	S	3	0
			27	7	11	8	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	BBB	1	Total	C	H	O	S	3	0
			27	7	11	8	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	AAA	406	Total	O	0	0
			406	406		
7	BBB	379	Total	O	0	0
			379	379		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-amylase

Chain AAA:  91% 5%



- Molecule 1: Alpha-amylase

Chain BBB:  92% 5%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.58Å 102.94Å 75.62Å 90.00° 103.81° 90.00°	Depositor
Resolution (Å)	59.78 – 1.50 59.78 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.5 (59.78-1.50) 98.5 (59.78-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.157 , 0.186 0.157 , 0.186	Depositor DCC
$R_{free}$ test set	7597 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.3	Xtrriage
Anisotropy	0.927	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 38.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	15507	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, NAG, EDO, CA, 93Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.79	1/3841 (0.0%)	0.88	3/5244 (0.1%)
1	BBB	0.78	0/3823	0.88	4/5220 (0.1%)
All	All	0.78	1/7664 (0.0%)	0.88	7/10464 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	294	GLU	CD-OE1	-6.00	1.19	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	204	ARG	NE-CZ-NH2	-11.54	114.53	120.30
1	BBB	204	ARG	NE-CZ-NH1	7.09	123.84	120.30
1	AAA	206	ASP	CB-CG-OD2	5.89	123.60	118.30
1	AAA	252	TYR	CB-CG-CD2	5.85	124.51	121.00
1	BBB	204	ARG	CD-NE-CZ	5.55	131.37	123.60
1	BBB	206	ASP	CB-CG-OD2	5.55	123.30	118.30
1	AAA	204	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3741	3531	3519	19	0
1	BBB	3723	3511	3501	10	0
2	AAA	8	12	12	1	0
2	BBB	16	24	24	0	0
3	AAA	14	14	13	0	0
3	BBB	14	14	13	2	0
4	AAA	1	0	0	0	0
4	BBB	1	0	0	0	0
5	AAA	11	11	10	0	0
5	BBB	11	11	10	0	0
6	AAA	16	11	0	7	0
6	BBB	16	11	0	0	0
7	AAA	406	0	0	5	0
7	BBB	379	0	0	5	0
All	All	8357	7150	7102	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:207[B]:THR:HG22	6:AAA:506:93Z:O14	1.44	1.14
1:AAA:207[B]:THR:CG2	6:AAA:506:93Z:O14	2.02	1.08
1:AAA:207[B]:THR:HG23	6:AAA:506:93Z:O7	1.68	0.92
1:AAA:383:LYS:HE3	7:AAA:834:HOH:O	1.72	0.88
3:BBB:501:NAG:H81	7:BBB:852:HOH:O	1.76	0.85
1:AAA:186[A]:GLU:OE1	7:AAA:601:HOH:O	1.99	0.81
1:BBB:230[B]:GLU:OE1	7:BBB:601:HOH:O	2.02	0.77
1:AAA:207[B]:THR:HG23	6:AAA:506:93Z:O14	1.88	0.73
1:AAA:207[B]:THR:CG2	6:AAA:506:93Z:S13	2.76	0.73
1:AAA:207[B]:THR:HG23	6:AAA:506:93Z:S13	2.30	0.71
1:AAA:207[B]:THR:CG2	6:AAA:506:93Z:O7	2.44	0.62
1:AAA:383:LYS:CE	7:AAA:834:HOH:O	2.40	0.54
1:BBB:319:ILE:HD13	1:BBB:325:PRO:HB3	1.92	0.52
1:AAA:467:GLU:HG2	7:AAA:743:HOH:O	2.11	0.50
1:BBB:37:TYR:CE1	1:BBB:79:TYR:HA	2.47	0.50
3:BBB:501:NAG:H82	7:BBB:793:HOH:O	2.13	0.48
1:AAA:155:TYR:HD2	2:AAA:502:EDO:O1	1.97	0.48
1:AAA:397[B]:ARG:HH21	1:AAA:397[B]:ARG:CG	2.28	0.46
1:BBB:230[B]:GLU:OE2	1:BBB:250:LEU:HD12	2.17	0.45
1:BBB:361:LEU:HD13	1:BBB:361:LEU:C	2.37	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:176:LEU:HD13	1:BBB:187:TRP:HE1	1.82	0.44
1:AAA:172:SER:HA	7:AAA:622:HOH:O	2.19	0.42
1:AAA:63:THR:HB	1:AAA:64:PRO:CD	2.50	0.42
1:AAA:302:ALA:HA	1:AAA:305:THR:O	2.19	0.42
1:BBB:204:ARG:NH2	7:BBB:613:HOH:O	2.53	0.42
1:AAA:270:ASP:OD1	1:AAA:389:LYS:HD2	2.20	0.41
1:AAA:37:TYR:CE1	1:AAA:79:TYR:HA	2.55	0.41
1:AAA:397[A]:ARG:HD3	1:AAA:397[A]:ARG:C	2.41	0.41
1:BBB:467:GLU:HG2	7:BBB:645:HOH:O	2.20	0.41
1:BBB:408:ILE:N	1:BBB:408:ILE:HD12	2.36	0.41
1:BBB:240:CYS:N	1:BBB:241:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	482/499 (97%)	470 (98%)	12 (2%)	0	100	100
1	BBB	480/499 (96%)	470 (98%)	10 (2%)	0	100	100
All	All	962/998 (96%)	940 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	405/414 (98%)	397 (98%)	8 (2%)	55	25
1	BBB	403/414 (97%)	398 (99%)	5 (1%)	71	48
All	All	808/828 (98%)	795 (98%)	13 (2%)	67	36

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	61	TRP
1	AAA	186[A]	GLU
1	AAA	186[B]	GLU
1	AAA	220	ASN
1	AAA	293	VAL
1	AAA	397[A]	ARG
1	AAA	397[B]	ARG
1	AAA	436	GLU
1	BBB	61	TRP
1	BBB	187	TRP
1	BBB	220	ASN
1	BBB	221	LYS
1	BBB	264	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	93Z	AAA	506	5,1	16,16,17	1.56	2 (12%)	18,24,26	1.66	2 (11%)
3	NAG	BBB	501	1	14,14,15	1.33	3 (21%)	17,19,21	2.17	7 (41%)
5	GLC	AAA	505	6	11,11,12	0.69	0	15,15,17	1.08	1 (6%)
5	GLC	BBB	507	6	11,11,12	0.66	0	15,15,17	1.11	1 (6%)
2	EDO	BBB	505	-	3,3,3	0.28	0	2,2,2	0.60	0
2	EDO	AAA	502	-	3,3,3	0.14	0	2,2,2	0.72	0
3	NAG	AAA	503	1	14,14,15	1.01	1 (7%)	17,19,21	1.38	2 (11%)
2	EDO	AAA	501	-	3,3,3	0.09	0	2,2,2	0.24	0
2	EDO	BBB	504	-	3,3,3	0.11	0	2,2,2	0.73	0
2	EDO	BBB	503	-	3,3,3	0.38	0	2,2,2	0.09	0
6	93Z	BBB	508	5,1	16,16,17	1.15	2 (12%)	18,24,26	1.35	3 (16%)
2	EDO	BBB	502	-	3,3,3	0.39	0	2,2,2	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	93Z	AAA	506	5,1	-	4/7/27/31	0/1/1/1
3	NAG	BBB	501	1	-	2/6/23/26	0/1/1/1
5	GLC	AAA	505	6	-	0/2/19/22	0/1/1/1
5	GLC	BBB	507	6	-	0/2/19/22	0/1/1/1
2	EDO	BBB	505	-	-	0/1/1/1	-
2	EDO	AAA	502	-	-	0/1/1/1	-
3	NAG	AAA	503	1	-	0/6/23/26	0/1/1/1
2	EDO	AAA	501	-	-	0/1/1/1	-
2	EDO	BBB	504	-	-	1/1/1/1	-
2	EDO	BBB	503	-	-	0/1/1/1	-
6	93Z	BBB	508	5,1	-	2/7/27/31	0/1/1/1
2	EDO	BBB	502	-	-	0/1/1/1	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	AAA	506	93Z	C2-C3	3.79	1.58	1.52
6	AAA	506	93Z	C5-C4	-3.15	1.50	1.53
3	BBB	501	NAG	O4-C4	2.49	1.48	1.43
3	BBB	501	NAG	O7-C7	2.42	1.28	1.23
6	BBB	508	93Z	C8-C5	2.33	1.56	1.52
6	BBB	508	93Z	C2-C3	2.31	1.55	1.52
3	BBB	501	NAG	O5-C5	2.21	1.47	1.43
3	AAA	503	NAG	C1-C2	2.13	1.55	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	AAA	506	93Z	C6-O7-S13	5.08	124.53	117.91
3	BBB	501	NAG	C2-N2-C7	4.09	128.73	122.90
3	BBB	501	NAG	C6-C5-C4	-3.64	104.47	113.00
3	BBB	501	NAG	C3-C4-C5	-3.63	103.77	110.24
6	BBB	508	93Z	C6-O7-S13	3.47	122.44	117.91
3	AAA	503	NAG	O5-C1-C2	-3.18	106.27	111.29
6	AAA	506	93Z	C2-C1-C6	2.78	117.66	111.21
3	BBB	501	NAG	O4-C4-C5	2.69	115.98	109.30
3	BBB	501	NAG	C1-O5-C5	2.69	115.83	112.19
5	AAA	505	GLC	C2-C3-C4	2.46	115.15	110.89
6	BBB	508	93Z	C1-C6-C5	2.45	115.86	112.47
3	BBB	501	NAG	O6-C6-C5	-2.36	103.19	111.29
5	BBB	507	GLC	O5-C5-C6	-2.23	103.70	107.20
3	BBB	501	NAG	O4-C4-C3	2.12	115.26	110.35
3	AAA	503	NAG	O5-C5-C6	2.12	110.53	107.20
6	BBB	508	93Z	C2-C1-C6	2.07	116.01	111.21

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	AAA	506	93Z	C5-C6-O7-S13
6	AAA	506	93Z	C6-O7-S13-O16
6	BBB	508	93Z	C5-C6-O7-S13
3	BBB	501	NAG	C8-C7-N2-C2
3	BBB	501	NAG	O7-C7-N2-C2
6	AAA	506	93Z	C1-C6-O7-S13
6	AAA	506	93Z	C6-O7-S13-O14
2	BBB	504	EDO	O1-C1-C2-O2
6	BBB	508	93Z	C1-C6-O7-S13


There are no ring outliers.


3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	AAA	506	93Z	7	0
3	BBB	501	NAG	2	0
2	AAA	502	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	476/499 (95%)	-0.55	1 (0%) 95   95	14, 20, 30, 56	1 (0%)
1	BBB	476/499 (95%)	-0.49	0 100   100	15, 21, 35, 63	3 (0%)
All	All	952/998 (95%)	-0.52	1 (0%) 95   95	14, 20, 33, 63	4 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	447	SER	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	BBB	501	14/15	0.88	0.09	28,37,46,50	3
2	EDO	AAA	502	4/4	0.89	0.15	29,33,43,43	1
3	NAG	AAA	503	14/15	0.93	0.11	29,35,38,41	3
5	GLC	AAA	505	11/12	0.93	0.10	17,19,20,22	22


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	AAA	501	4/4	0.95	0.11	24,29,37,37	1
2	EDO	BBB	504	4/4	0.96	0.08	25,30,31,31	1
2	EDO	BBB	502	4/4	0.96	0.07	25,29,31,31	1
5	GLC	BBB	507	11/12	0.96	0.06	17,18,19,20	4
6	93Z	AAA	506	16/17	0.96	0.08	16,20,21,24	27
2	EDO	BBB	505	4/4	0.97	0.08	22,27,29,29	1
2	EDO	BBB	503	4/4	0.97	0.10	28,30,36,36	1
6	93Z	BBB	508	16/17	0.98	0.06	16,19,22,25	3
4	CA	BBB	506	1/1	0.99	0.08	17,17,17,17	0
4	CA	AAA	504	1/1	1.00	0.07	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.