



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 01:05 am GMT

PDB ID : 7AD0
Title : X-ray structure of Mdm2 with modified p53 peptide
Authors : Twarda-Clapa, A.; Fortuna, P.; Grudnik, P.; Dubin, G.; Berlicki, L.; Holak, T.A.
Deposited on : 2020-09-13
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	90	 4% 80% 17% .
1	B	90	 87% 10% .
1	C	90	 % 83% 11% . .
1	D	90	 % 87% 8% . .
1	E	90	 88% 7% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	90	<p>%</p> <p>84% 11% ..</p>
2	H	14	<p>86% 14%</p>
2	I	14	<p>7%</p> <p>93% 7%</p>
2	J	14	<p>93% 7%</p>
2	K	14	<p>93% 7%</p>
2	L	14	<p>7%</p> <p>100%</p>
2	M	14	<p>14%</p> <p>86% 7% 7%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5145 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase Mdm2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	90	Total 747	C 486	N 125	O 132	S 4	0	0	0
1	B	87	Total 725	C 475	N 119	O 126	S 5	0	1	0
1	C	86	Total 714	C 470	N 116	O 123	S 5	0	1	0
1	D	87	Total 725	C 475	N 119	O 126	S 5	0	1	0
1	E	87	Total 720	C 471	N 119	O 126	S 4	0	0	0
1	F	87	Total 725	C 475	N 119	O 126	S 5	0	1	0

- Molecule 2 is a protein called Modified p53 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	H	14	Total 104	C 72	N 14	O 18	0	0	1
2	I	14	Total 104	C 72	N 14	O 18	0	0	1
2	J	14	Total 104	C 72	N 14	O 18	0	0	1
2	K	14	Total 104	C 72	N 14	O 18	0	0	1
2	L	14	Total 104	C 72	N 14	O 18	0	0	1
2	M	14	Total 104	C 72	N 14	O 18	0	0	1


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	12	Total O 12 12	0	0
3	B	29	Total O 29 29	0	0
3	C	5	Total O 5 5	0	0
3	D	33	Total O 33 33	0	0
3	E	20	Total O 20 20	0	0
3	F	15	Total O 15 15	0	0
3	H	9	Total O 9 9	0	0
3	I	11	Total O 11 11	0	0
3	J	7	Total O 7 7	0	0
3	K	6	Total O 6 6	0	0
3	L	9	Total O 9 9	0	0
3	M	9	Total O 9 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase Mdm2




- Molecule 1: E3 ubiquitin-protein ligase Mdm2




- Molecule 1: E3 ubiquitin-protein ligase Mdm2




- Molecule 1: E3 ubiquitin-protein ligase Mdm2





- Molecule 1: E3 ubiquitin-protein ligase Mdm2





- Molecule 1: E3 ubiquitin-protein ligase Mdm2



- Molecule 2: Modified p53 peptide



- Molecule 2: Modified p53 peptide



- Molecule 2: Modified p53 peptide





- Molecule 2: Modified p53 peptide



- Molecule 2: Modified p53 peptide



- Molecule 2: Modified p53 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.81Å 76.08Å 197.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.32 – 2.07 49.32 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.32-2.07) 99.5 (49.32-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.07Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.187 , 0.238 0.195 , 0.249	Depositor DCC
R_{free} test set	2100 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtrriage
Anisotropy	0.664	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5145	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 9FB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.69	0/761	0.82	0/1025
1	B	0.80	0/742	0.85	0/999
1	C	0.67	0/731	0.81	0/984
1	D	1.01	1/742 (0.1%)	1.13	4/999 (0.4%)
1	E	0.75	0/733	0.89	0/986
1	F	0.70	0/742	0.80	0/999
2	H	0.88	0/99	0.80	0/134
2	I	1.00	0/99	0.79	0/134
2	J	0.75	0/99	0.69	0/134
2	K	0.70	0/99	0.76	0/134
2	L	0.78	0/99	0.70	0/134
2	M	0.76	0/99	0.68	0/134
All	All	0.78	1/5045 (0.0%)	0.87	4/6796 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1
2	I	0	1
2	J	0	1
2	M	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	109	VAL	C-N	17.40	1.74	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	109	VAL	O-C-N	-17.22	95.15	122.70
1	D	109	VAL	C-N-CA	9.18	144.64	121.70
1	D	50[A]	MET	CG-SD-CE	6.49	110.59	100.20
1	D	50[B]	MET	CG-SD-CE	6.49	110.59	100.20

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	27	9FB	Mainchain
2	I	27	9FB	Mainchain
2	J	27	9FB	Mainchain
2	M	27	9FB	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	747	0	779	18	0
1	B	725	0	764	5	0
1	C	714	0	754	11	0
1	D	725	0	763	8	0
1	E	720	0	754	9	0
1	F	725	0	764	5	0
2	H	104	0	85	1	0
2	I	104	0	85	0	0
2	J	104	0	85	0	0
2	K	104	0	85	1	0
2	L	104	0	85	0	0
2	M	104	0	85	2	0
3	A	12	0	0	0	0
3	B	29	0	0	0	0
3	C	5	0	0	0	0
3	D	33	0	0	0	0
3	E	20	0	0	0	0
3	F	15	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	9	0	0	0	0
3	I	11	0	0	0	0
3	J	7	0	0	0	0
3	K	6	0	0	0	0
3	L	9	0	0	0	0
3	M	9	0	0	0	0
All	All	5145	0	5088	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:109:VAL:C	1:D:110:VAL:N	1.74	1.40
1:C:28:VAL:HG11	1:C:107:LEU:HD22	1.62	0.81
1:F:101:THR:O	1:F:105:ARG:HG3	1.85	0.77
1:E:109:VAL:O	1:E:110:VAL:N	2.20	0.74
1:D:109:VAL:O	1:D:110:VAL:N	2.21	0.72
1:A:82:LEU:HA	1:A:85:LEU:CD1	2.26	0.66
1:C:28:VAL:HG13	1:C:107:LEU:HD13	1.78	0.66
1:C:86:PHE:CE1	1:C:102[A]:MET:HG2	2.31	0.66
1:A:28:VAL:HG22	1:A:107:LEU:HD21	1.77	0.65
1:D:50[A]:MET:HG2	1:D:51:LYS:N	2.12	0.64
1:F:30:PRO:HA	1:F:107:LEU:HD23	1.82	0.61
1:C:37:LEU:HD12	1:C:81:LEU:HD23	1.82	0.61
1:A:82:LEU:HA	1:A:85:LEU:HD12	1.83	0.60
1:A:78:SER:O	1:A:79:ASN:C	2.38	0.60
1:E:109:VAL:C	1:E:110:VAL:N	2.55	0.60
1:A:107:LEU:HD23	1:A:108:VAL:N	2.16	0.59
1:A:28:VAL:HG13	1:A:107:LEU:HD21	1.87	0.57
1:A:33:LEU:CD2	1:A:85:LEU:HG	2.37	0.55
1:C:28:VAL:CG1	1:C:107:LEU:HD22	2.36	0.53
1:A:28:VAL:CG2	1:A:107:LEU:HD21	2.39	0.52
1:E:37:LEU:C	1:E:37:LEU:HD23	2.30	0.52
1:A:47:THR:HG22	1:A:47:THR:O	2.09	0.52
1:A:28:VAL:HG13	1:A:107:LEU:CD2	2.40	0.52
1:C:37:LEU:CD1	1:C:81:LEU:HD23	2.40	0.52
1:C:37:LEU:HD12	1:C:81:LEU:CD2	2.40	0.50
1:D:50[A]:MET:HE2	1:D:51:LYS:HB3	1.94	0.50
1:A:78:SER:O	1:A:80:ASP:N	2.45	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:108:VAL:HG12	1:C:109:VAL:N	2.27	0.50
1:E:73:HIS:CD2	1:E:94:LYS:HE3	2.48	0.49
1:F:37:LEU:C	1:F:37:LEU:HD23	2.33	0.49
1:A:91:PHE:CD1	1:A:99:ILE:HD11	2.48	0.49
1:D:37:LEU:HD23	1:D:37:LEU:C	2.33	0.49
1:A:31:LYS:O	1:A:32:PRO:C	2.52	0.48
1:B:107:LEU:C	1:B:107:LEU:HD12	2.33	0.48
1:E:109:VAL:O	1:E:110:VAL:CA	2.62	0.47
2:M:27:9FB:O	2:M:27:9FB:N	2.49	0.45
1:B:37:LEU:C	1:B:37:LEU:HD23	2.37	0.45
1:D:50[A]:MET:CG	1:D:51:LYS:N	2.79	0.44
1:C:28:VAL:CG1	1:C:107:LEU:HD13	2.46	0.44
1:A:99:ILE:HG21	2:H:26:LEU:HD21	1.99	0.44
1:A:28:VAL:CG1	1:A:107:LEU:HD21	2.47	0.44
1:A:33:LEU:HD23	1:A:85:LEU:HG	1.99	0.43
1:D:95:GLU:O	1:D:99:ILE:HG12	2.18	0.43
1:B:76:TYR:CZ	1:B:78:SER:OG	2.71	0.43
1:A:39:LYS:HG2	1:A:43:ALA:O	2.19	0.42
1:B:86:PHE:HB3	1:B:88:VAL:HG22	2.00	0.42
1:F:51:LYS:HA	2:M:28:PRO:O	2.19	0.42
1:A:77:CYS:HB3	1:A:83:GLY:HA2	2.03	0.41
1:E:50:MET:HE3	1:E:50:MET:HB2	1.78	0.41
1:F:50:MET:CE	1:F:103:ILE:HG21	2.51	0.41
1:C:68:ASP:HB3	1:C:71:GLN:O	2.20	0.41
1:E:93:VAL:O	1:E:93:VAL:HG12	2.19	0.41
1:E:109:VAL:HA	1:E:110:VAL:N	2.36	0.40
1:D:51:LYS:HE2	2:K:28:PRO:HG2	2.04	0.40
1:E:109:VAL:O	1:E:110:VAL:HA	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/90 (98%)	86 (98%)	1 (1%)	1 (1%)	14	5
1	B	86/90 (96%)	86 (100%)	0	0	100	100
1	C	85/90 (94%)	82 (96%)	3 (4%)	0	100	100
1	D	86/90 (96%)	85 (99%)	1 (1%)	0	100	100
1	E	83/90 (92%)	83 (100%)	0	0	100	100
1	F	86/90 (96%)	85 (99%)	1 (1%)	0	100	100
2	H	11/14 (79%)	11 (100%)	0	0	100	100
2	I	11/14 (79%)	11 (100%)	0	0	100	100
2	J	11/14 (79%)	11 (100%)	0	0	100	100
2	K	11/14 (79%)	11 (100%)	0	0	100	100
2	L	11/14 (79%)	11 (100%)	0	0	100	100
2	M	11/14 (79%)	11 (100%)	0	0	100	100
All	All	580/624 (93%)	573 (99%)	6 (1%)	1 (0%)	47	39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	79	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	83 (98%)	2 (2%)	49	43
1	B	83/85 (98%)	81 (98%)	2 (2%)	49	43
1	C	81/85 (95%)	78 (96%)	3 (4%)	34	27
1	D	83/85 (98%)	79 (95%)	4 (5%)	25	18
1	E	82/85 (96%)	79 (96%)	3 (4%)	34	27
1	F	83/85 (98%)	79 (95%)	4 (5%)	25	18
2	H	9/9 (100%)	9 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	I	9/9 (100%)	9 (100%)	0	100	100
2	J	9/9 (100%)	9 (100%)	0	100	100
2	K	9/9 (100%)	9 (100%)	0	100	100
2	L	9/9 (100%)	9 (100%)	0	100	100
2	M	9/9 (100%)	9 (100%)	0	100	100
All	All	551/564 (98%)	533 (97%)	18 (3%)	42	31

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LEU
1	A	47	THR
1	B	27	LEU
1	B	69	GLU
1	C	78	SER
1	C	102[A]	MET
1	C	102[B]	MET
1	D	25	GLU
1	D	50[A]	MET
1	D	50[B]	MET
1	D	77	CYS
1	E	44	GLN
1	E	109	VAL
1	E	110	VAL
1	F	45	LYS
1	F	51	LYS
1	F	65	ARG
1	F	77	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	44	GLN
1	A	111	ASN
1	C	44	GLN
1	C	96	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	E	1
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	E	109:VAL	C	110:VAL	N	2.55
1	D	109:VAL	C	110:VAL	N	1.74

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	90/90 (100%)	0.36	4 (4%) 34 35	35, 58, 90, 109	0
1	B	87/90 (96%)	-0.13	0 100 100	30, 41, 60, 80	0
1	C	86/90 (95%)	0.15	1 (1%) 79 80	35, 57, 81, 100	0
1	D	87/90 (96%)	-0.00	1 (1%) 80 81	30, 39, 65, 86	0
1	E	87/90 (96%)	-0.06	0 100 100	33, 48, 66, 89	0
1	F	87/90 (96%)	0.12	1 (1%) 80 81	34, 53, 74, 103	0
2	H	13/14 (92%)	-0.00	0 100 100	36, 40, 56, 59	0
2	I	13/14 (92%)	0.03	1 (7%) 13 13	32, 40, 49, 56	0
2	J	13/14 (92%)	0.08	0 100 100	31, 36, 52, 65	0
2	K	13/14 (92%)	-0.10	0 100 100	33, 38, 43, 50	0
2	L	13/14 (92%)	0.11	1 (7%) 13 13	38, 44, 52, 52	0
2	M	13/14 (92%)	0.69	2 (15%) 2 1	40, 47, 65, 80	0
All	All	602/624 (96%)	0.08	11 (1%) 68 70	30, 47, 79, 109	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	26	THR	5.0
1	F	111	ASN	4.7
2	M	16	ALA	4.0
2	M	28	PRO	3.4
1	A	24	GLN	3.1
1	A	79	ASN	2.7
1	C	26	THR	2.5
2	L	16	ALA	2.3
1	A	111	ASN	2.2
1	D	50[A]	MET	2.2
2	I	16	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.