



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2021 – 05:09 am BST

PDB ID : 7AYE
Title : Crystal structure of the computationally designed chemically disruptable heterodimer LD6-MDM2
Authors : Yang, C.; Lau, K.; Pojer, F.; Correia, B.E.
Deposited on : 2020-11-12
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	 3% 51% 24% • 25%
2	B	137	 3% 67% 18% • 13%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform 11 of E3 ubiquitin-protein ligase Mdm2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	86	712	467	117	124	4	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q00987
A	1	GLU	-	expression tag	UNP Q00987
A	2	PHE	-	expression tag	UNP Q00987
A	112	LEU	-	expression tag	UNP Q00987
A	113	GLU	-	expression tag	UNP Q00987

- Molecule 2 is a protein called Thiol:disulfide interchange protein DsbD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	119	944	607	158	175	4	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP Q328D2
B	20	ALA	VAL	engineered mutation	UNP Q328D2
B	21	PHE	ASP	engineered mutation	UNP Q328D2
B	22	ALA	GLU	engineered mutation	UNP Q328D2
B	24	TYR	ASN	engineered mutation	UNP Q328D2
B	25	TRP	GLN	engineered mutation	UNP Q328D2
B	28	LEU	VAL	engineered mutation	UNP Q328D2
B	31	GLN	LYS	engineered mutation	UNP Q328D2
B	130	LEU	-	expression tag	UNP Q328D2
B	131	GLU	-	expression tag	UNP Q328D2
B	132	HIS	-	expression tag	UNP Q328D2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	133	HIS	-	expression tag	UNP Q328D2
B	134	HIS	-	expression tag	UNP Q328D2
B	135	HIS	-	expression tag	UNP Q328D2
B	136	HIS	-	expression tag	UNP Q328D2
B	137	HIS	-	expression tag	UNP Q328D2

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	73.24Å 73.24Å 92.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.20 – 2.95 45.16 – 2.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.20-2.95) 99.9 (45.16-2.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.197 , 0.260 0.201 , 0.259	Depositor DCC
R_{free} test set	285 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	90.9	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1656	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/726	0.88	0/978
2	B	0.67	0/966	0.87	0/1314
All	All	0.69	0/1692	0.87	0/2292

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	712	0	749	12	0
2	B	944	0	920	11	0
All	All	1656	0	1669	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:THR:OG1	1:A:38:GLU:HG3	1.92	0.69
2:B:30:ALA:O	2:B:31:GLN:O	2.16	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:24:TYR:OH	2:B:87:HIS:CE1	2.52	0.62
2:B:67:THR:OG1	2:B:127:ARG:NH1	2.32	0.62
1:A:63:CYS:HB2	1:A:69:GLY:N	2.23	0.54
1:A:69:GLY:O	1:A:73:GLY:N	2.42	0.51
1:A:31:LYS:HB2	1:A:34:TYR:CE2	2.45	0.51
1:A:47:ILE:HG23	1:A:52:LEU:HB2	1.93	0.50
2:B:127:ARG:HD2	2:B:128:GLN:HB2	1.94	0.49
2:B:76:ALA:O	2:B:77:ASN:C	2.52	0.48
2:B:88:LEU:O	2:B:110:ARG:NH2	2.46	0.47
1:A:37:LYS:HG3	1:A:38:GLU:N	2.29	0.46
1:A:64:SER:O	1:A:65:ASN:HB2	2.16	0.46
1:A:63:CYS:HB2	1:A:69:GLY:CA	2.46	0.46
2:B:57:ASP:OD1	2:B:57:ASP:C	2.54	0.45
1:A:89:ILE:O	1:A:92:ASN:N	2.49	0.44
2:B:24:TYR:HH	2:B:87:HIS:CE1	2.35	0.43
2:B:78:ASP:O	2:B:81:ASP:HB2	2.20	0.42
1:A:94:VAL:HG22	1:A:95:VAL:H	1.84	0.42
2:B:42:ASP:O	2:B:48:LYS:HD2	2.20	0.42
2:B:24:TYR:OH	2:B:87:HIS:NE2	2.53	0.41
1:A:84:LYS:O	1:A:88:MET:HG3	2.21	0.41
1:A:17:LYS:HB3	1:A:18:PRO:HD2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	84/114 (74%)	77 (92%)	7 (8%)	0	100	100
2	B	117/137 (85%)	110 (94%)	4 (3%)	3 (3%)	5	24
All	All	201/251 (80%)	187 (93%)	11 (6%)	3 (2%)	10	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	31	GLN
2	B	77	ASN
2	B	60	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/107 (76%)	72 (89%)	9 (11%)	6	22
2	B	99/115 (86%)	89 (90%)	10 (10%)	7	26
All	All	180/222 (81%)	161 (89%)	19 (11%)	6	24

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	ARG
1	A	19	LEU
1	A	24	LEU
1	A	26	SER
1	A	33	THR
1	A	57	GLN
1	A	63	CYS
1	A	83	ARG
1	A	96	VAL
2	B	10	THR
2	B	15	THR
2	B	17	ILE
2	B	39	LEU
2	B	45	VAL
2	B	50	PHE
2	B	52	LYS
2	B	86	LYS
2	B	105	GLU
2	B	119	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	87	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	86/114 (75%)	-0.08	0 100 100	60, 86, 118, 127	0
2	B	119/137 (86%)	0.15	4 (3%) 45 29	62, 88, 119, 149	0
All	All	205/251 (81%)	0.06	4 (1%) 65 48	60, 87, 118, 149	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	64	LEU	3.7
2	B	106	HIS	2.8
2	B	102	GLN	2.5
2	B	66	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.