



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 02:26 AM JST

PDB ID : 7EKG
Title : Structure of SARS-CoV-2 Beta variant spike receptor-binding domain complexed with human ACE2
Authors : Han, P.C.; Su, C.; Zhang, Y.F.; Qi, J.X.; Gao, G.F.
Deposited on : 2021-04-05
Resolution : 2.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	603	
2	B	229	
3	C	2	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiotensin-converting enzyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	596	4862	3111	805	917	29	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	616	HIS	-	expression tag	UNP Q9BYF1
A	617	HIS	-	expression tag	UNP Q9BYF1
A	618	HIS	-	expression tag	UNP Q9BYF1
A	619	HIS	-	expression tag	UNP Q9BYF1
A	620	HIS	-	expression tag	UNP Q9BYF1
A	621	HIS	-	expression tag	UNP Q9BYF1


- Molecule 2 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	195	1552	997	258	289	8	0	1	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	417	ASN	LYS	engineered mutation	UNP P0DTC2
B	484	LYS	GLU	engineered mutation	UNP P0DTC2
B	501	TYR	ASN	engineered mutation	UNP P0DTC2
B	542	HIS	-	expression tag	UNP P0DTC2
B	543	HIS	-	expression tag	UNP P0DTC2
B	544	HIS	-	expression tag	UNP P0DTC2
B	545	HIS	-	expression tag	UNP P0DTC2
B	546	HIS	-	expression tag	UNP P0DTC2
B	547	HIS	-	expression tag	UNP P0DTC2

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Zn 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	18	Total 18	O 18	0	0
6	B	3	Total 3	O 3	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.15Å 105.15Å 230.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.91 – 2.63 27.01 – 2.63	Depositor EDS
% Data completeness (in resolution range)	99.9 (26.91-2.63) 100.0 (27.01-2.63)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.64Å)	Xtrriage
Refinement program	PHENIX 1.19_4092	Depositor
R, R_{free}	0.214 , 0.248 0.214 , 0.245	Depositor DCC
R_{free} test set	1944 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	64.8	Xtrriage
Anisotropy	0.360	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6534	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.27	0/4999	0.45	0/6792
2	B	0.47	4/1600 (0.2%)	0.55	2/2180 (0.1%)
All	All	0.33	4/6599 (0.1%)	0.48	2/8972 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	498[A]	GLN	C-N	8.39	1.50	1.34
2	B	498[B]	GLN	C-N	8.39	1.50	1.34
2	B	498[A]	GLN	C-O	6.22	1.35	1.23
2	B	498[B]	GLN	C-O	6.22	1.35	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	498[A]	GLN	O-C-N	-6.32	109.09	121.10
2	B	498[B]	GLN	O-C-N	-6.32	109.09	121.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4862	0	4634	35	1
2	B	1552	0	1470	20	0
3	C	28	0	25	0	0
4	A	56	0	52	0	0
4	B	14	0	13	0	0
5	A	1	0	0	0	0
6	A	18	0	0	0	0
6	B	3	0	0	0	0
All	All	6534	0	6194	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:VAL:HG11	1:A:194:ASN:ND2	1.66	1.10
1:A:107:VAL:HG11	1:A:194:ASN:HD21	1.28	0.89
1:A:107:VAL:CG1	1:A:194:ASN:HD21	2.00	0.73
1:A:490:PRO:HA	1:A:612:PRO:HG2	1.73	0.69
1:A:105:SER:O	1:A:105:SER:OG	2.16	0.64
2:B:520:ALA:HB1	2:B:521:PRO:HD2	1.79	0.64
1:A:607:SER:OG	1:A:609:ASP:OD1	2.20	0.60
2:B:366:SER:HB3	2:B:388:ASN:HD21	1.66	0.60
1:A:107:VAL:HG21	1:A:193:ALA:HB3	1.85	0.59
2:B:393:THR:HA	2:B:522:ALA:HA	1.84	0.58
1:A:560:LEU:HD22	1:A:564:GLU:HG3	1.84	0.57
1:A:432:ASN:HA	1:A:435:GLU:HG3	1.87	0.55
1:A:107:VAL:CG2	1:A:193:ALA:HB3	2.37	0.55
1:A:85:LEU:HG	1:A:101:GLN:HE22	1.72	0.54
1:A:131:LYS:HD3	1:A:141:CYS:HB2	1.90	0.52
2:B:521:PRO:HB2	2:B:523:THR:HG23	1.92	0.52
1:A:419:LYS:HE3	1:A:426:PRO:HA	1.92	0.52
2:B:379:CYS:HB2	2:B:384:PRO:HD3	1.91	0.51
2:B:393:THR:HG21	2:B:518:LEU:H	1.76	0.50
2:B:474:GLN:NE2	2:B:479:PRO:HA	2.27	0.50
2:B:334:ASN:O	2:B:362:VAL:N	2.39	0.50
1:A:55:THR:O	1:A:59:VAL:HG23	2.11	0.50
1:A:379:ILE:O	1:A:383:MET:HG3	2.12	0.49
2:B:383:SER:H	2:B:387:LEU:HD22	1.76	0.49
1:A:81:GLN:HG3	1:A:101:GLN:HG2	1.95	0.49
2:B:455:LEU:HD22	2:B:493:GLN:HG3	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:PRO:HB3	1:A:360:MET:HG3	1.94	0.48
1:A:144:LEU:HA	1:A:148:LEU:HB2	1.96	0.47
2:B:366:SER:HB3	2:B:388:ASN:ND2	2.30	0.47
1:A:140:GLU:HA	1:A:140:GLU:OE1	2.15	0.46
2:B:395:VAL:HG22	2:B:515:PHE:HD1	1.79	0.46
2:B:382:VAL:HG11	2:B:515:PHE:HE2	1.80	0.46
2:B:383:SER:O	2:B:387:LEU:HB2	2.16	0.46
1:A:107:VAL:HG21	1:A:193:ALA:CB	2.46	0.46
1:A:107:VAL:CG2	1:A:193:ALA:CB	2.94	0.46
1:A:239:HIS:CE1	1:A:596:LYS:HG2	2.51	0.45
2:B:447:GLY:HA2	2:B:498[A]:GLN:HG2	1.98	0.45
1:A:293:VAL:O	1:A:297:MET:HG3	2.17	0.45
1:A:294:THR:HG23	1:A:365:THR:HA	1.98	0.45
1:A:524:GLN:HG2	1:A:583:PRO:HG2	1.98	0.44
1:A:315:PHE:CZ	1:A:408:MET:HG3	2.52	0.44
1:A:131:LYS:HD3	1:A:141:CYS:CB	2.47	0.44
1:A:339:VAL:O	1:A:341:LYS:HG2	2.18	0.44
2:B:362:VAL:CG1	2:B:526:GLY:HA2	2.47	0.44
2:B:391:CYS:HA	2:B:525:CYS:HA	2.00	0.43
1:A:64:ASN:O	1:A:68:LYS:HG3	2.18	0.43
1:A:57:GLU:O	1:A:61:ASN:ND2	2.49	0.43
1:A:100:LEU:C	1:A:102:GLN:N	2.72	0.43
1:A:549:GLU:H	1:A:549:GLU:CD	2.23	0.42
2:B:502:GLY:O	2:B:506:GLN:HG3	2.20	0.41
2:B:362:VAL:HG13	2:B:526:GLY:HA2	2.01	0.41
1:A:74:LYS:HG2	1:A:105:SER:HB2	2.03	0.41
1:A:415:PRO:O	1:A:419:LYS:HG3	2.20	0.41
2:B:334:ASN:O	2:B:362:VAL:HG23	2.21	0.41
1:A:524:GLN:HB3	1:A:574:VAL:HG11	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:609:ASP:OD1	1:A:609:ASP:OD1[8_554]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	594/603 (98%)	588 (99%)	6 (1%)	0	100	100
2	B	194/229 (85%)	184 (95%)	10 (5%)	0	100	100
All	All	788/832 (95%)	772 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	526/533 (99%)	513 (98%)	13 (2%)	47	66
2	B	169/202 (84%)	165 (98%)	4 (2%)	49	67
All	All	695/735 (95%)	678 (98%)	17 (2%)	49	67

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	LEU
1	A	56	GLU
1	A	102	GLN
1	A	103	ASN
1	A	107	VAL
1	A	167	SER
1	A	236	LEU
1	A	338	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	363	LYS
1	A	381	TYR
1	A	401	HIS
1	A	430	GLU
1	A	559	ARG
2	B	377	PHE
2	B	382	VAL
2	B	424	LYS
2	B	445	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	A	194	ASN
2	B	474	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,1	14,14,15	0.28	0	17,19,21	0.56	0
3	NAG	C	2	3	14,14,15	0.32	0	17,19,21	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	NAG	O5-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	702	1	14,14,15	0.28	0	17,19,21	0.46	0
4	NAG	A	701	1	14,14,15	0.40	0	17,19,21	0.61	1 (5%)
4	NAG	A	705	1	14,14,15	0.31	0	17,19,21	0.72	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	704	1	14,14,15	0.28	0	17,19,21	0.51	0
4	NAG	B	601	2	14,14,15	0.28	0	17,19,21	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	702	1	-	0/6/23/26	0/1/1/1
4	NAG	A	701	1	-	0/6/23/26	0/1/1/1
4	NAG	A	705	1	-	0/6/23/26	0/1/1/1
4	NAG	A	704	1	-	2/6/23/26	0/1/1/1
4	NAG	B	601	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	705	NAG	C1-O5-C5	2.63	115.76	112.19
4	A	701	NAG	C1-O5-C5	2.08	115.01	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	704	NAG	C8-C7-N2-C2
4	A	704	NAG	O7-C7-N2-C2



There are no ring outliers.



No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	596/603 (98%)	0.08	31 (5%) 27 24	46, 63, 108, 183	0
2	B	195/229 (85%)	0.45	22 (11%) 5 3	52, 75, 149, 183	0
All	All	791/832 (95%)	0.17	53 (6%) 17 15	46, 66, 125, 183	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	519	HIS	7.0
2	B	333	THR	5.6
2	B	521	PRO	5.6
1	A	136	ASP	5.5
1	A	138	PRO	5.4
1	A	103	ASN	5.3
2	B	520	ALA	5.3
1	A	107	VAL	5.2
2	B	518	LEU	4.5
1	A	105	SER	4.4
1	A	338	ASN	4.1
2	B	385	THR	4.0
1	A	108	LEU	3.7
2	B	525	CYS	3.7
2	B	523	THR	3.6
2	B	372	ALA	3.6
1	A	455	MET	3.5
1	A	427	ASP	3.5
2	B	522	ALA	3.4
1	A	139	GLN	3.3
2	B	432	CYS	3.1
2	B	384	PRO	3.1
1	A	339	VAL	2.9
2	B	335	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	452	PHE	2.8
1	A	614	ALA	2.7
2	B	386	LYS	2.7
1	A	429	GLN	2.6
2	B	370	ASN	2.6
2	B	479	PRO	2.5
1	A	133	CYS	2.5
2	B	527	PRO	2.5
1	A	456	LEU	2.5
2	B	369	TYR	2.4
1	A	141	CYS	2.4
1	A	140	GLU	2.4
2	B	389	ASP	2.4
1	A	408	MET	2.3
1	A	341	LYS	2.3
1	A	315	PHE	2.3
1	A	68	LYS	2.3
1	A	377	GLY	2.3
1	A	79	LEU	2.3
2	B	460	ASN	2.2
1	A	428	PHE	2.2
2	B	511	VAL	2.2
1	A	311	ALA	2.2
1	A	411	SER	2.2
1	A	568	LEU	2.1
1	A	69	TRP	2.1
2	B	526	GLY	2.1
1	A	603	PHE	2.1
1	A	251	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	C	2	14/15	0.79	0.36	123,137,145,147	0
3	NAG	C	1	14/15	0.93	0.13	97,105,109,112	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	704	14/15	0.69	0.35	119,132,139,139	0
4	NAG	A	705	14/15	0.75	0.25	114,130,135,139	0
4	NAG	A	702	14/15	0.77	0.33	116,120,127,128	0
4	NAG	A	701	14/15	0.87	0.23	104,111,116,118	0
4	NAG	B	601	14/15	0.88	0.14	97,105,108,109	0
5	ZN	A	703	1/1	0.97	0.16	48,48,48,48	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NAG A 705:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAG A 702:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NAG A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.