



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 30, 2022 – 10:05 pm BST

PDB ID : 7NFV
Title : Structure of SARS-CoV-2 Papain-like protease PLpro
Authors : Srinivasan, V.; Gunther, S.; Reinke, P.; Werner, N.; Falke, S.; Brognaro, H.; Ullah, N.; Andaleeb, H.; Perbandt, M.; Alves Franca, B.; Schwinzer, M.; Wang, M.; Sprenger, J.; Lieske, J.; Ginn, H.; Lane, T.J.; Yefanov, O.; Gelisio, L.; Koua, F.; Saouane, S.; Tolstikova, A.; Groessler, M.; Fleckenstein, H.; Ewert, W.; Trost, F.; Lorenzen, K.; Schubert, R.; Han, H.; Schmidt, C.; Brings, L.; Ehrt, C.; Rarey, M.; Galchenkova, M.; Gevorkov, Y.; Li, C.; Perk, A.; Awel, S.; Hinrichs, W.; Meents, A.; Betzel, C.
Deposited on : 2021-02-07
Resolution : 1.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.30
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2579 (1.44-1.40)
Clashscore	141614	2696 (1.44-1.40)
Ramachandran outliers	138981	2632 (1.44-1.40)
Sidechain outliers	138945	2631 (1.44-1.40)
RSRZ outliers	127900	2528 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	315	

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.30

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Papain-like protease nsp3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	315	2674	1692	445	517	20	0	20	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	AAA	1	1	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	AAA	1	12	6	6	0	1

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	2	Total Cl 2 2	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	AAA	1	Total O P 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AAA	530	Total O 530 530	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Papain-like protease nsp3



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	82.33Å 82.33Å 134.32Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.89 – 1.42 48.89 – 1.42	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.89-1.42) 100.0 (48.89-1.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.42Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.154 , 0.171 0.160 , 0.175	Depositor DCC
R_{free} test set	4919 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtrriage
Anisotropy	0.053	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3224	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	1.09	11/2736 (0.4%)	1.24	22/3705 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AAA	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	51[A]	GLU	CD-OE2	8.39	1.34	1.25
1	AAA	51[B]	GLU	CD-OE2	8.39	1.34	1.25
1	AAA	203	GLU	CD-OE2	8.39	1.34	1.25
1	AAA	161	GLU	CD-OE2	-8.35	1.16	1.25
1	AAA	280	GLU	CD-OE2	8.28	1.34	1.25
1	AAA	78	SER	CA-CB	-7.14	1.42	1.52
1	AAA	67	GLU	CD-OE2	6.47	1.32	1.25
1	AAA	167	GLU	CD-OE2	5.40	1.31	1.25
1	AAA	171	TYR	CG-CD2	-5.25	1.32	1.39
1	AAA	180[A]	SER	CA-CB	-5.12	1.45	1.52
1	AAA	180[B]	SER	CA-CB	-5.12	1.45	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	140	ARG	NE-CZ-NH1	9.10	124.85	120.30
1	AAA	140	ARG	NE-CZ-NH2	-8.94	115.83	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	82	ARG	NE-CZ-NH1	-8.51	116.05	120.30
1	AAA	108	ASP	CB-CG-OD1	-8.44	110.70	118.30
1	AAA	82	ARG	NE-CZ-NH2	8.05	124.33	120.30
1	AAA	268[A]	TYR	CA-C-O	-6.63	106.17	120.10
1	AAA	268[B]	TYR	CA-C-O	-6.63	106.17	120.10
1	AAA	35	TYR	CB-CG-CD2	-6.46	117.12	121.00
1	AAA	166	ARG	NE-CZ-NH1	6.46	123.53	120.30
1	AAA	12	ASP	CB-CG-OD1	-6.39	112.55	118.30
1	AAA	270	CYS	CB-CA-C	6.09	122.59	110.40
1	AAA	83	TYR	CB-CG-CD1	6.00	124.60	121.00
1	AAA	268[A]	TYR	O-C-N	5.89	132.13	122.70
1	AAA	268[B]	TYR	O-C-N	5.89	132.13	122.70
1	AAA	138	ARG	CD-NE-CZ	5.69	131.57	123.60
1	AAA	183	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	AAA	108	ASP	CB-CG-OD2	5.36	123.12	118.30
1	AAA	138	ARG	NE-CZ-NH1	5.35	122.98	120.30
1	AAA	207	TYR	CB-CG-CD1	-5.31	117.81	121.00
1	AAA	138	ARG	NE-CZ-NH2	-5.27	117.67	120.30
1	AAA	35	TYR	CB-CG-CD1	5.17	124.10	121.00
1	AAA	137	TYR	CB-CG-CD2	-5.03	117.98	121.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AAA	15	ASN	Sidechain
1	AAA	162	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2674	0	2578	19	0
2	AAA	1	0	0	0	0
3	AAA	12	0	16	0	0
4	AAA	2	0	0	0	0
5	AAA	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	AAA	530	0	0	13	3
All	All	3224	0	2594	19	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:174[B]:GLN:CG	6:AAA:572:HOH:O	2.22	0.87
1:AAA:174[B]:GLN:HG3	6:AAA:572:HOH:O	1.73	0.86
1:AAA:60:ASN:ND2	6:AAA:504:HOH:O	2.08	0.81
1:AAA:267:ASN:ND2	6:AAA:505:HOH:O	2.13	0.81
1:AAA:174[A]:GLN:HG2	1:AAA:174[A]:GLN:O	1.80	0.81
1:AAA:143[A]:GLU:OE2	6:AAA:503:HOH:O	2.02	0.78
1:AAA:236[B]:GLN:NE2	1:AAA:238[B]:GLU:OE2	2.28	0.66
1:AAA:126[B]:LYS:NZ	6:AAA:506:HOH:O	2.19	0.61
1:AAA:171:TYR:O	1:AAA:175[B]:HIS:HD2	1.83	0.60
1:AAA:174[A]:GLN:NE2	6:AAA:514:HOH:O	2.35	0.58
1:AAA:200:LYS:HD2	6:AAA:632:HOH:O	2.05	0.55
1:AAA:306:LYS:HE3	6:AAA:596:HOH:O	2.07	0.54
1:AAA:126[B]:LYS:HE2	6:AAA:869:HOH:O	2.10	0.52
1:AAA:174[B]:GLN:CD	6:AAA:572:HOH:O	2.47	0.51
1:AAA:197:THR:HG23	6:AAA:709:HOH:O	2.12	0.50
1:AAA:279[B]:LYS:NZ	6:AAA:518:HOH:O	2.41	0.45
1:AAA:59:PRO:HD3	1:AAA:80:LEU:HD12	2.00	0.44
1:AAA:263:GLU:O	1:AAA:273:TYR:HA	2.19	0.42
1:AAA:188:VAL:HG22	1:AAA:194:GLN:HG2	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:AAA:780:HOH:O	6:AAA:864:HOH:O[5_555]	1.97	0.23
6:AAA:579:HOH:O	6:AAA:1008:HOH:O[2_654]	2.09	0.11
6:AAA:692:HOH:O	6:AAA:831:HOH:O[5_555]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	333/315 (106%)	318 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	295/275 (107%)	291 (99%)	4 (1%)	67	38

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	60	ASN
1	AAA	229[A]	GLN
1	AAA	229[B]	GLN
1	AAA	270	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PO4	AAA	404	-	4,4,4	2.17	1 (25%)	6,6,6	0.77	0
3	GOL	AAA	402[B]	-	5,5,5	0.26	0	5,5,5	0.40	0
3	GOL	AAA	402[A]	-	5,5,5	0.29	0	5,5,5	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	AAA	402[B]	-	-	0/4/4/4	-
3	GOL	AAA	402[A]	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	AAA	404	PO4	P-O1	3.95	1.60	1.50

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	AAA	402[A]	GOL	O1-C1-C2-O2
3	AAA	402[A]	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	315/315 (100%)	0.25	12 (3%) 40 40	23, 30, 45, 71	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	2	VAL	6.7
1	AAA	315	LYS	6.3
1	AAA	314	ILE	4.6
1	AAA	3	ARG	4.2
1	AAA	1	GLU	4.0
1	AAA	47[A]	HIS	3.7
1	AAA	24	SER	3.1
1	AAA	313	THR	3.0
1	AAA	5	ILE	2.8
1	AAA	48	ASN	2.6
1	AAA	44	ILE	2.4
1	AAA	4	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	AAA	402[A]	6/6	0.78	0.23	42,51,52,52	6
3	GOL	AAA	402[B]	6/6	0.78	0.23	37,38,40,42	6
4	CL	AAA	405	1/1	0.96	0.07	42,42,42,42	0
5	PO4	AAA	404	5/5	0.97	0.21	25,31,33,36	5
2	ZN	AAA	401	1/1	0.99	0.03	35,35,35,35	0
4	CL	AAA	403	1/1	1.00	0.11	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around GOL AAA 402 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around CL AAA 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 AAA 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN AAA 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CL AAA 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.