



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2022 – 01:18 pm BST

PDB ID : 7P2D
Title : Structure of alphaMbeta2/Cd11bCD18 headpiece in complex with a nanobody
Authors : Jensen, R.K.; Andersen, G.R.
Deposited on : 2021-07-05
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	763	 6% 84% 13% ...
2	B	473	 6% 85% 11% ..
3	C	129	 28% 87% 7% 6%
4	D	2	 100%
4	E	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	2	 50% 50%
4	G	2	 50% 50%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform 2 of Integrin alpha-M.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	751	5802	3633	1040	1104	25	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	ARG	ASN	conflict	UNP P11215
A	681	ARG	ASN	conflict	UNP P11215
A	755	THR	CYS	conflict	UNP P11215
A	757	GLY	-	expression tag	UNP P11215
A	758	LEU	-	expression tag	UNP P11215
A	759	GLU	-	expression tag	UNP P11215
A	760	VAL	-	expression tag	UNP P11215
A	761	LEU	-	expression tag	UNP P11215
A	762	PHE	-	expression tag	UNP P11215
A	763	GLN	-	expression tag	UNP P11215

- Molecule 2 is a protein called Integrin beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	459	3573	2225	634	685	29	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	461	LYS	-	expression tag	UNP A0A494C0X7
B	462	ASN	-	expression tag	UNP A0A494C0X7
B	463	CYS	-	expression tag	UNP A0A494C0X7
B	464	GLU	-	expression tag	UNP A0A494C0X7
B	465	PRO	-	expression tag	UNP A0A494C0X7
B	466	ALA	-	expression tag	UNP A0A494C0X7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	467	ALA	-	expression tag	UNP A0A494C0X7
B	468	LEU	-	expression tag	UNP A0A494C0X7
B	469	GLN	-	expression tag	UNP A0A494C0X7
B	470	THR	-	expression tag	UNP A0A494C0X7
B	471	LEU	-	expression tag	UNP A0A494C0X7
B	472	PHE	-	expression tag	UNP A0A494C0X7
B	473	GLN	-	expression tag	UNP A0A494C0X7

- Molecule 3 is a protein called hCD11bNb1 nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	121	916	570	166	175	5	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	2	28	16	2	10	0	0	0
4	E	2	28	16	2	10	0	0	0
4	F	2	28	16	2	10	0	0	0
4	G	2	28	16	2	10	0	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	B	1	14	8	1	5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isoform 2 of Integrin alpha-M




- Molecule 2: Integrin beta



- Molecule 3: hCD11bNb1 nanobody

Chain C: 



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 


MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.10Å 114.10Å 250.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.95 – 3.20 49.41 – 3.01	Depositor EDS
% Data completeness (in resolution range)	82.8 (45.95-3.20) 72.7 (49.41-3.01)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.19rc5_4047	Depositor
R, R_{free}	0.261 , 0.295 0.260 , 0.291	Depositor DCC
R_{free} test set	1207 reflections (4.33%)	wwPDB-VP
Wilson B-factor (Å ²)	97.8	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	10419	wwPDB-VP
Average B, all atoms (Å ²)	127.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	1/5913 (0.0%)	0.79	16/7999 (0.2%)
2	B	0.35	0/3642	0.74	6/4927 (0.1%)
3	C	0.33	0/933	0.68	0/1262
All	All	0.36	1/10488 (0.0%)	0.76	22/14188 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	649	ARG	CG-CD	6.91	1.69	1.51

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	149	PHE	CB-CG-CD1	11.54	128.88	120.80
2	B	149	PHE	CB-CG-CD2	-10.24	113.63	120.80
1	A	176	TYR	CA-CB-CG	8.20	128.99	113.40
1	A	720	PHE	CB-CG-CD2	-7.99	115.21	120.80
2	B	247	PHE	CB-CG-CD2	-7.84	115.31	120.80
1	A	749	PHE	CB-CG-CD2	-7.57	115.50	120.80
1	A	720	PHE	CB-CG-CD1	7.33	125.93	120.80
1	A	649	ARG	CG-CD-NE	7.13	126.78	111.80
1	A	749	PHE	CB-CG-CD1	7.01	125.71	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	247	PHE	CB-CA-C	7.00	124.39	110.40
1	A	176	TYR	CB-CG-CD1	6.95	125.17	121.00
1	A	176	TYR	CB-CG-CD2	-6.79	116.93	121.00
2	B	247	PHE	CB-CG-CD1	6.49	125.34	120.80
1	A	720	PHE	CB-CA-C	6.11	122.61	110.40
1	A	390	TYR	CB-CG-CD2	-6.01	117.39	121.00
1	A	650	LEU	CA-CB-CG	5.98	129.06	115.30
1	A	487	ARG	CB-CG-CD	5.85	126.81	111.60
1	A	390	TYR	CA-CB-CG	5.51	123.86	113.40
1	A	649	ARG	CB-CA-C	-5.31	99.78	110.40
2	B	149	PHE	CB-CA-C	5.11	120.62	110.40
1	A	654	GLN	CA-CB-CG	5.07	124.56	113.40
1	A	487	ARG	CD-NE-CZ	5.06	130.69	123.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	115	ARG	Sidechain
1	A	487	ARG	Sidechain
1	A	649	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5802	0	5716	59	0
2	B	3573	0	3489	33	0
3	C	916	0	894	4	0
4	D	28	0	25	0	0
4	E	28	0	25	0	0
4	F	28	0	25	1	0
4	G	28	0	25	0	0
5	A	2	0	0	0	0
6	B	14	0	13	0	0
All	All	10419	0	10212	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:430:GLN:HG3	2:B:443:LEU:HB2	1.66	0.77
1:A:414:GLY:HA3	1:A:441:PHE:HB3	1.76	0.68
1:A:650:LEU:HG	1:A:652:GLU:H	1.60	0.67
1:A:663:LEU:HB2	1:A:684:ARG:HB3	1.77	0.66
2:B:322:GLU:O	2:B:329:ASN:ND2	2.30	0.65
1:A:665:LEU:HD11	1:A:701:LEU:HD22	1.82	0.62
1:A:103:VAL:HG12	1:A:334:MET:HE3	1.83	0.61
1:A:389:GLY:HA2	1:A:405:LEU:HD12	1.82	0.60
1:A:71:MET:HE2	1:A:73:LEU:HD13	1.84	0.60
1:A:662:ASP:OD2	4:F:1:NAG:N2	2.34	0.59
1:A:364:LEU:HB3	1:A:372:THR:HB	1.86	0.57
1:A:737:LEU:HD23	1:A:743:ARG:HH21	1.70	0.57
1:A:86:LEU:HB3	1:A:110:PHE:HB2	1.86	0.57
2:B:177:GLN:HB2	2:B:203:LEU:HB2	1.87	0.57
2:B:242:ASP:O	2:B:303:ARG:NH2	2.37	0.57
1:A:1:PHE:HA	1:A:554:GLN:HB2	1.87	0.57
3:C:4:GLN:HB2	3:C:26:SER:HB3	1.88	0.56
2:B:67:GLU:HG2	2:B:78:PRO:HB3	1.87	0.56
1:A:335:SER:HB2	1:A:352:THR:HB	1.86	0.56
1:A:123:GLU:H	1:A:368:LYS:HE3	1.71	0.55
1:A:103:VAL:HB	2:B:155:LEU:HD13	1.89	0.55
1:A:562:SER:HB3	1:A:565:LEU:HG	1.88	0.55
1:A:713:PRO:HB3	1:A:750:PRO:HA	1.89	0.54
1:A:337:GLU:HB3	1:A:350:LEU:HD13	1.89	0.54
1:A:347:GLY:HA3	1:A:366:THR:HA	1.88	0.54
1:A:579:THR:HG21	1:A:598:ARG:HD2	1.90	0.53
1:A:178:GLU:HG3	1:A:208:ARG:H	1.73	0.53
1:A:385:ASP:OD2	2:B:210:ALA:HB3	2.09	0.53
2:B:358:VAL:HG22	2:B:393:VAL:HG22	1.90	0.53
2:B:361:ASP:HB2	2:B:390:GLN:HB3	1.91	0.53
1:A:360:GLY:HA3	1:A:388:LEU:HB3	1.89	0.53
2:B:287:HIS:NE2	2:B:291:GLU:OE2	2.42	0.53
1:A:24:GLN:HE21	1:A:587:THR:HG23	1.75	0.52
1:A:330:PHE:HB3	1:A:333:GLU:HG3	1.92	0.52
2:B:120:ASP:N	2:B:120:ASP:OD1	2.43	0.51
1:A:362:VAL:HG21	1:A:405:LEU:HD11	1.93	0.50
2:B:317:LYS:NZ	2:B:344:ARG:O	2.45	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:500:GLN:HG2	1:A:531:GLU:H	1.76	0.50
1:A:579:THR:HG23	1:A:585:ASP:OD2	2.13	0.49
1:A:644:LYS:NZ	1:A:648:ASP:OD1	2.45	0.49
1:A:145:ILE:HD12	1:A:302:PHE:HZ	1.76	0.49
1:A:446:CYS:HB3	1:A:460:LEU:HB2	1.95	0.49
1:A:720:PHE:CD1	1:A:743:ARG:HG3	2.48	0.49
2:B:105:ILE:HG12	2:B:234:THR:HG21	1.95	0.49
1:A:245:LYS:HD3	1:A:248:ASP:HB3	1.94	0.48
2:B:41:ASP:OD2	2:B:45:GLN:HG2	2.14	0.48
1:A:172:SER:HB3	1:A:222:LEU:HD11	1.94	0.48
1:A:337:GLU:HG2	1:A:352:THR:HG22	1.95	0.48
2:B:150:VAL:HG12	2:B:151:ASP:H	1.80	0.47
1:A:515:VAL:HG12	1:A:522:ASP:OD2	2.15	0.47
1:A:149:ASP:HB3	1:A:302:PHE:CG	2.51	0.46
1:A:716:LEU:HB3	1:A:747:ALA:HB3	1.97	0.46
1:A:666:ASP:OD2	1:A:673:ARG:HB2	2.15	0.46
2:B:301:THR:HG21	2:B:325:GLU:OE2	2.15	0.46
3:C:108:TRP:O	3:C:111:TRP:NE1	2.49	0.46
1:A:94:HIS:CD2	2:B:155:LEU:HD12	2.51	0.46
1:A:76:SER:OG	1:A:339:PHE:O	2.33	0.46
2:B:435:SER:HB2	2:B:443:LEU:HG	1.97	0.46
3:C:7:GLU:OE2	3:C:112:GLY:HA3	2.16	0.46
1:A:138:LEU:HG	1:A:176:TYR:HD2	1.79	0.45
1:A:339:PHE:HE2	1:A:390:TYR:CD1	2.34	0.45
1:A:415:LEU:HD13	1:A:433:LYS:HG2	1.97	0.45
1:A:139:ILE:HG22	1:A:240:ILE:HD12	1.99	0.45
1:A:354:GLY:HA2	1:A:358:TRP:HA	1.99	0.45
2:B:152:LYS:NZ	2:B:216:ASP:OD2	2.40	0.45
2:B:330:VAL:HB	2:B:332:HIS:HB3	1.99	0.45
1:A:31:VAL:HG11	1:A:86:LEU:HD21	1.98	0.44
2:B:428:ARG:HD3	2:B:428:ARG:HA	1.72	0.44
1:A:140:ASP:HB2	1:A:241:THR:HA	1.99	0.44
3:C:91:THR:HG23	3:C:118:THR:HA	1.99	0.44
1:A:51:ASP:OD1	1:A:52:TYR:N	2.51	0.43
1:A:61:ARG:HD2	1:A:115:ARG:HA	1.99	0.43
1:A:106:LEU:HD21	1:A:120:LYS:HE3	2.01	0.43
2:B:173:GLU:OE2	2:B:175:GLU:HB2	2.18	0.43
2:B:106:ASP:OD2	2:B:231:ARG:HB2	2.18	0.43
2:B:43:ARG:HG3	2:B:56:ILE:HD13	2.01	0.43
1:A:439:ALA:HB1	1:A:466:TYR:HB2	2.01	0.42
1:A:440:TYR:CD1	2:B:247:PHE:HE1	2.37	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:105:ILE:HA	2:B:234:THR:HB	2.01	0.42
2:B:76:LEU:HD13	2:B:97:PHE:HD1	1.84	0.42
1:A:7:ASN:ND2	1:A:731:GLY:O	2.53	0.42
2:B:260:ASN:O	2:B:284:GLN:NE2	2.47	0.41
1:A:218:VAL:HG11	1:A:237:LEU:HD13	2.02	0.41
2:B:430:GLN:O	2:B:432:ARG:HG2	2.21	0.41
1:A:19:GLY:HA3	1:A:590:ALA:O	2.20	0.41
1:A:447:SER:HB2	1:A:456:THR:HG21	2.02	0.41
1:A:598:ARG:HE	1:A:733:LEU:HD22	1.86	0.41
2:B:51:CYS:HB3	2:B:56:ILE:HG13	2.03	0.41
2:B:332:HIS:HA	2:B:335:LYS:HB3	2.03	0.41
1:A:291:PRO:HA	1:A:292:PRO:HD3	1.91	0.40
2:B:399:ILE:H	2:B:399:ILE:HG13	1.73	0.40
2:B:407:ARG:HD3	2:B:409:LEU:HD23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	749/763 (98%)	711 (95%)	37 (5%)	1 (0%)	51 83
2	B	457/473 (97%)	439 (96%)	17 (4%)	1 (0%)	47 79
3	C	119/129 (92%)	117 (98%)	2 (2%)	0	100 100
All	All	1325/1365 (97%)	1267 (96%)	56 (4%)	2 (0%)	47 79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	368	VAL
1	A	650	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	634/644 (98%)	619 (98%)	15 (2%)	49	77
2	B	400/412 (97%)	390 (98%)	10 (2%)	47	77
3	C	95/103 (92%)	94 (99%)	1 (1%)	73	88
All	All	1129/1159 (97%)	1103 (98%)	26 (2%)	50	78

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	TYR
1	A	115	ARG
1	A	176	TYR
1	A	208	ARG
1	A	366	THR
1	A	390	TYR
1	A	419	PHE
1	A	440	TYR
1	A	466	TYR
1	A	538	TYR
1	A	540	PHE
1	A	676	PHE
1	A	720	PHE
1	A	749	PHE
1	A	751	PHE
2	B	103	TYR
2	B	109	TYR
2	B	120	ASP
2	B	149	PHE
2	B	247	PHE
2	B	308	TYR
2	B	333	LEU
2	B	360	TYR
2	B	368	VAL
2	B	407	ARG
3	C	107	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	HIS
1	A	210	HIS
1	A	336	GLN
1	A	465	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	1	1,4	14,14,15	0.59	0	17,19,21	1.08	1 (5%)
4	NAG	D	2	4	14,14,15	0.37	0	17,19,21	0.65	1 (5%)
4	NAG	E	1	1,4	14,14,15	0.63	0	17,19,21	0.89	1 (5%)
4	NAG	E	2	4	14,14,15	1.05	2 (14%)	17,19,21	0.58	1 (5%)
4	NAG	F	1	1,4	14,14,15	0.53	0	17,19,21	0.72	1 (5%)
4	NAG	F	2	4	14,14,15	0.32	0	17,19,21	0.68	1 (5%)
4	NAG	G	1	2,4	14,14,15	0.89	1 (7%)	17,19,21	1.20	1 (5%)
4	NAG	G	2	4	14,14,15	0.25	0	17,19,21	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	D	2	4	-	4/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	E	2	4	-	1/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	0/6/23/26	0/1/1/1
4	NAG	G	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	1	NAG	O5-C1	-3.11	1.38	1.43
4	E	2	NAG	C1-C2	2.95	1.56	1.52
4	E	2	NAG	O5-C1	2.52	1.47	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	C2-N2-C7	3.25	127.53	122.90
4	G	1	NAG	C3-C4-C5	2.70	115.06	110.24
4	F	1	NAG	C1-O5-C5	2.47	115.54	112.19
4	E	2	NAG	O5-C1-C2	-2.19	107.83	111.29
4	E	1	NAG	C3-C4-C5	2.13	114.03	110.24
4	D	2	NAG	C1-O5-C5	2.02	114.93	112.19
4	F	2	NAG	C1-O5-C5	2.01	114.91	112.19

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	1	NAG	O5-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6
4	D	2	NAG	C8-C7-N2-C2
4	D	2	NAG	O7-C7-N2-C2
4	G	1	NAG	C8-C7-N2-C2
4	G	1	NAG	O7-C7-N2-C2
4	D	1	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	D	2	NAG	C4-C5-C6-O6
4	E	2	NAG	O5-C5-C6-O6
4	E	1	NAG	C4-C5-C6-O6
4	D	1	NAG	C1-C2-N2-C7
4	D	1	NAG	C3-C2-N2-C7


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	B	501	2	14,14,15	0.67	0	17,19,21	0.72	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	B	501	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	501	NAG	C1-O5-C5	2.60	115.72	112.19

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	501	NAG	O5-C5-C6-O6
6	B	501	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	751/763 (98%)	0.33	49 (6%) 18 11	41, 109, 197, 261	0
2	B	459/473 (97%)	0.46	29 (6%) 20 11	72, 127, 183, 226	0
3	C	121/129 (93%)	1.71	36 (29%) 0 0	98, 172, 226, 251	0
All	All	1331/1365 (97%)	0.50	114 (8%) 10 5	41, 122, 201, 261	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	115	THR	9.2
1	A	708	GLU	8.1
3	C	61	ALA	7.9
1	A	702	GLN	7.7
1	A	628	VAL	7.2
3	C	60	TYR	6.9
3	C	11	GLY	6.9
3	C	12	LEU	6.8
3	C	94	TYR	6.8
1	A	627	VAL	6.0
1	A	631	LYS	6.0
3	C	36	GLY	5.9
1	A	703	LEU	5.8
1	A	296	VAL	5.7
2	B	50	GLY	5.5
2	B	35	PRO	5.1
3	C	50	ALA	5.0
2	B	433	ASP	5.0
2	B	47	LEU	5.0
2	B	438	HIS	4.7
3	C	89	GLU	4.7
1	A	636	VAL	4.5
1	A	287	ILE	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	269	ILE	4.5
1	A	748	LEU	4.4
3	C	43	GLY	4.3
3	C	118	THR	4.3
1	A	701	LEU	4.3
3	C	42	PRO	4.2
1	A	705	ASN	4.2
3	C	96	CYS	4.2
1	A	292	PRO	4.2
1	A	243	GLY	4.2
2	B	389	PHE	4.2
1	A	629	LYS	4.1
1	A	632	GLU	4.0
1	A	630	GLY	4.0
3	C	117	VAL	3.8
1	A	664	ALA	3.8
2	B	36	ASP	3.7
3	C	38	PHE	3.7
1	A	638	VAL	3.6
3	C	85	SER	3.6
1	A	707	ILE	3.5
2	B	368	VAL	3.4
3	C	49	VAL	3.3
3	C	44	LYS	3.3
2	B	51	CYS	3.3
2	B	366	ASN	3.3
1	A	751	PHE	3.3
3	C	14	GLN	3.3
3	C	18	SER	3.2
1	A	623	CYS	3.2
2	B	66	GLN	3.2
1	A	297	PHE	3.2
3	C	48	LEU	3.2
2	B	46	LEU	3.2
1	A	246	PHE	3.1
1	A	699	LEU	3.1
1	A	678	GLU	3.0
1	A	255	VAL	3.0
1	A	679	THR	3.0
3	C	86	LEU	3.0
1	A	131	GLU	3.0
1	A	284	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	454	TYR	3.0
3	C	121	SER	2.9
2	B	99	ARG	2.8
1	A	709	ASP	2.8
3	C	67	ARG	2.8
2	B	228	ILE	2.8
1	A	640	LEU	2.8
3	C	84	ASN	2.8
2	B	369	THR	2.7
1	A	704	PRO	2.7
3	C	41	ALA	2.7
1	A	613	PRO	2.7
3	C	119	VAL	2.7
2	B	338	TYR	2.7
3	C	59	ASN	2.7
1	A	612	ASN	2.6
1	A	639	CYS	2.6
1	A	682	SER	2.6
2	B	171	ASN	2.6
1	A	294	ASP	2.5
2	B	28	ASN	2.5
1	A	293	ARG	2.5
2	B	30	THR	2.5
1	A	267	TYR	2.5
3	C	39	ARG	2.5
2	B	231	ARG	2.5
2	B	109	TYR	2.4
2	B	146	PHE	2.4
1	A	240	ILE	2.4
1	A	484	GLN	2.3
1	A	604	ARG	2.3
1	A	668	GLY	2.3
3	C	37	TRP	2.3
2	B	93	PHE	2.3
2	B	289	LEU	2.3
2	B	33	GLY	2.2
3	C	120	SER	2.2
1	A	214	GLY	2.2
2	B	37	SER	2.2
1	A	675	VAL	2.2
3	C	108	TRP	2.1
1	A	652	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	443	LEU	2.1
3	C	8	THR	2.1
3	C	62	ASP	2.0
2	B	233	VAL	2.0
1	A	645	SER	2.0
3	C	51	ALA	2.0
3	C	111	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	G	2	14/15	0.80	0.22	131,169,207,216	0
4	NAG	E	2	14/15	0.85	0.21	134,163,175,182	0
4	NAG	F	2	14/15	0.86	0.18	167,177,189,191	0
4	NAG	F	1	14/15	0.87	0.18	99,139,178,196	0
4	NAG	G	1	14/15	0.88	0.19	100,127,155,157	0
4	NAG	D	2	14/15	0.88	0.17	104,128,146,154	0
4	NAG	E	1	14/15	0.90	0.23	110,140,161,176	0
4	NAG	D	1	14/15	0.90	0.19	80,104,127,143	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	NAG	B	501	14/15	0.74	0.30	172,188,204,213	0
5	CA	A	2002	1/1	0.75	0.20	78,78,78,78	0
5	CA	A	2001	1/1	0.93	0.15	100,100,100,100	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.