



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 13, 2022 – 12:07 pm GMT

PDB ID : 7QHJ
Title : Peptide GAKSAA in complex with human cathepsin V C25A mutant
Authors : Loboda, J.; Sosnowski, P.; Tusar, L.; Vidmar, R.; Vizovisek, M.; Horvat, J.;
Kosec, G.; Impens, F.; Demol, H.; Turk, B.; Gevaert, K.; Turk, D.
Deposited on : 2021-12-13
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.3
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AA	221	 91% 9%
1	BA	221	 6% 93% 6%
2	PA	8	 12% 38% 12% 50%
2	PB	8	 25% 50% 50%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8520 atoms, of which 4388 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cathepsin L2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	AA	221	3476	1114	1707	305	340	10	1710	10	0
1	BA	221	3355	1078	1645	294	328	10	1647	3	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AA	25	ALA	CYS	engineered mutation	UNP O60911
AA	108	GLN	ASN	engineered mutation	UNP O60911
AA	179	GLN	ASN	engineered mutation	UNP O60911
BA	246	ALA	CYS	engineered mutation	UNP O60911
BA	329	GLN	ASN	engineered mutation	UNP O60911
BA	400	GLN	ASN	engineered mutation	UNP O60911

- Molecule 2 is a protein called SER-ALA-ALA-AMI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	PA	4	34	9	17	4	4	17	0	1
2	PB	4	34	9	17	4	4	17	0	1

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AA	1	Total C O 8 6 2	0	0
3	AA	1	Total C O 8 6 2	0	0
3	AA	1	Total C O 8 6 2	0	0
3	AA	1	Total C O 8 6 2	0	0
3	BA	1	Total C O 8 6 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	AA	1	Total C O 6 3 3	0	0
4	BA	1	Total C O 6 3 3	0	0
4	BA	1	Total C O 6 3 3	0	0
4	BA	1	Total C O 6 3 3	0	0
4	BA	1	Total C O 6 3 3	0	0
4	BA	1	Total C O 6 3 3	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	AA	4	Total Cl 4 4	0	0
5	BA	1	Total Cl 1 1	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AA	1	Total Na 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	AA	256	Total 768	H 512	O 256	512	0
7	BA	237	Total 711	H 474	O 237	474	0
7	PA	5	Total 15	H 10	O 5	10	0
7	PB	3	Total 9	H 6	O 3	6	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cathepsin L2




- Molecule 1: Cathepsin L2



- Molecule 2: SER-ALA-ALA-AMI



- Molecule 2: SER-ALA-ALA-AMI



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.31Å 94.31Å 125.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.67 – 1.40 47.16 – 1.35	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.67-1.40) 100.0 (47.16-1.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.35Å)	Xtriage
Refinement program	MAIN	Depositor
R, R_{free}	0.186 , 0.213 0.192 , 0.192	Depositor DCC
R_{free} test set	2097 reflections (1.69%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8520	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NH2, GOL, CL, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AA	0.73	0/1816	0.79	0/2451
1	BA	0.69	0/1756	0.79	0/2373
2	PA	0.44	0/15	0.57	0/19
2	PB	0.72	0/15	0.88	0/19
All	All	0.70	0/3602	0.79	0/4862

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AA	1769	1707	1694	11	0
1	BA	1710	1645	1638	10	0
2	PA	17	17	14	2	0
2	PB	17	17	14	0	0
3	AA	32	0	56	5	0
3	BA	8	0	14	5	0
4	AA	42	0	55	1	0
4	BA	30	0	40	3	0
5	AA	4	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	BA	1	0	0	0	0
6	AA	1	0	0	0	0
7	AA	256	512	0	3	3
7	BA	237	474	0	2	2
7	PA	5	10	0	2	0
7	PB	3	6	0	0	0
All	All	4132	4388	3525	31	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:BA:504:MPD:H12	7:BA:712:HOH:O	1.65	0.96
3:BA:504:MPD:H53	3:BA:504:MPD:H11	1.52	0.91
3:BA:504:MPD:H52	3:BA:504:MPD:HM3	1.61	0.80
3:BA:504:MPD:HM3	3:BA:504:MPD:C5	2.16	0.75
1:BA:384:ASP:OD2	3:BA:504:MPD:H51	1.89	0.73
3:AA:311:MPD:H12	3:AA:311:MPD:H52	1.83	0.59
1:BA:369:LYS:HE3	7:BA:679:HOH:O	2.03	0.57
1:AA:8:ARG:HB3	4:BA:505:GOL:H12	1.87	0.56
1:BA:320:LYS:O	4:BA:503:GOL:H31	2.07	0.54
4:AA:313:GOL:H11	7:AA:502:HOH:O	2.09	0.53
2:PA:443:SER:N	7:PA:501:HOH:O	2.42	0.52
1:BA:398:ASN:ND2	1:BA:400:GLN:H	2.08	0.51
1:AA:201[A]:LYS:NZ	5:AA:314:CL:CL	2.82	0.50
1:BA:239:ASN:ND2	1:BA:241:LYS:H	2.10	0.49
1:AA:7:TRP:CE2	1:AA:131:GLY:HA2	2.47	0.49
1:AA:141:HIS:HD2	7:AA:422:HOH:O	1.96	0.47
1:BA:228:TRP:CE2	1:BA:352:GLY:HA2	2.50	0.46
1:AA:116:ALA:HB3	1:AA:119:LYS:HD2	1.96	0.46
1:BA:398:ASN:HD22	1:BA:399:SER:N	2.14	0.46
1:AA:163:ASP:CG	3:AA:301:MPD:H13	2.36	0.46
1:AA:7:TRP:CZ3	1:AA:10:LYS:HE2	2.53	0.43
7:AA:411:HOH:O	4:BA:505:GOL:H32	2.18	0.43
1:BA:239:ASN:HD22	1:BA:241:LYS:H	1.66	0.42
1:BA:398:ASN:HD22	1:BA:400:GLN:H	1.68	0.42
1:AA:116:ALA:HA	1:AA:117[B]:PRO:HD2	1.94	0.42
3:AA:301:MPD:H4	3:AA:301:MPD:H11	1.85	0.42
1:BA:224:LYS:N	1:BA:224:LYS:HD3	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:PA:443:SER:N	7:PA:502:HOH:O	2.53	0.41
1:AA:140:GLY:HA2	3:AA:301:MPD:H4	2.02	0.41
1:AA:186:VAL:HG21	1:AA:202:ILE:HD12	2.03	0.41
1:AA:140:GLY:HA2	3:AA:301:MPD:H11	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:AA:608:HOH:O	7:BA:775:HOH:O[6_555]	1.85	0.35
7:AA:538:HOH:O	7:BA:756:HOH:O[4_445]	2.10	0.10
7:AA:516:HOH:O	7:AA:577:HOH:O[3_554]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AA	229/221 (104%)	221 (96%)	8 (4%)	0	100	100
1	BA	222/221 (100%)	214 (96%)	8 (4%)	0	100	100
2	PA	2/8 (25%)	2 (100%)	0	0	100	100
2	PB	2/8 (25%)	2 (100%)	0	0	100	100
All	All	455/458 (99%)	439 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AA	188/178 (106%)	181 (96%)	7 (4%)	34	7
1	BA	181/178 (102%)	173 (96%)	8 (4%)	28	4
2	PA	1/2 (50%)	1 (100%)	0	100	100
2	PB	1/2 (50%)	1 (100%)	0	100	100
All	All	371/360 (103%)	356 (96%)	15 (4%)	35	5

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AA	52	ASN
1	AA	87	GLU
1	AA	89	TYR
1	AA	157	CYS
1	AA	196	SER
1	AA	212[A]	ILE
1	AA	212[B]	ILE
1	BA	239	ASN
1	BA	273	ASN
1	BA	310	TYR
1	BA	316	ASP
1	BA	378	CYS
1	BA	398	ASN
1	BA	433[A]	ILE
1	BA	433[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	BA	239	ASN
1	BA	398	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 6 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	AA	305	-	5,5,5	0.64	0	5,5,5	0.24	0
4	GOL	BA	502	-	5,5,5	0.34	0	5,5,5	0.36	0
3	MPD	AA	311	-	7,7,7	0.25	0	9,10,10	0.39	0
4	GOL	AA	309	-	5,5,5	0.41	0	5,5,5	0.47	0
3	MPD	AA	303	-	7,7,7	0.32	0	9,10,10	0.34	0
4	GOL	AA	312	-	5,5,5	0.17	0	5,5,5	0.26	0
4	GOL	BA	506	-	5,5,5	0.22	0	5,5,5	0.21	0
4	GOL	BA	503	-	5,5,5	0.47	0	5,5,5	0.73	0
4	GOL	AA	308	-	5,5,5	0.15	0	5,5,5	0.25	0
3	MPD	BA	504	-	7,7,7	0.26	0	9,10,10	0.25	0
4	GOL	AA	304	-	5,5,5	0.38	0	5,5,5	0.53	0
4	GOL	BA	507	-	5,5,5	0.25	0	5,5,5	0.27	0
3	MPD	AA	302	-	7,7,7	0.25	0	9,10,10	0.51	0
4	GOL	AA	310	-	5,5,5	0.34	0	5,5,5	0.51	0
3	MPD	AA	301	-	7,7,7	0.46	0	9,10,10	0.47	0
4	GOL	AA	313	-	5,5,5	0.21	0	5,5,5	0.41	0
4	GOL	BA	505	-	5,5,5	0.70	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	AA	305	-	-	0/4/4/4	-
4	GOL	BA	502	-	-	0/4/4/4	-
3	MPD	AA	311	-	-	1/5/5/5	-
4	GOL	AA	309	-	-	2/4/4/4	-
3	MPD	AA	303	-	-	0/5/5/5	-
4	GOL	AA	312	-	-	2/4/4/4	-
4	GOL	BA	506	-	-	0/4/4/4	-
4	GOL	BA	503	-	-	2/4/4/4	-
4	GOL	AA	308	-	-	0/4/4/4	-
3	MPD	BA	504	-	-	2/5/5/5	-
4	GOL	AA	304	-	-	0/4/4/4	-
4	GOL	BA	507	-	-	0/4/4/4	-
3	MPD	AA	302	-	-	3/5/5/5	-
4	GOL	AA	310	-	-	1/4/4/4	-
3	MPD	AA	301	-	-	2/5/5/5	-
4	GOL	AA	313	-	-	0/4/4/4	-
4	GOL	BA	505	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	AA	301	MPD	C2-C3-C4-O4
3	AA	302	MPD	C1-C2-C3-C4
3	AA	302	MPD	O2-C2-C3-C4
3	AA	311	MPD	C2-C3-C4-O4
3	BA	504	MPD	C2-C3-C4-C5
4	BA	503	GOL	O1-C1-C2-C3
4	AA	310	GOL	C1-C2-C3-O3
4	AA	312	GOL	O1-C1-C2-C3
4	BA	503	GOL	O1-C1-C2-O2
4	AA	312	GOL	O1-C1-C2-O2
3	AA	302	MPD	CM-C2-C3-C4
4	AA	309	GOL	O1-C1-C2-C3
4	AA	309	GOL	O1-C1-C2-O2
4	BA	505	GOL	O1-C1-C2-C3
4	BA	505	GOL	C1-C2-C3-O3
3	AA	301	MPD	C2-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	BA	504	MPD	C2-C3-C4-O4

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AA	311	MPD	1	0
4	BA	503	GOL	1	0
3	BA	504	MPD	5	0
3	AA	301	MPD	4	0
4	AA	313	GOL	1	0
4	BA	505	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AA	221/221 (100%)	0.10	2 (0%) 84 82	12, 17, 29, 48	4 (1%)
1	BA	221/221 (100%)	0.50	14 (6%) 20 18	15, 23, 41, 64	1 (0%)
2	PA	3/8 (37%)	1.21	1 (33%) 0 0	23, 23, 27, 35	0
2	PB	3/8 (37%)	1.80	2 (66%) 0 0	30, 30, 39, 62	0
All	All	448/458 (97%)	0.31	19 (4%) 36 37	12, 19, 38, 64	5 (1%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BA	396	GLY	5.6
1	AA	87	GLU	4.1
1	BA	401	ASN	4.0
1	BA	398	ASN	3.7
1	BA	428	ASN	3.0
1	BA	400	GLN	3.0
1	BA	397	ALA	2.8
1	AA	116	ALA	2.7
2	PA	443	SER	2.7
1	BA	338	PRO	2.7
1	BA	399	SER	2.6
1	BA	222	LEU	2.5
1	BA	377	ASP	2.3
1	BA	337	ALA	2.1
1	BA	349	ALA	2.1
1	BA	378	CYS	2.1
1	BA	442	VAL	2.0
2	PB	448	ALA	2.0
2	PB	449	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	AA	302	8/8	0.07	0.36	51,59,67,69	0
4	GOL	AA	310	6/6	0.53	0.27	58,76,79,83	0
4	GOL	AA	312	6/6	0.54	0.17	56,70,75,80	0
3	MPD	AA	311	8/8	0.55	0.27	66,70,72,73	0
4	GOL	BA	506	6/6	0.63	0.23	72,77,89,94	0
4	GOL	BA	507	6/6	0.66	0.21	42,62,76,82	0
4	GOL	AA	308	6/6	0.67	0.27	70,71,76,77	0
3	MPD	AA	301	8/8	0.68	0.19	68,70,73,73	0
3	MPD	AA	303	8/8	0.70	0.17	27,30,34,35	0
4	GOL	AA	313	6/6	0.70	0.21	37,61,72,74	0
4	GOL	BA	505	6/6	0.77	0.18	37,67,69,71	0
4	GOL	AA	309	6/6	0.78	0.17	30,49,76,80	0
4	GOL	AA	304	6/6	0.84	0.12	21,25,36,39	0
3	MPD	BA	504	8/8	0.84	0.14	37,45,49,51	0
4	GOL	BA	503	6/6	0.84	0.12	33,38,45,45	0
4	GOL	BA	502	6/6	0.92	0.11	24,37,54,62	0
4	GOL	AA	305	6/6	0.93	0.18	23,42,62,63	0
5	CL	AA	307	1/1	0.94	0.09	25,25,25,25	0
6	NA	AA	316	1/1	0.94	0.08	17,17,17,17	0
5	CL	BA	501	1/1	0.96	0.06	28,28,28,28	0
5	CL	AA	315	1/1	0.97	0.05	26,26,26,26	0
5	CL	AA	306	1/1	0.98	0.05	22,22,22,22	0
5	CL	AA	314	1/1	0.99	0.04	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.