



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 2, 2022 – 02:38 pm BST

PDB ID : 7R0U
Title : Structure of a cytosolic sulfotransferase of *Anopheles gambiae* (AGAP001425) in complex with 3'-phosphoadenosine 5-phosphate and vanillin.
Authors : Esposito Verza, A.; Miggiano, R.; Rizzi, R.; Rossi, F.
Deposited on : 2022-02-02
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.29
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	360	83% 10% 7%
1	B	360	81% 12% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	405	-	-	X	-

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AGAP001425-PA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2821	1811	477	522	11	0	0	0
1	B	336	2821	1811	477	522	11	0	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP Q7PXJ0
A	-19	GLY	-	expression tag	UNP Q7PXJ0
A	-18	HIS	-	expression tag	UNP Q7PXJ0
A	-17	HIS	-	expression tag	UNP Q7PXJ0
A	-16	HIS	-	expression tag	UNP Q7PXJ0
A	-15	HIS	-	expression tag	UNP Q7PXJ0
A	-14	HIS	-	expression tag	UNP Q7PXJ0
A	-13	HIS	-	expression tag	UNP Q7PXJ0
A	-12	HIS	-	expression tag	UNP Q7PXJ0
A	-11	HIS	-	expression tag	UNP Q7PXJ0
A	-10	HIS	-	expression tag	UNP Q7PXJ0
A	-9	HIS	-	expression tag	UNP Q7PXJ0
A	-8	SER	-	expression tag	UNP Q7PXJ0
A	-7	SER	-	expression tag	UNP Q7PXJ0
A	-6	GLY	-	expression tag	UNP Q7PXJ0
A	-5	HIS	-	expression tag	UNP Q7PXJ0
A	-4	ILE	-	expression tag	UNP Q7PXJ0
A	-3	GLU	-	expression tag	UNP Q7PXJ0
A	-2	GLY	-	expression tag	UNP Q7PXJ0
A	-1	ARG	-	expression tag	UNP Q7PXJ0
A	0	HIS	-	expression tag	UNP Q7PXJ0
B	-20	MET	-	initiating methionine	UNP Q7PXJ0
B	-19	GLY	-	expression tag	UNP Q7PXJ0
B	-18	HIS	-	expression tag	UNP Q7PXJ0
B	-17	HIS	-	expression tag	UNP Q7PXJ0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP Q7PXJ0
B	-15	HIS	-	expression tag	UNP Q7PXJ0
B	-14	HIS	-	expression tag	UNP Q7PXJ0
B	-13	HIS	-	expression tag	UNP Q7PXJ0
B	-12	HIS	-	expression tag	UNP Q7PXJ0
B	-11	HIS	-	expression tag	UNP Q7PXJ0
B	-10	HIS	-	expression tag	UNP Q7PXJ0
B	-9	HIS	-	expression tag	UNP Q7PXJ0
B	-8	SER	-	expression tag	UNP Q7PXJ0
B	-7	SER	-	expression tag	UNP Q7PXJ0
B	-6	GLY	-	expression tag	UNP Q7PXJ0
B	-5	HIS	-	expression tag	UNP Q7PXJ0
B	-4	ILE	-	expression tag	UNP Q7PXJ0
B	-3	GLU	-	expression tag	UNP Q7PXJ0
B	-2	GLY	-	expression tag	UNP Q7PXJ0
B	-1	ARG	-	expression tag	UNP Q7PXJ0
B	0	HIS	-	expression tag	UNP Q7PXJ0

- Molecule 2 is 4-hydroxy-3-methoxybenzaldehyde (three-letter code: V55) (formula: C₈H₈O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	8	3		
2	B	1	Total	C	O	0	0
			11	8	3		

- Molecule 3 is ADENOSINE-3'-5'-DIPHOSPHATE (three-letter code: A3P) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	37	Total	O	0	0
			37	37		
5	B	47	Total	O	0	0
			47	47		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: AGAP001425-PA

Chain A: 83% 10% 7%



- Molecule 1: AGAP001425-PA

Chain B: 81% 12% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.24Å 91.82Å 103.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.40 – 2.50 62.40 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.9 (62.40-2.50) 98.9 (62.40-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.13 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.206 , 0.252 0.216 , 0.209	Depositor DCC
R_{free} test set	1275 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtrriage
Anisotropy	0.643	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5820	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.75 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1867e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: V55, A3P, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2903	0.86	0/3927
1	B	0.73	1/2903 (0.0%)	0.85	0/3927
All	All	0.72	1/5806 (0.0%)	0.85	0/7854

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	102	GLU	CD-OE2	5.18	1.31	1.25

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	297	GLU	Peptide
1	B	297	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2821	0	2699	21	0
1	B	2821	0	2699	26	0
2	A	11	0	8	0	0
2	B	11	0	7	0	0
3	A	27	0	11	1	0
3	B	27	0	11	0	0
4	B	18	0	24	4	0
5	A	37	0	0	2	0
5	B	47	0	0	3	0
All	All	5820	0	5459	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:215:PRO:HB3	4:B:405:GOL:H31	1.24	1.13
1:B:215:PRO:CB	4:B:405:GOL:H31	2.07	0.81
1:A:67:ARG:HD3	1:A:186:LEU:HD13	1.67	0.77
1:B:131:GLN:HG2	5:B:530:HOH:O	1.89	0.71
1:B:36:ARG:HH21	1:B:36:ARG:HG3	1.62	0.65
1:B:258:SER:HA	5:B:543:HOH:O	2.00	0.60
1:A:72:TRP:HH2	1:A:241:LYS:HA	1.68	0.59
1:A:244:LEU:HB3	1:A:245:PRO:HD3	1.87	0.57
1:A:260:SER:O	1:A:264:ILE:HD12	2.04	0.56
1:B:72:TRP:HH2	1:B:241:LYS:HA	1.70	0.56
1:A:67:ARG:HD3	1:A:186:LEU:CD1	2.35	0.56
1:B:288:LYS:HA	5:B:501:HOH:O	2.07	0.54
1:A:84:LEU:CD1	1:A:263:GLN:HB3	2.42	0.50
1:A:63:VAL:O	1:A:171:TYR:HA	2.11	0.49
1:B:63:VAL:O	1:B:171:TYR:HA	2.13	0.49
1:B:217:TRP:HB2	4:B:405:GOL:H32	1.94	0.48
1:A:161:SER:HA	1:A:164:GLU:HB2	1.96	0.48
3:A:402:A3P:H8	5:A:510:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:86:PHE:CZ	1:B:266:THR:HG22	2.49	0.47
1:B:298:GLN:HG2	1:B:302:ARG:HG3	1.97	0.47
1:A:176:PRO:HG3	1:A:217:TRP:CH2	2.51	0.46
1:B:84:LEU:CD1	1:B:263:GLN:HB3	2.46	0.46
1:A:86:PHE:CZ	1:A:266:THR:HG22	2.52	0.45
1:A:112:LYS:HD3	1:A:133:SER:OG	2.17	0.45
1:B:100:PHE:CD2	1:B:103:PHE:HB2	2.51	0.45
1:B:176:PRO:HG3	1:B:217:TRP:CH2	2.52	0.45
1:A:84:LEU:HD11	1:A:263:GLN:HB3	1.99	0.44
1:B:152:HIS:HB2	1:B:213:TRP:CE2	2.52	0.44
1:B:161:SER:HA	1:B:164:GLU:HB2	1.98	0.44
1:A:65:TYR:CG	1:A:66:PRO:HD2	2.53	0.44
1:A:100:PHE:CD2	1:A:103:PHE:HB2	2.53	0.44
1:A:177:SER:HA	1:A:321:LEU:HD21	2.00	0.43
1:B:35:LYS:HD2	1:B:218:GLU:HG2	1.99	0.43
1:B:65:TYR:CG	1:B:66:PRO:HD2	2.53	0.43
1:B:298:GLN:HG2	1:B:302:ARG:CG	2.49	0.43
1:B:23:HIS:CE1	1:B:196:VAL:HG11	2.53	0.43
1:B:215:PRO:HB3	4:B:405:GOL:C3	2.17	0.43
1:A:23:HIS:CE1	1:A:196:VAL:HG11	2.54	0.42
1:B:84:LEU:HD11	1:B:263:GLN:HB3	2.02	0.42
1:A:152:HIS:HB2	1:A:213:TRP:CE2	2.54	0.42
1:A:176:PRO:HG3	1:A:217:TRP:CZ2	2.55	0.42
1:B:244:LEU:HD23	1:B:244:LEU:HA	1.82	0.42
1:B:32:VAL:HG21	1:B:43:PHE:CE1	2.55	0.42
1:A:244:LEU:HD12	1:A:244:LEU:HA	1.85	0.41
1:A:232:VAL:O	5:A:501:HOH:O	2.22	0.41
1:B:333:ASP:OD1	1:B:333:ASP:N	2.54	0.41
1:A:336:PHE:HA	1:A:337:PRO:HD3	1.93	0.40
1:B:120:GLU:HA	1:B:126:MET:SD	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/360 (93%)	326 (98%)	8 (2%)	0	100	100
1	B	334/360 (93%)	325 (97%)	9 (3%)	0	100	100
All	All	668/720 (93%)	651 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/328 (94%)	302 (98%)	5 (2%)	62	84
1	B	307/328 (94%)	296 (96%)	11 (4%)	35	61
All	All	614/656 (94%)	598 (97%)	16 (3%)	46	72

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	PHE
1	A	144	LYS
1	A	311	GLU
1	A	321	LEU
1	A	322	ASN
1	B	34	GLU
1	B	36	ARG
1	B	90	LYS
1	B	98	PHE
1	B	133	SER
1	B	229	ARG
1	B	230	GLU
1	B	265	ASP
1	B	298	GLN
1	B	318	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	321	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	V55	B	402	-	11,11,11	0.41	0	14,14,14	0.53	0
2	V55	A	401	-	11,11,11	0.46	0	14,14,14	0.59	0
4	GOL	B	403	-	5,5,5	0.18	0	5,5,5	0.49	0
3	A3P	A	402	-	26,29,29	1.33	3 (11%)	31,45,45	1.87	10 (32%)
4	GOL	B	404	-	5,5,5	0.21	0	5,5,5	0.49	0
4	GOL	B	405	-	5,5,5	0.26	0	5,5,5	0.69	0
3	A3P	B	401	-	26,29,29	1.21	2 (7%)	31,45,45	1.89	10 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	V55	B	402	-	-	2/4/4/4	0/1/1/1
2	V55	A	401	-	-	0/4/4/4	0/1/1/1
4	GOL	B	403	-	-	2/4/4/4	-
3	A3P	A	402	-	-	3/11/31/31	0/3/3/3
4	GOL	B	404	-	-	4/4/4/4	-
4	GOL	B	405	-	-	4/4/4/4	-
3	A3P	B	401	-	-	2/11/31/31	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	A3P	C2-N3	2.90	1.36	1.32
3	A	402	A3P	C5-C4	2.88	1.48	1.40
3	B	401	A3P	C5-C4	2.78	1.48	1.40
3	B	401	A3P	C2-N3	2.35	1.35	1.32
3	A	402	A3P	O5'-C5'	-2.09	1.36	1.44

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	A3P	N3-C2-N1	-4.97	120.90	128.68
3	B	401	A3P	O4'-C1'-C2'	-3.68	101.55	106.93
3	A	402	A3P	N3-C2-N1	-3.56	123.11	128.68
3	A	402	A3P	O5'-P2-O4P	-3.38	96.98	106.47
3	A	402	A3P	O2P-P1-O1P	3.25	123.39	110.68
3	A	402	A3P	O3'-P1-O1P	-2.91	98.17	109.39
3	A	402	A3P	O4'-C4'-C3'	2.90	111.09	104.87
3	A	402	A3P	O4'-C1'-C2'	-2.89	102.70	106.93
3	B	401	A3P	O5'-P2-O4P	-2.71	98.86	106.47
3	A	402	A3P	N6-C6-N1	2.67	124.11	118.57
3	B	401	A3P	O2P-P1-O1P	2.66	121.10	110.68
3	B	401	A3P	O5P-P2-O5'	-2.65	99.68	106.73
3	A	402	A3P	C2'-C3'-C4'	-2.65	98.54	103.22
3	B	401	A3P	O5P-P2-O4P	2.62	120.95	110.68
3	B	401	A3P	C2-N1-C6	2.62	123.23	118.75
3	B	401	A3P	C2'-C3'-C4'	-2.55	98.71	103.22
3	B	401	A3P	N6-C6-N1	2.48	123.72	118.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	A3P	O3'-P1-O1P	-2.35	100.32	109.39
3	A	402	A3P	O4'-C4'-C5'	-2.10	102.48	109.37
3	A	402	A3P	C5'-C4'-C3'	-2.05	107.61	114.40

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	A3P	C3'-O3'-P1-O3P
4	B	403	GOL	O1-C1-C2-C3
4	B	404	GOL	O1-C1-C2-C3
4	B	404	GOL	C1-C2-C3-O3
4	B	404	GOL	O2-C2-C3-O3
4	B	405	GOL	O1-C1-C2-C3
2	B	402	V55	OAB-CAD-CAI-CAG
2	B	402	V55	OAB-CAD-CAI-CAE
4	B	403	GOL	O1-C1-C2-O2
3	A	402	A3P	C4'-C3'-O3'-P1
3	A	402	A3P	C2'-C3'-O3'-P1
4	B	404	GOL	O1-C1-C2-O2
4	B	405	GOL	O1-C1-C2-O2
4	B	405	GOL	O2-C2-C3-O3
3	A	402	A3P	C5'-O5'-P2-O4P
3	B	401	A3P	C4'-C3'-O3'-P1
4	B	405	GOL	C1-C2-C3-O3



There are no ring outliers.



2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	A3P	1	0
4	B	405	GOL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.



6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around V55 A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around V55 B 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A3P A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.