



Full wwPDB X-ray Structure Validation Report i

Oct 14, 2023 – 09:05 PM EDT

PDB ID : 7T7U
Title : Light Harvesting complex phycocyanin PC 630, from the cryptophyte Chroomonas sp. M1627
Authors : Michie, K.A.; Harrop, S.J.; Rathbone, H.W.; Wilk, K.E.; Curmi, P.M.G.
Deposited on : 2021-12-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 8530 atoms, of which 4048 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phycoerythrin alpha subunit L1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	81	1349	414	694	109	127	5	0	6	0


- Molecule 2 is a protein called Phycoerythrin beta subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	162	2415	744	1210	206	245	10	0	7	0
2	D	168	2526	780	1267	217	252	10	0	6	0

- Molecule 3 is a protein called Phycoerythrin alpha subunit S1.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	C	70	1140	345	579	96	113	7	0	5	0

- Molecule 4 is mesobiliverdin IX(alpha) (three-letter code: M1V) (formula: C₃₃H₃₈N₄O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
4	A	1	78	33	35	4	6	0	0
4	C	1	78	33	35	4	6	0	0

- Molecule 5 is DiCys-(15,16)-Dihydrobiliverdin (three-letter code: AX9) (formula: C₃₃H₄₀N₄O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
5	B	1	79	33	36	4	6	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
5	D	1	79	33	36	4	6	0	0

- Molecule 6 is PHYCOCYANOBILIN (three-letter code: CYC) (formula: C₃₃H₄₀N₄O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
6	B	1	80	33	37	4	6	0	0
6	B	1	80	33	37	4	6	0	0
6	D	1	80	33	37	4	6	0	0
6	D	1	80	33	37	4	6	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C H O 14 3 8 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	98	Total O 98 98	0	0
8	B	122	Total O 122 122	0	0
8	C	85	Total O 85 85	0	0
8	D	147	Total O 147 147	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phycoerythrin alpha subunit L1




- Molecule 2: Phycoerythrin beta subunit



- Molecule 2: Phycoerythrin beta subunit



- Molecule 3: Phycoerythrin alpha subunit S1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	90.29 Å 93.41 Å 132.02 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.58 – 1.80 39.58 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (39.58-1.80) 98.5 (39.58-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.96 (at 1.79 Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R , R_{free}	0.158 , 0.185 0.158 , 0.184	Depositor DCC
R_{free} test set	2622 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 44.2	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.001 for -1/2*h+1/2*k+1/2*l, 1/2*h-1/2*k +1/2*l, h+k 0.014 for -1/2*h+1/2*k-1/2*l, 1/2*h-1/2*k- 1/2*l, -h-k 0.017 for k, h, -l 0.013 for -1/2*h-1/2*k+1/2*l, -1/2*h-1/2*k- 1/2*l, h-k 0.013 for -1/2*h-1/2*k-1/2*l, -1/2*h-1/2*k+ 1/2*l, -h+k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8530	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: LYZ, MEN, AX9, CYC, GOL, M1V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/663	0.61	0/883
2	B	0.56	0/1224	0.63	0/1652
2	D	0.59	0/1273	0.63	0/1716
3	C	0.61	0/559	0.64	0/740
All	All	0.57	0/3719	0.63	0/4991

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	655	694	703	3	0
2	B	1205	1210	1205	5	0
2	D	1259	1267	1271	3	0
3	C	561	579	579	0	0
4	A	43	35	35	1	0
4	C	43	35	35	2	0
5	B	43	36	0	0	0
5	D	43	36	0	0	0
6	B	86	74	74	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	86	74	74	4	0
7	B	6	8	8	0	0
8	A	98	0	0	2	0
8	B	122	0	0	0	0
8	C	85	0	0	0	0
8	D	147	0	0	0	0
All	All	4482	4048	3984	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:101:M1V:H41	4:A:101:M1V:H13	1.58	0.68
4:C:101:M1V:H13	4:C:101:M1V:H41	1.58	0.67
6:B:203:CYC:NB	6:B:203:CYC:HMA1	2.23	0.54
6:D:302:CYC:HMD2	6:D:302:CYC:HC	1.73	0.54
6:B:203:CYC:HMA1	6:B:203:CYC:HB	1.72	0.53
6:D:301:CYC:NB	6:D:301:CYC:HMA1	2.24	0.53
6:D:302:CYC:NB	6:D:302:CYC:HMA1	2.25	0.50
2:B:114:LYS:HA	2:B:114:LYS:HE2	1.95	0.49
6:B:202:CYC:HMA1	6:B:202:CYC:NB	2.28	0.48
6:D:302:CYC:HMA1	6:D:302:CYC:HB	1.78	0.48
2:B:114:LYS:O	2:B:114:LYS:HD3	2.13	0.48
8:A:229:HOH:O	2:D:150:LYS:HE3	2.13	0.47
6:B:203:CYC:HC	6:B:203:CYC:HMD2	1.79	0.47
1:A:34[B]:MET:HE2	1:A:38:GLN:HE22	1.79	0.46
2:B:171:LYS:N	2:B:171:LYS:HD3	2.31	0.46
1:A:61:PHE:CE1	2:B:57:SER:HB2	2.51	0.46
6:B:202:CYC:HBC3	6:B:202:CYC:HHD	1.99	0.45
2:D:77[B]:ARG:NH2	2:D:78:ARG:HE	2.14	0.45
1:A:34[B]:MET:HE3	8:A:267:HOH:O	2.16	0.44
4:C:101:M1V:H13	4:C:101:M1V:NA	2.30	0.43
2:D:26:ALA:O	2:D:29:LYS:HG3	2.19	0.42
2:B:167:GLY:O	2:B:171:LYS:HE2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	84/81 (104%)	79 (94%)	5 (6%)	0	100 100
2	B	166/173 (96%)	164 (99%)	2 (1%)	0	100 100
2	D	169/173 (98%)	167 (99%)	1 (1%)	1 (1%)	25 12
3	C	72/70 (103%)	71 (99%)	1 (1%)	0	100 100
All	All	491/497 (99%)	481 (98%)	9 (2%)	1 (0%)	47 33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	75	THR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	73/67 (109%)	72 (99%)	1 (1%)	67 59
2	B	133/136 (98%)	130 (98%)	3 (2%)	50 37
2	D	138/136 (102%)	137 (99%)	1 (1%)	84 81
3	C	61/56 (109%)	60 (98%)	1 (2%)	62 54
All	All	405/395 (102%)	399 (98%)	6 (2%)	62 56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LYS
2	B	50	CYS
2	B	54	ASP
2	B	114	LYS
3	C	64	SER
2	D	101	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
2	B	148	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LYZ	A	4	1	7,9,10	0.65	0	4,10,12	0.79	0
2	MEN	B	72	2	7,8,9	0.94	0	6,9,11	1.75	2 (33%)
3	LYZ	C	4	3	7,9,10	0.76	0	4,10,12	0.75	0
2	MEN	D	72	2	7,8,9	0.67	0	6,9,11	1.83	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LYZ	A	4	1	-	1/8/9/11	-
2	MEN	B	72	2	-	4/7/8/10	-
3	LYZ	C	4	3	-	0/8/9/11	-
2	MEN	D	72	2	-	2/7/8/10	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	72	MEN	CB-CG-ND2	3.61	120.33	115.48
2	B	72	MEN	CB-CA-C	-3.12	105.62	111.47
2	B	72	MEN	CB-CG-ND2	2.53	118.88	115.48
2	D	72	MEN	CB-CA-C	-2.39	106.98	111.47

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	4	LYZ	CG-CD-CE-NZ
2	B	72	MEN	C-CA-CB-CG
2	B	72	MEN	N-CA-CB-CG
2	B	72	MEN	CA-CB-CG-OD1
2	B	72	MEN	CA-CB-CG-ND2
2	D	72	MEN	CA-CB-CG-OD1
2	D	72	MEN	CA-CB-CG-ND2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	M1V	A	101	1	42,46,46	1.94	10 (23%)	53,67,67	1.37	8 (15%)
5	AX9	B	201	2	41,46,46	1.57	9 (21%)	41,67,67	1.10	3 (7%)
5	AX9	D	303	2	41,46,46	1.56	7 (17%)	41,67,67	1.41	6 (14%)
6	CYC	B	203	2	42,46,46	2.74	14 (33%)	50,67,67	2.05	11 (22%)
4	M1V	C	101	3	42,46,46	2.16	11 (26%)	53,67,67	1.25	6 (11%)
6	CYC	D	302	2	42,46,46	2.40	13 (30%)	50,67,67	2.15	11 (22%)
7	GOL	B	204	-	5,5,5	0.62	0	5,5,5	1.13	0
6	CYC	D	301	2	42,46,46	2.63	11 (26%)	50,67,67	2.35	13 (26%)
6	CYC	B	202	2	42,46,46	2.66	12 (28%)	50,67,67	1.95	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	M1V	A	101	1	-	10/26/74/74	0/4/4/4
5	AX9	B	201	2	-	5/26/74/74	0/4/4/4
5	AX9	D	303	2	-	8/26/74/74	0/4/4/4
6	CYC	B	203	2	-	7/25/74/74	0/4/4/4
4	M1V	C	101	3	-	7/26/74/74	0/4/4/4
6	CYC	D	302	2	-	8/25/74/74	0/4/4/4
7	GOL	B	204	-	-	2/4/4/4	-
6	CYC	D	301	2	-	10/25/74/74	0/4/4/4
6	CYC	B	202	2	-	6/25/74/74	0/4/4/4

All (87) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	301	CYC	C1C-NC	-10.97	1.23	1.37
6	B	203	CYC	C1C-NC	-9.39	1.25	1.37
6	B	202	CYC	C1C-NC	-9.19	1.25	1.37
6	D	302	CYC	C1C-NC	-7.96	1.27	1.37
6	B	202	CYC	C4C-NC	-7.04	1.22	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	203	CYC	C4C-NC	-6.22	1.24	1.37
6	D	302	CYC	CHB-C4A	5.73	1.53	1.40
4	C	101	M1V	CHC-C4C	5.49	1.53	1.40
6	D	301	CYC	C4C-NC	-5.48	1.25	1.37
4	C	101	M1V	CHB-C1C	5.24	1.39	1.35
4	C	101	M1V	C1C-C2C	-5.20	1.37	1.45
6	B	203	CYC	CHB-C4A	5.19	1.52	1.40
6	D	301	CYC	CHB-C4A	5.18	1.52	1.40
6	D	302	CYC	C4C-NC	-5.15	1.26	1.37
4	C	101	M1V	C4D-C3D	-5.00	1.38	1.48
6	B	203	CYC	C4B-C3B	-4.89	1.38	1.48
4	A	101	M1V	C4D-C3D	-4.72	1.39	1.48
4	A	101	M1V	C1C-C2C	-4.71	1.38	1.45
6	B	202	CYC	CAC-C3C	-4.70	1.44	1.54
6	B	203	CYC	CHA-C1A	4.67	1.39	1.35
5	B	201	AX9	C1C-C2C	-4.57	1.38	1.45
6	B	202	CYC	CHB-C4A	4.55	1.51	1.40
4	C	101	M1V	C4C-C3C	-4.54	1.36	1.45
5	D	303	AX9	C1C-C2C	-4.40	1.38	1.45
6	D	302	CYC	C4B-C3B	-4.23	1.40	1.48
6	B	203	CYC	C1B-C2B	-4.22	1.37	1.45
4	A	101	M1V	C4C-C3C	-4.03	1.37	1.45
4	A	101	M1V	CHC-C4C	4.03	1.49	1.40
6	D	301	CYC	C4B-C3B	-3.94	1.40	1.48
6	B	202	CYC	C4B-C3B	-3.94	1.40	1.48
5	D	303	AX9	C1A-C2A	-3.92	1.37	1.47
6	B	202	CYC	CHA-C1A	3.91	1.38	1.35
6	D	301	CYC	C1B-C2B	-3.91	1.38	1.45
4	A	101	M1V	C1D-C2D	-3.80	1.38	1.45
6	B	202	CYC	C2C-C3C	3.80	1.65	1.54
6	B	203	CYC	C4D-CHA	3.66	1.55	1.41
6	D	302	CYC	CAC-C3C	-3.61	1.46	1.54
6	B	203	CYC	C1D-CHD	3.61	1.55	1.41
6	B	202	CYC	C1D-CHD	3.56	1.55	1.41
6	D	301	CYC	CHA-C1A	3.47	1.38	1.35
6	D	301	CYC	C2C-C3C	3.43	1.64	1.54
6	D	301	CYC	C1D-CHD	3.35	1.54	1.41
6	B	203	CYC	C2C-C1C	-3.34	1.49	1.52
4	A	101	M1V	C1A-C2A	-3.30	1.38	1.47
6	D	301	CYC	C4D-CHA	3.26	1.53	1.41
6	D	302	CYC	CHA-C1A	3.25	1.37	1.35
6	B	203	CYC	CAC-C3C	-3.24	1.47	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	101	M1V	C4A-C3A	-3.23	1.38	1.45
6	B	202	CYC	C1B-C2B	-3.23	1.39	1.45
6	D	302	CYC	C1D-CHD	3.23	1.53	1.41
4	A	101	M1V	C4A-C3A	-3.22	1.38	1.45
6	B	202	CYC	C4D-CHA	3.19	1.53	1.41
5	B	201	AX9	C4D-ND	3.14	1.39	1.35
6	B	203	CYC	C2C-C3C	3.10	1.63	1.54
6	D	302	CYC	C1B-C2B	-3.08	1.39	1.45
5	D	303	AX9	CHB-C1C	3.08	1.37	1.35
6	D	302	CYC	C4D-CHA	3.06	1.53	1.41
4	C	101	M1V	C1A-C2A	-3.02	1.39	1.47
6	D	301	CYC	CAC-C3C	-3.01	1.48	1.54
6	D	302	CYC	C2C-C3C	2.94	1.62	1.54
6	B	202	CYC	C2C-C1C	-2.94	1.49	1.52
6	D	302	CYC	C1A-C2A	-2.91	1.41	1.45
6	B	202	CYC	C1A-C2A	-2.84	1.41	1.45
5	B	201	AX9	C4D-C3D	-2.81	1.42	1.48
5	D	303	AX9	C4D-ND	2.80	1.38	1.35
5	B	201	AX9	C1D-ND	-2.78	1.41	1.45
5	B	201	AX9	C1B-CHA	2.70	1.51	1.41
5	B	201	AX9	C4B-CHB	2.65	1.51	1.41
4	A	101	M1V	C4B-CHB	2.63	1.51	1.41
4	A	101	M1V	C1B-CHA	2.60	1.51	1.41
5	D	303	AX9	C4B-CHB	2.55	1.51	1.41
4	C	101	M1V	C4B-CHB	2.53	1.50	1.41
5	B	201	AX9	CHC-C1D	-2.48	1.48	1.53
6	B	203	CYC	C3C-C4C	-2.46	1.47	1.50
5	D	303	AX9	C1B-CHA	2.44	1.50	1.41
4	C	101	M1V	C1B-CHA	2.42	1.50	1.41
6	D	301	CYC	C1A-C2A	-2.35	1.42	1.45
5	B	201	AX9	C1A-C2A	-2.35	1.41	1.47
4	A	101	M1V	CHB-C1C	2.34	1.37	1.35
4	C	101	M1V	C1D-C2D	-2.34	1.40	1.45
5	B	201	AX9	C4A-C3A	-2.24	1.40	1.45
6	D	302	CYC	O1A-CGA	2.17	1.29	1.22
4	C	101	M1V	C1C-NC	-2.17	1.33	1.38
5	D	303	AX9	C4A-C3A	-2.16	1.40	1.45
6	D	302	CYC	C3C-C4C	-2.13	1.47	1.50
6	B	203	CYC	C4A-C3A	-2.06	1.41	1.45
6	B	203	CYC	C1A-C2A	-2.04	1.42	1.45

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	301	CYC	OC-C1C-C2C	-9.03	118.99	126.17
6	D	302	CYC	OC-C1C-C2C	-8.16	119.69	126.17
6	D	301	CYC	C2C-C1C-NC	7.92	115.10	108.27
6	B	202	CYC	C2C-C1C-NC	7.39	114.64	108.27
6	B	203	CYC	C2C-C1C-NC	7.32	114.58	108.27
6	D	302	CYC	C2C-C1C-NC	7.25	114.52	108.27
6	B	202	CYC	OC-C1C-C2C	-7.01	120.60	126.17
6	B	203	CYC	OC-C1C-C2C	-5.47	121.83	126.17
6	D	301	CYC	CAB-C3B-C4B	5.40	129.91	121.38
6	D	302	CYC	CBD-CAD-C3D	-4.70	104.61	112.62
6	B	203	CYC	CAB-C3B-C4B	4.06	127.79	121.38
5	B	201	AX9	CBB-CAB-C3B	-3.48	106.67	112.62
6	B	202	CYC	C4D-CHA-C1A	-3.46	124.68	128.81
6	B	203	CYC	CHB-C4A-C3A	3.27	133.30	124.90
4	A	101	M1V	CAC-CBC-CGC	-3.21	106.69	113.60
5	D	303	AX9	CAA-C3A-C4A	3.17	128.65	124.38
4	A	101	M1V	C4B-CHB-C1C	-3.08	125.13	128.81
6	D	301	CYC	C2C-C3C-C4C	-3.04	96.79	101.34
4	C	101	M1V	OD-C4D-C3D	-3.02	124.77	128.04
6	D	302	CYC	CHB-C4A-C3A	2.98	132.57	124.90
5	D	303	AX9	C4A-NA-C1A	-2.95	106.92	110.67
6	B	203	CYC	CBD-CAD-C3D	-2.94	107.61	112.62
5	D	303	AX9	C4B-CHB-C1C	-2.85	125.40	128.81
5	D	303	AX9	CHC-C1D-ND	-2.84	110.14	113.72
6	D	301	CYC	CAC-C3C-C4C	2.83	119.95	112.67
6	B	203	CYC	O2A-CGA-CBA	2.83	123.11	114.03
4	A	101	M1V	CAA-C3A-C4A	2.80	128.16	124.38
6	B	202	CYC	C2C-C3C-C4C	-2.79	97.16	101.34
4	A	101	M1V	C1D-ND-C4D	-2.70	107.23	110.67
5	D	303	AX9	O1C-CGC-CBC	2.68	122.63	114.03
5	D	303	AX9	CAD-C3D-C4D	2.62	125.52	121.38
6	D	302	CYC	O2A-CGA-CBA	2.57	122.29	114.03
6	B	202	CYC	C3B-C4B-NB	2.57	108.85	106.78
6	D	302	CYC	CHD-C4C-NC	2.52	128.20	125.20
6	D	301	CYC	C1B-NB-C4B	-2.42	107.59	110.67
6	D	301	CYC	CAC-C3C-C2C	-2.42	108.21	114.26
4	A	101	M1V	OD-C4D-C3D	-2.41	125.43	128.04
4	C	101	M1V	C1D-ND-C4D	-2.35	107.68	110.67
6	B	203	CYC	C4D-CHA-C1A	-2.33	126.02	128.81
5	B	201	AX9	CAD-C3D-C4D	2.32	125.05	121.38
6	D	301	CYC	CHB-C4A-C3A	2.31	130.85	124.90
6	D	302	CYC	CAB-C3B-C4B	2.30	125.02	121.38
6	B	203	CYC	C2A-C1A-NA	-2.30	106.70	110.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	203	CYC	C3A-C4A-NA	-2.30	105.61	110.53
6	D	302	CYC	O2D-CGD-CBD	2.28	121.34	114.03
4	C	101	M1V	O1C-CGC-O2C	-2.27	117.64	123.30
6	D	302	CYC	C1B-NB-C4B	-2.25	107.80	110.67
6	D	301	CYC	O2A-CGA-CBA	2.24	121.23	114.03
6	D	301	CYC	CAB-C3B-C2B	-2.23	123.72	127.53
6	B	202	CYC	CMA-C3A-C4A	2.21	128.47	125.06
6	B	202	CYC	O2A-CGA-O1A	-2.20	117.81	123.30
4	A	101	M1V	CAA-C3A-C2A	-2.19	123.79	127.53
6	D	302	CYC	CHB-C1B-C2B	-2.18	122.63	126.95
6	D	302	CYC	CHB-C4A-NA	-2.17	120.40	124.93
6	B	202	CYC	CAB-C3B-C4B	2.15	124.78	121.38
4	A	101	M1V	O2B-CGB-O1B	-2.14	117.96	123.30
6	D	301	CYC	O2D-CGD-CBD	2.14	120.91	114.03
6	D	301	CYC	C2A-C1A-NA	-2.14	106.94	110.05
4	C	101	M1V	C4B-CHB-C1C	-2.10	126.30	128.81
4	C	101	M1V	C1C-NC-C4C	2.10	110.46	106.51
5	B	201	AX9	O2B-CGB-CBB	2.10	120.77	114.03
4	C	101	M1V	O2B-CGB-O1B	-2.07	118.14	123.30
6	B	203	CYC	CMA-C3A-C4A	2.06	128.23	125.06
4	A	101	M1V	C3D-C4D-ND	2.03	108.42	106.78
6	B	203	CYC	C4A-C3A-C2A	2.03	108.84	106.51
6	D	301	CYC	O2A-CGA-O1A	-2.00	118.31	123.30

There are no chirality outliers.

All (63) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	101	M1V	C2B-C1B-CHA-C4A
4	A	101	M1V	NB-C1B-CHA-C4A
4	A	101	M1V	C3C-C4C-CHC-C1D
4	A	101	M1V	NC-C4C-CHC-C1D
4	C	101	M1V	C2B-C1B-CHA-C4A
4	C	101	M1V	NB-C1B-CHA-C4A
4	C	101	M1V	C3C-C4C-CHC-C1D
4	C	101	M1V	NC-C4C-CHC-C1D
5	B	201	AX9	NB-C1B-CHA-C4A
5	B	201	AX9	C2B-C1B-CHA-C4A
5	B	201	AX9	NC-C4C-CHC-C1D
5	D	303	AX9	NB-C1B-CHA-C4A
5	D	303	AX9	C2B-C1B-CHA-C4A
5	D	303	AX9	NC-C4C-CHC-C1D

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	202	CYC	NA-C4A-CHB-C1B
6	B	202	CYC	C2C-C3C-CAC-CBC
6	B	202	CYC	C4C-C3C-CAC-CBC
6	B	202	CYC	ND-C1D-CHD-C4C
6	B	203	CYC	NA-C4A-CHB-C1B
6	B	203	CYC	C3A-C4A-CHB-C1B
6	D	301	CYC	NA-C4A-CHB-C1B
6	D	301	CYC	C4C-C3C-CAC-CBC
6	D	301	CYC	ND-C1D-CHD-C4C
6	D	302	CYC	NA-C4A-CHB-C1B
6	D	302	CYC	C3A-C4A-CHB-C1B
6	D	302	CYC	ND-C1D-CHD-C4C
6	D	302	CYC	C2D-C1D-CHD-C4C
7	B	204	GOL	O1-C1-C2-C3
6	B	202	CYC	C3A-C4A-CHB-C1B
6	D	302	CYC	C2B-C3B-CAB-CBB
6	B	203	CYC	C2B-C3B-CAB-CBB
4	A	101	M1V	C2A-C3A-CAA-CBA
6	D	301	CYC	C3A-C4A-CHB-C1B
7	B	204	GOL	O1-C1-C2-O2
6	D	302	CYC	C1A-C2A-CAA-CBA
6	D	302	CYC	C3A-C2A-CAA-CBA
5	D	303	AX9	NA-C4A-CHA-C1B
6	D	301	CYC	C2B-C3B-CAB-CBB
6	B	203	CYC	C4B-C3B-CAB-CBB
4	A	101	M1V	CAC-CBC-CGC-O1C
5	D	303	AX9	CAC-CBC-CGC-O2C
6	D	301	CYC	CAD-CBD-CGD-O1D
5	D	303	AX9	CAC-CBC-CGC-O1C
6	B	203	CYC	NC-C4C-CHD-C1D
4	C	101	M1V	CAB-CBB-CGB-O2B
4	C	101	M1V	CAB-CBB-CGB-O1B
4	A	101	M1V	CAC-CBC-CGC-O2C
6	D	301	CYC	CAA-CBA-CGA-O2A
4	A	101	M1V	CAB-CBB-CGB-O2B
4	A	101	M1V	CAB-CBB-CGB-O1B
6	D	301	CYC	CAA-CBA-CGA-O1A
6	D	301	CYC	CAD-CBD-CGD-O2D
5	B	201	AX9	CAB-CBB-CGB-O1B
5	B	201	AX9	CAB-CBB-CGB-O2B
6	B	203	CYC	CAD-CBD-CGD-O1D
4	C	101	M1V	C3B-CAB-CBB-CGB

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	D	303	AX9	CAB-CBB-CGB-O2B
6	B	202	CYC	C2B-C3B-CAB-CBB
4	A	101	M1V	C4A-C3A-CAA-CBA
5	D	303	AX9	CAB-CBB-CGB-O1B
6	D	301	CYC	C2C-C3C-CAC-CBC
6	D	302	CYC	CAA-CBA-CGA-O2A
6	B	203	CYC	CAD-CBD-CGD-O2D


There are no ring outliers.


6 monomers are involved in 12 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	101	M1V	1	0
6	B	203	CYC	3	0
4	C	101	M1V	2	0
6	D	302	CYC	3	0
6	D	301	CYC	1	0
6	B	202	CYC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	80/81 (98%)	-0.43	0	100	100	11, 15, 26, 31	0
2	B	161/173 (93%)	-0.30	0	100	100	11, 19, 29, 33	0
2	D	167/173 (96%)	-0.37	2 (1%)	79	76	10, 15, 27, 62	0
3	C	69/70 (98%)	-0.51	0	100	100	11, 16, 21, 24	0
All	All	477/497 (95%)	-0.38	2 (0%)	92	90	10, 16, 28, 62	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	30	PHE	2.6
2	D	7	ARG	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MEN	B	72	9/10	0.94	0.09	17,22,28,28	0
1	LYZ	A	4	10/11	0.96	0.12	23,31,39,43	0
3	LYZ	C	4	10/11	0.97	0.09	15,20,27,29	0
2	MEN	D	72	9/10	0.98	0.07	12,15,19,19	0

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	B	204	6/6	0.92	0.12	16,20,23,27	0
6	CYC	B	203	43/43	0.93	0.10	13,21,30,35	0
6	CYC	B	202	43/43	0.93	0.10	12,21,28,30	0
6	CYC	D	301	43/43	0.94	0.11	12,17,24,27	0
5	AX9	B	201	43/43	0.95	0.10	9,13,22,28	0
4	M1V	C	101	43/43	0.96	0.09	12,15,19,26	0
6	CYC	D	302	43/43	0.96	0.08	9,14,23,28	0
4	M1V	A	101	43/43	0.96	0.09	12,16,21,27	0
5	AX9	D	303	43/43	0.97	0.10	10,13,18,22	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




















6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.