



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 14, 2023 – 02:52 PM EDT

PDB ID : 7UV7  
Title : Isoreticular, interpenetrating co-crystal of Replication Initiator Protein REPE54 and scaffold duplex (21mer) containing the cognate REPE54 sequence and an insert duplex (10mer) with guest TAMRA-labelled thymine and T-A rich sequence.  
Authors : Orun, A.R.; Snow, C.D.  
Deposited on : 2022-04-29  
Resolution : 3.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2399 (3.04-3.00)
Clashscore	141614	2734 (3.04-3.00)
Ramachandran outliers	138981	2640 (3.04-3.00)
Sidechain outliers	138945	2643 (3.04-3.00)
RSRZ outliers	127900	2287 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	21	 29% 71%
2	B	21	 57% 43%
3	C	263	 65% 16% 19%
4	E	10	 40% 60%
5	F	10	 10% 90%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*AP\*CP\*CP\*TP\*GP\*TP\*GP\*AP\*CP\*AP\*AP\*AP\*TP\*TP\*GP\*CP\*CP\*CP\*TP\*CP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	21	422	203	76	123	20	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(\*TP\*CP\*TP\*GP\*AP\*GP\*GP\*GP\*CP\*AP\*AP\*TP\*TP\*TP\*GP\*TP\*CP\*AP\*CP\*AP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	21	430	206	79	125	20	0	0	0

- Molecule 3 is a protein called Replication initiation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	212	1785	1148	314	316	7	0	3	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-11	MET	-	initiating methionine	UNP P03856
C	-10	ARG	-	expression tag	UNP P03856
C	-9	GLY	-	expression tag	UNP P03856
C	-8	SER	-	expression tag	UNP P03856
C	-7	HIS	-	expression tag	UNP P03856
C	-6	HIS	-	expression tag	UNP P03856
C	-5	HIS	-	expression tag	UNP P03856
C	-4	HIS	-	expression tag	UNP P03856
C	-3	HIS	-	expression tag	UNP P03856
C	-2	HIS	-	expression tag	UNP P03856
C	-1	GLY	-	expression tag	UNP P03856
C	0	SER	-	expression tag	UNP P03856

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	118	PRO	ARG	engineered mutation	UNP P03856

- Molecule 4 is a DNA chain called DNA (5'-D(\*GP\*AP\*CP\*GP\*GP\*TP\*AP\*AP\*TP\*T)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
4	E	10	205	99	39	58	9	0	0	0

- Molecule 5 is a DNA chain called DNA (5'-D(\*GP\*T(TAMRA)P\*AP\*AP\*TP\*TP\*AP\*CP\*CP\*G)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
5	F	10	202	98	37	58	9	0	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	C	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	2	2	2	0	0
7	B	2	2	2	0	0
7	C	4	4	4	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(\*AP\*CP\*CP\*TP\*GP\*TP\*GP\*AP\*CP\*AP\*AP\*AP\*TP\*TP\*GP\*CP\*CP\*CP\*TP\*CP\*A)-3')

Chain A: 




- Molecule 2: DNA (5'-D(\*TP\*CP\*TP\*GP\*AP\*GP\*GP\*GP\*CP\*AP\*AP\*TP\*TP\*TP\*GP\*TP\*CP\*AP\*CP\*AP\*G)-3')

Chain B: 



- Molecule 3: Replication initiation protein

Chain C: 




- Molecule 4: DNA (5'-D(\*GP\*AP\*CP\*GP\*GP\*TP\*AP\*AP\*TP\*T)-3')

Chain E: 



- Molecule 5: DNA (5'-D(\*GP\*T(TAMRA)P\*AP\*AP\*TP\*TP\*AP\*CP\*CP\*G)-3')

Chain F:  10% 90%

G12  
T13  
A14  
A15  
T16  
T17  
A18  
C19  
C20  
G21

## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.49Å 131.39Å 132.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.85 – 3.02 37.85 – 3.02	Depositor EDS
% Data completeness (in resolution range)	96.8 (37.85-3.02) 96.8 (37.85-3.02)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.231 , 0.269 0.229 , 0.267	Depositor DCC
$R_{free}$ test set	1253 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.9	Xtrriage
Anisotropy	0.451	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 82.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.008 for -h,-l,-k	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3053	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	114.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.62	0/472	0.95	0/725
2	B	0.63	0/482	1.00	0/743
3	C	0.29	0/1828	0.59	0/2461
4	E	0.76	0/230	1.18	1/354 (0.3%)
5	F	0.98	2/226 (0.9%)	1.22	2/347 (0.6%)
All	All	0.52	2/3238 (0.1%)	0.84	3/4630 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	13	DT	C5-C6	-6.63	1.29	1.34
5	F	13	DT	N1-C6	-5.34	1.34	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	13	DT	C6-C5-C7	-5.73	119.46	122.90
4	E	26	DT	N3-C4-O4	5.18	123.01	119.90
5	F	13	DT	C4-C5-C7	5.05	122.03	119.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	422	0	238	14	0
2	B	430	0	239	7	0
3	C	1785	0	1766	27	0
4	E	205	0	115	4	0
5	F	202	0	115	8	0
6	C	1	0	0	0	0
7	A	2	0	0	0	0
7	B	2	0	0	0	0
7	C	4	0	0	0	0
All	All	3053	0	2473	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:DT:H2'	1:A:14:DG:C8	2.14	0.82
3:C:152:SER:HA	3:C:155:LYS:HD3	1.70	0.71
4:E:29:DT:H2''	4:E:30:DT:H3'	1.74	0.69
3:C:94:VAL:HG22	3:C:114:TRP:CE2	2.29	0.67
5:F:20:DC:H2''	5:F:21:DG:C8	2.35	0.61
2:B:22:DT:H2''	2:B:23:DC:C6	2.36	0.60
3:C:122:PRO:HD2	3:C:126:LEU:HD23	1.83	0.60
1:A:6:DG:O6	3:C:206:ARG:NH2	2.36	0.58
3:C:78:ALA:O	3:C:82:ILE:HD12	2.04	0.57
3:C:190[A]:GLU:OE1	3:C:191:ARG:NH1	2.39	0.56
2:B:40:DC:H2''	2:B:41:DA:H8	1.70	0.55
1:A:5:DT:H2''	1:A:6:DG:C8	2.42	0.54
2:B:40:DC:H2''	2:B:41:DA:C8	2.41	0.54
5:F:20:DC:OP2	5:F:20:DC:H6	1.90	0.54
2:B:37:DT:OP2	3:C:195:PRO:HB2	2.08	0.54
5:F:19:DC:H2''	5:F:20:DC:C5	2.43	0.54
3:C:152:SER:HA	3:C:155:LYS:CD	2.36	0.53
5:F:14:DA:H2'	5:F:14:DA:OP2	2.08	0.53
3:C:21:SER:HB3	3:C:138:PHE:O	2.10	0.52
3:C:45:GLN:OE1	3:C:60:CYS:HB3	2.09	0.52
5:F:19:DC:H2''	5:F:20:DC:C6	2.44	0.52
3:C:28:ALA:HB3	3:C:95:VAL:HB	1.93	0.51
3:C:174:LYS:HB3	3:C:176:ASP:OD1	2.11	0.51
3:C:94:VAL:O	3:C:111:SER:HA	2.11	0.50
1:A:4:DG:C8	1:A:5:DT:H72	2.47	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:27:DG:H2'	2:B:28:DG:C8	2.48	0.48
1:A:0:DA:HO5'	1:A:0:DA:H8	1.58	0.48
1:A:19:DC:H2''	1:A:20:DA:C8	2.49	0.48
2:B:27:DG:H2'	2:B:28:DG:H8	1.79	0.47
3:C:185:ILE:HD11	3:C:238:ILE:HG12	1.96	0.47
4:E:23:DC:H2''	4:E:24:DG:C8	2.49	0.47
5:F:15:DA:H1'	5:F:16:DT:H5'	1.96	0.46
2:B:30:DC:H2'	2:B:31:DA:C8	2.51	0.46
3:C:14:ASN:OD1	3:C:150:ARG:NH1	2.49	0.45
1:A:15:DC:H2'	1:A:16:DC:C6	2.51	0.45
3:C:19:VAL:HG22	3:C:47:ARG:HD2	1.97	0.45
3:C:196:GLN:HG3	3:C:197:SER:N	2.31	0.45
1:A:6:DG:H1'	1:A:7:DA:H5''	1.98	0.45
4:E:30:DT:O4	5:F:14:DA:N1	2.50	0.45
1:A:15:DC:H2'	1:A:16:DC:H6	1.81	0.44
3:C:19:VAL:HG12	3:C:148:GLN:HB2	1.99	0.44
5:F:17:DT:H2''	5:F:18:DA:C8	2.53	0.43
1:A:8:DC:H2''	1:A:9:DA:C8	2.53	0.43
3:C:56:HIS:CG	3:C:57:ASP:H	2.37	0.43
4:E:21:DG:H5''	4:E:21:DG:H8	1.84	0.43
3:C:164[B]:ARG:HA	3:C:164[B]:ARG:HD2	1.83	0.42
3:C:83:ARG:O	3:C:87:LYS:HG3	2.20	0.42
1:A:8:DC:H2''	1:A:9:DA:H8	1.85	0.42
3:C:47:ARG:HH11	3:C:47:ARG:HG3	1.84	0.42
1:A:12:DT:H2'	1:A:13:DT:H72	2.01	0.42
1:A:2:DC:H4'	3:C:233:ARG:HB3	2.02	0.41
3:C:136:ILE:HB	3:C:137:PRO:HD3	2.02	0.41
3:C:182:SER:HB2	3:C:239:VAL:HG22	2.03	0.41
1:A:5:DT:H2''	1:A:6:DG:H8	1.85	0.40
3:C:150:ARG:HG3	3:C:152:SER:OG	2.22	0.40
3:C:180:ILE:HA	3:C:240:PHE:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	207/263 (79%)	204 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	194/236 (82%)	194 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	21/21 (100%)	-0.46	0 100 100	80, 116, 126, 146	0
2	B	21/21 (100%)	-0.44	0 100 100	82, 113, 135, 148	0
3	C	212/263 (80%)	-0.31	0 100 100	56, 93, 147, 209	0
4	E	10/10 (100%)	0.18	0 100 100	173, 178, 184, 189	0
5	F	10/10 (100%)	0.31	0 100 100	165, 175, 195, 199	0
All	All	274/325 (84%)	-0.29	0 100 100	56, 100, 176, 209	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	MG	C	301	1/1	0.97	0.12	117,117,117,117	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.