



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 03:48 PM JST

PDB ID : 7VJE  
Title : class II photolyase MmCPDII semiquinone to fully reduced TR-SFX studies (300 ns time-point)  
Authors : Maestre-Reyna, M.; Yang, C.-H.; Huang, W.-C.; Nango, E.; Ngura Putu, E.P.G.; Franz-Badur, S.; Wu, W.-J.; Wu, H.-Y.; Wang, P.-H.; Hosokawa, Y.; Saft, M.; Emmerich, H.-J.; Liao, J.-H.; Lee, C.-C.; Huang, K.-F.; Chang, Y.-K.; Weng, J.-H.; Royant, A.; Gad, W.; Pang, A.H.; Chang, C.-W.; Sugahara, M.; Owada, S.; Joti, Y.; Yamashita, A.; Tanaka, R.; Tanaka, T.; Luo, F.J.; Tono, K.; Kiontke, S.; Yamamoto, J.; Iwata, S.; Essen, L.-O.; Bessho, Y.; Tsai, M.-D.  
Deposited on : 2021-09-28  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	482	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DTT	A	505	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA photolyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	453	3572	2315	584	657	16	0	7	0

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP Q8PYK9
A	-16	GLY	-	expression tag	UNP Q8PYK9
A	-15	SER	-	expression tag	UNP Q8PYK9
A	-14	SER	-	expression tag	UNP Q8PYK9
A	-13	HIS	-	expression tag	UNP Q8PYK9
A	-12	HIS	-	expression tag	UNP Q8PYK9
A	-11	HIS	-	expression tag	UNP Q8PYK9
A	-10	HIS	-	expression tag	UNP Q8PYK9
A	-9	HIS	-	expression tag	UNP Q8PYK9
A	-8	HIS	-	expression tag	UNP Q8PYK9
A	-7	SER	-	expression tag	UNP Q8PYK9
A	-6	SER	-	expression tag	UNP Q8PYK9
A	-5	GLY	-	expression tag	UNP Q8PYK9
A	-4	LEU	-	expression tag	UNP Q8PYK9
A	-3	VAL	-	expression tag	UNP Q8PYK9
A	-2	PRO	-	expression tag	UNP Q8PYK9
A	-1	ARG	-	expression tag	UNP Q8PYK9
A	0	GLY	-	expression tag	UNP Q8PYK9
A	1	SER	-	expression tag	UNP Q8PYK9
A	2	HIS	-	expression tag	UNP Q8PYK9
A	377	THR	MET	engineered mutation	UNP Q8PYK9

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	53	27	9	15	2	0	0

- Molecule 4 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula:  $C_4H_{10}O_2S_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
4	A	1	8	4	2	2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	188	Total	O	0	0
			188	188		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA photolyase

Chain A:  76% 17% 6%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.52Å 70.52Å 246.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.68 – 2.50 32.40 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (31.68-2.50) 100.0 (32.40-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.41 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.19_4092: ???)	Depositor
R, $R_{free}$	0.223 , 0.248 0.279 , 0.290	Depositor DCC
$R_{free}$ test set	2095 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.3	Xtrriage
Anisotropy	0.009	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 58.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	3836	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FAD, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/3692	0.45	0/5017

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3572	0	3341	51	0
2	A	15	0	0	0	0
3	A	53	0	28	0	0
4	A	8	0	10	4	0
5	A	188	0	0	3	0
All	All	3836	0	3379	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:TYR:CD1	1:A:158:TYR:CZ	2.38	1.03
1:A:379:MET:CE	4:A:505:DTT:H2	1.90	1.00
1:A:39:PHE:CG	1:A:39:PHE:CE1	2.40	0.97
1:A:379:MET:HE3	4:A:505:DTT:H2	1.52	0.90
1:A:379:MET:HE1	4:A:505:DTT:H2	1.60	0.82
1:A:378:ARG:NH2	1:A:407:GLU:OE1	2.23	0.71
1:A:51:VAL:HG23	1:A:88:ILE:HG23	1.87	0.57
1:A:86:LYS:NZ	5:A:619:HOH:O	2.39	0.56
1:A:383:LYS:HD2	1:A:443:MET:HG3	1.86	0.56
1:A:256:ARG:NH1	5:A:620:HOH:O	2.40	0.54
1:A:118:SER:HB3	1:A:123:LYS:HG2	1.89	0.54
1:A:266:ASN:HA	1:A:372:LYS:HE3	1.91	0.51
1:A:366:GLU:OE1	1:A:457:TYR:OH	2.22	0.51
1:A:66:ILE:HD12	1:A:215:ARG:HG3	1.93	0.50
1:A:92:PHE:O	1:A:197:VAL:N	2.45	0.49
1:A:378:ARG:NH2	1:A:407:GLU:CD	2.66	0.49
1:A:150:TRP:CD1	1:A:151:GLU:HG2	2.48	0.49
1:A:386:LEU:HD13	1:A:423:ILE:HG23	1.95	0.48
1:A:224:LEU:HD13	1:A:368:LEU:HB3	1.96	0.48
1:A:379:MET:HE1	4:A:505:DTT:C2	2.38	0.47
1:A:373:MET:O	1:A:378:ARG:HD2	2.14	0.47
1:A:73[A]:LEU:HD23	1:A:204:LEU:HB2	1.96	0.47
1:A:129:LYS:HE3	1:A:129:LYS:HB2	1.63	0.47
1:A:41:ARG:HA	1:A:51:VAL:HG11	1.97	0.46
1:A:378:ARG:NH2	1:A:409:ASP:OD1	2.48	0.45
1:A:255:LEU:HB2	1:A:263:MET:HG2	1.99	0.44
1:A:98:GLY:HA3	1:A:129:LYS:HG2	1.98	0.44
1:A:27:ARG:NH2	5:A:615:HOH:O	2.48	0.44
1:A:345:TYR:OH	1:A:356:HIS:ND1	2.49	0.44
1:A:442:TYR:HE1	1:A:444:SER:HB3	1.82	0.44
1:A:62:LEU:HD12	1:A:211:LEU:HG	2.01	0.43
1:A:435:GLU:O	1:A:437:THR:N	2.46	0.43
1:A:28:ASP:HA	1:A:274:GLY:HA3	2.01	0.43
1:A:72:MET:O	1:A:76:LEU:HG	2.18	0.43
1:A:259:PRO:HA	1:A:374:HIS:CD2	2.53	0.43
1:A:23:TYR:CE2	1:A:25:MET:HB2	2.54	0.43
1:A:58:THR:HG21	1:A:126[B]:TRP:CZ2	2.53	0.43
1:A:157:GLU:HG3	1:A:163:PHE:CD1	2.54	0.42
1:A:22:VAL:HA	1:A:52:VAL:HG23	2.00	0.42
1:A:384:LYS:HD3	1:A:384:LYS:HA	1.81	0.42
1:A:403:ASN:ND2	1:A:403:ASN:C	2.73	0.42
1:A:24:TRP:N	1:A:114:VAL:O	2.40	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:PHE:HD1	1:A:248:ARG:HB2	1.84	0.41
1:A:34:ASN:O	1:A:38:LEU:HD12	2.20	0.41
1:A:62:LEU:HA	1:A:211:LEU:HD21	2.00	0.41
1:A:30:ARG:NE	1:A:33:ASP:O	2.44	0.41
1:A:147:VAL:HG13	1:A:174:PHE:CD2	2.55	0.41
1:A:282:VAL:HG21	1:A:303:LEU:HD21	2.02	0.41
1:A:290:SER:HB2	1:A:291:ASN:H	1.74	0.41
1:A:359:LEU:HB2	1:A:445:TYR:HE1	1.86	0.41
1:A:268:SER:OG	1:A:269:PRO:HD3	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	456/482 (95%)	439 (96%)	17 (4%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	355/416 (85%)	343 (97%)	12 (3%)	<a href="#">37</a> <a href="#">63</a>

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	MET
1	A	49	VAL
1	A	52	VAL
1	A	62	LEU
1	A	79	LEU
1	A	85	ARG
1	A	199	THR
1	A	218	LEU
1	A	389	SER
1	A	403	ASN
1	A	443	MET
1	A	462	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	343	HIS
1	A	403	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DTT	A	505	-	7,7,7	0.34	0	4,8,8	0.62	0
3	FAD	A	503	-	53,58,58	3.48	23 (43%)	68,89,89	1.65	17 (25%)
2	SO4	A	504	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	A	501	-	4,4,4	0.13	0	6,6,6	0.05	0
2	SO4	A	502	-	4,4,4	0.14	0	6,6,6	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTT	A	505	-	-	3/8/8/8	-
3	FAD	A	503	-	-	7/30/50/50	0/6/6/6

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	FAD	C2B-C1B	-12.12	1.35	1.53
3	A	503	FAD	O2'-C2'	-7.43	1.27	1.43
3	A	503	FAD	O4'-C4'	-6.86	1.28	1.43
3	A	503	FAD	C3B-C4B	-6.73	1.35	1.53
3	A	503	FAD	C9A-N10	-6.36	1.29	1.41
3	A	503	FAD	C4'-C3'	-6.13	1.41	1.53
3	A	503	FAD	O4-C4	5.91	1.34	1.23
3	A	503	FAD	O2-C2	5.90	1.35	1.24
3	A	503	FAD	C2B-C3B	-5.65	1.37	1.53
3	A	503	FAD	C2'-C3'	-5.49	1.43	1.53
3	A	503	FAD	C4-N3	-4.32	1.30	1.38
3	A	503	FAD	O3'-C3'	-3.75	1.34	1.43
3	A	503	FAD	C6A-N6A	3.22	1.45	1.34
3	A	503	FAD	P-O5'	2.86	1.70	1.59
3	A	503	FAD	C5B-C4B	-2.78	1.42	1.51
3	A	503	FAD	C10-N10	-2.58	1.31	1.37
3	A	503	FAD	C2-N3	-2.55	1.33	1.39
3	A	503	FAD	PA-O5B	2.48	1.69	1.59
3	A	503	FAD	PA-O2A	2.48	1.67	1.55
3	A	503	FAD	P-O2P	2.29	1.66	1.55

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	FAD	C8M-C8	2.29	1.55	1.51
3	A	503	FAD	O4B-C1B	2.16	1.44	1.41
3	A	503	FAD	C5X-N5	-2.15	1.35	1.39

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	FAD	N3A-C2A-N1A	-4.19	122.13	128.68
3	A	503	FAD	C5X-N5-C4X	-4.13	111.21	118.07
3	A	503	FAD	C4A-C5A-N7A	-3.86	105.37	109.40
3	A	503	FAD	C4X-C10-N1	-3.66	116.23	124.73
3	A	503	FAD	C5'-C4'-C3'	-3.50	105.44	112.20
3	A	503	FAD	C4X-C10-N10	3.03	120.91	116.48
3	A	503	FAD	C3B-C2B-C1B	2.98	105.47	100.98
3	A	503	FAD	C4X-C4-N3	2.97	120.73	113.19
3	A	503	FAD	C4-N3-C2	-2.86	120.36	125.64
3	A	503	FAD	C1B-N9A-C4A	-2.83	121.66	126.64
3	A	503	FAD	C9A-N10-C10	-2.64	116.65	120.77
3	A	503	FAD	C4-C4X-N5	-2.42	114.79	118.23
3	A	503	FAD	O4-C4-C4X	-2.34	120.39	126.60
3	A	503	FAD	C4-C4X-C10	2.24	120.55	116.79
3	A	503	FAD	P-O3P-PA	-2.24	125.16	132.83
3	A	503	FAD	C10-N1-C2	2.20	121.29	116.90
3	A	503	FAD	C5X-C9A-N10	2.11	120.13	117.95

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	503	FAD	N10-C1'-C2'-C3'
4	A	505	DTT	C1-C2-C3-O3
4	A	505	DTT	O2-C2-C3-O3
4	A	505	DTT	O2-C2-C3-C4
3	A	503	FAD	P-O3P-PA-O1A
3	A	503	FAD	C4'-C5'-O5'-P
3	A	503	FAD	O4'-C4'-C5'-O5'
3	A	503	FAD	C5B-O5B-PA-O3P
3	A	503	FAD	P-O3P-PA-O2A
3	A	503	FAD	C5B-O5B-PA-O1A

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	505	DTT	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.