

# Full wwPDB X-ray Structure Validation Report (i)

Nov 29, 2022 – 12:30 pm GMT

PDB ID : 8APP

Title : AbLys1 endolysin from Acinetobacter baumannii phage AbTZA1 Authors : Premetis, G.E.; Stathi, A.; Papageorgiou, A.C.; Labrou, N.E.

Deposited on : 2022-08-10

Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.31.3

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0267

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	193	78%	%	
1	В	193	90%	8%	•••
1	С	193	89%	9%	••
1	D	193	82%	15%	• •



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6539 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Endolysin.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	187	Total	С	N	О	S	0	0	0
1	A	107	1468	936	259	271	2	0	0	U
1	В	189	Total	С	N	О	S	0	0	0
1	Ъ	109	1486	947	263	273	3	0	0	U
1	С	190	Total	С	N	О	S	0	0	0
1		190	1496	953	266	274	3	0	0	U
1	D	188	Total	С	N	О	S	0	0	0
1	D	100	1476	941	260	272	3	0	0	U

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	188	HIS	-	expression tag	UNP A0A3T0IGR7
A	189	HIS	-	expression tag	UNP A0A3T0IGR7
A	190	HIS	-	expression tag	UNP A0A3T0IGR7
A	191	HIS	-	expression tag	UNP A0A3T0IGR7
A	192	HIS	-	expression tag	UNP A0A3T0IGR7
A	193	HIS	-	expression tag	UNP A0A3T0IGR7
В	188	HIS	-	expression tag	UNP A0A3T0IGR7
В	189	HIS	-	expression tag	UNP A0A3T0IGR7
В	190	HIS	-	expression tag	UNP A0A3T0IGR7
В	191	HIS	-	expression tag	UNP A0A3T0IGR7
В	192	HIS	-	expression tag	UNP A0A3T0IGR7
В	193	HIS	-	expression tag	UNP A0A3T0IGR7
С	188	HIS	-	expression tag	UNP A0A3T0IGR7
С	189	HIS	-	expression tag	UNP A0A3T0IGR7
С	190	HIS	-	expression tag	UNP A0A3T0IGR7
С	191	HIS	-	expression tag	UNP A0A3T0IGR7
С	192	HIS	-	expression tag	UNP A0A3T0IGR7
С	193	HIS	-	expression tag	UNP A0A3T0IGR7
D	188	HIS	-	expression tag	UNP A0A3T0IGR7
D	189	HIS	-	expression tag	UNP A0A3T0IGR7
D	190	HIS	-	expression tag	UNP A0A3T0IGR7



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
D	191	HIS	-	expression tag	UNP A0A3T0IGR7
D	192	HIS	-	expression tag	UNP A0A3T0IGR7
D	193	HIS	-	expression tag	UNP A0A3T0IGR7


• Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	С	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0

 $\bullet$  Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $\mathrm{C_3H_8O_3}).$ 





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	В	1	Total 6	C 3	O 3	0	0

#### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	153	Total O 153 153	0	0
4	В	157	Total O 157 157	0	0
4	С	146	Total O 146 146	0	0
4	D	131	Total O 131 131	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endolysin Chain A: 18% • Molecule 1: Endolysin Chain B: 90% • Molecule 1: Endolysin Chain C: 89% • Molecule 1: Endolysin Chain D: 82% 15%



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	41.65Å 184.78Å 57.90Å	Donositon
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $110.69^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	40.68 - 1.82	Depositor
Resolution (A)	46.19 - 1.82	EDS
% Data completeness	96.1 (40.68-1.82)	Depositor
(in resolution range)	88.8 (46.19-1.82)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.38 \; (at \; 1.82 \text{Å})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
$R, R_{free}$	0.211 , 0.252	Depositor
10, 10 free	0.211 , $0.253$	DCC
$R_{free}$ test set	1992 reflections $(2.84\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.9	Xtriage
Anisotropy	0.199	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6539	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.44	0/1493	0.63	0/2014	
1	В	0.42	0/1512	0.62	0/2039	
1	С	0.41	0/1523	0.64	0/2054	
1	D	0.39	0/1501	0.63	0/2024	
All	All	0.42	0/6029	0.63	0/8131	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1468	0	1496	29	0
1	В	1486	0	1515	11	0
1	С	1496	0	1522	13	0
1	D	1476	0	1508	27	0
2	A	10	0	0	0	0
2	С	5	0	0	0	0
2	D	5	0	0	1	0
3	В	6	0	8	1	0
4	A	153	0	0	9	2



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	В	157	0	0	3	2
4	С	146	0	0	10	0
4	D	131	0	0	10	0
All	All	6539	0	6049	76	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A.1 1	A	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\rm \mathring{A})$	overlap (Å)
1:C:176:ASN:ND2	4:C:304:HOH:O	2.04	0.88
1:A:50:GLN:NE2	4:A:303:HOH:O	2.07	0.88
1:D:160:ASN:HD22	1:D:175:THR:HG22	1.42	0.84
1:C:8:ASP:OD1	4:C:301:HOH:O	1.95	0.83
1:C:176:ASN:OD1	4:C:302:HOH:O	1.95	0.82
1:A:50:GLN:O	4:A:301:HOH:O	1.99	0.79
1:C:136:ALA:O	4:C:303:HOH:O	2.00	0.79
1:A:111:VAL:O	4:A:302:HOH:O	2.03	0.75
1:D:167:LYS:N	4:D:303:HOH:O	2.21	0.72
1:A:116:LYS:HD3	1:A:117:GLN:H	1.52	0.71
1:A:180:LYS:HE3	1:D:57:ARG:HH22	1.59	0.68
1:D:188:HIS:O	4:D:301:HOH:O	2.11	0.67
1:C:142:LYS:NZ	4:C:306:HOH:O	2.28	0.66
1:D:164:VAL:N	4:D:303:HOH:O	2.28	0.65
1:A:180:LYS:HE3	1:D:57:ARG:NH2	2.13	0.64
1:A:111:VAL:HG11	1:A:115:LEU:HD11	1.81	0.63
1:A:102:LYS:HB3	4:A:327:HOH:O	1.99	0.62
1:C:142:LYS:HE3	4:C:303:HOH:O	2.00	0.60
1:A:45:VAL:O	1:A:49:ARG:HG3	2.04	0.58
1:D:50:GLN:HE22	1:D:180:LYS:HE3	1.69	0.58
1:D:182:LYS:NZ	1:D:186:GLU:OE2	2.37	0.57
1:A:2:ASP:N	4:A:307:HOH:O	2.37	0.57
1:D:40:ILE:HD12	1:D:121:ASP:OD1	2.05	0.56
1:A:140:LEU:HD12	1:A:155:GLN:HB3	1.87	0.56
1:A:97:LYS:NZ	4:A:310:HOH:O	2.39	0.54
1:A:54:LEU:HD22	1:A:89:GLU:HG3	1.90	0.53
1:D:157:LEU:HD21	1:D:182:LYS:HD3	1.90	0.53
1:A:47:LEU:O	1:A:50:GLN:HG2	2.08	0.53
1:A:116:LYS:HD3	1:A:117:GLN:N	2.24	0.51
1:D:145:ASN:O	4:D:302:HOH:O	2.19	0.51



 $Continued\ from\ previous\ page...$ 

Continued from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap(Å)	
1:A:32:ARG:HD3	4:A:323:HOH:O	2.10	0.50	
1:D:74:ARG:NH1	4:D:308:HOH:O	2.44	0.50	
1:B:28:LYS:NZ	4:B:307:HOH:O	2.44	0.50	
1:C:82:LYS:HE3	4:C:369:HOH:O	2.10	0.50	
1:A:144:LEU:O	1:A:146:SER:N	2.43	0.50	
1:B:1:MET:SD	1:B:34:VAL:HG11	2.52	0.50	
1:D:1:MET:N	4:D:309:HOH:O	2.45	0.50	
1:C:163:ARG:HG3	1:C:167:LYS:O	2.12	0.50	
1:C:121:ASP:OD2	4:C:305:HOH:O	2.19	0.49	
1:A:21:ALA:O	1:A:25:GLU:HG3	2.12	0.48	
1:B:142:LYS:HE3	4:B:312:HOH:O	2.12	0.48	
1:B:57:ARG:HH11	1:B:57:ARG:HG2	1.79	0.47	
1:B:66:TRP:CZ2	1:B:83:ALA:HB2	2.49	0.47	
1:D:1:MET:HG2	1:D:2:ASP:N	2.29	0.47	
1:B:53:GLY:HA3	3:B:201:GOL:H2	1.97	0.46	
1:A:155:GLN:OE1	4:A:304:HOH:O	2.21	0.46	
1:B:57:ARG:HD3	1:B:86:VAL:HG12	1.98	0.46	
1:D:20:GLN:HG2	2:D:201:PO4:O1	2.16	0.46	
1:D:40:ILE:HD11	1:D:44:GLY:HA3	1.98	0.46	
1:A:161:LYS:HD2	1:A:168:LEU:HB3	1.98	0.46	
1:A:165:LYS:HA	1:A:165:LYS:HD3	1.56	0.45	
1:B:116:LYS:NZ	4:B:301:HOH:O	2.08	0.45	
1:A:10:ALA:CB	1:A:26:LEU:HD11	2.47	0.45	
1:A:89:GLU:OE2	1:D:89:GLU:OE1	2.34	0.44	
1:C:57:ARG:HD3	1:C:86:VAL:HG22	1.99	0.44	
1:A:134:ALA:O	1:A:138:SER:HB3	2.18	0.44	
1:C:2:ASP:OD1	1:C:4:LYS:N	2.49	0.44	
1:D:115:LEU:N	4:D:305:HOH:O	2.35	0.44	
1:D:167:LYS:HE2	1:D:167:LYS:HB3	1.87	0.43	
1:B:188:HIS:HB2	1:B:189:HIS:H	1.69	0.42	
1:D:179:LYS:NZ	4:D:307:HOH:O	2.42	0.42	
1:B:45:VAL:O	1:B:49:ARG:HG3	2.19	0.42	
1:A:143:LYS:HD3	1:A:151:GLY:HA3	2.01	0.42	
1:A:73:ILE:HG13	1:A:74:ARG:HG3	2.01	0.42	
1:D:61:ASP:HB2	4:D:326:HOH:O	2.20	0.42	
1:A:113:VAL:HG13	1:A:145:ASN:HA	2.02	0.41	
1:D:165:LYS:HA	1:D:165:LYS:HD3	1.91	0.41	
1:D:52:GLU:OE1	4:D:304:HOH:O	2.22	0.41	
1:A:109:LYS:HE2	4:A:309:HOH:O	2.20	0.41	
1:D:1:MET:HG2	1:D:2:ASP:H	1.85	0.41	
1:A:53:GLY:HA2	1:D:54:LEU:O	2.21	0.41	



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:C:161:LYS:HE2	4:C:316:HOH:O	2.20	0.41
1:D:174:LEU:HD23	1:D:174:LEU:HA	1.86	0.41
1:D:58:ALA:HA	1:D:67:THR:O	2.20	0.40
1:B:55:ARG:HG3	1:B:59:TYR:CE1	2.56	0.40
1:C:180:LYS:NZ	4:C:309:HOH:O	2.45	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$	
4:A:307:HOH:O	4:B:375:HOH:O[1_556]	1.82	0.38	
4:A:439:HOH:O	4:B:410:HOH:O[1_556]	2.18	0.02	

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	185/193 (96%)	176 (95%)	7 (4%)	2 (1%)	14	4
1	В	187/193 (97%)	183 (98%)	4 (2%)	0	100	100
1	$\mathbf{C}$	188/193 (97%)	180 (96%)	8 (4%)	0	100	100
1	D	186/193 (96%)	182 (98%)	4 (2%)	0	100	100
All	All	$746/772 \ (97\%)$	721 (97%)	23 (3%)	2 (0%)	41	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	SER
1	A	145	ASN



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	154/160~(96%)	153 (99%)	1 (1%)		86	83
1	В	156/160 (98%)	155 (99%)	1 (1%)		86	83
1	C	157/160 (98%)	152 (97%)	5 (3%)		39	24
1	D	155/160 (97%)	154 (99%)	1 (1%)		86	83
All	All	622/640 (97%)	614 (99%)	8 (1%)		69	61

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	В	189	HIS
1	С	32	ARG
1	С	36	ASP
1	С	74	ARG
1	С	82	LYS
1	С	190	HIS
1	D	74	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	В	20	GLN
1	В	50	GLN
1	С	190	HIS
1	D	50	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain Res		Link	Bond lengths			Bond angles				
Mol	Type	Chain	nes	Res   Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	PO4	С	201	-	4,4,4	1.10	0	6,6,6	0.34	0
2	PO4	D	201	-	4,4,4	1.12	0	6,6,6	0.49	0
2	PO4	A	202	-	4,4,4	1.16	0	6,6,6	0.71	0
2	PO4	A	201	-	4,4,4	0.97	0	6,6,6	0.69	0
3	GOL	В	201	-	5,5,5	0.97	0	5,5,5	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	В	201	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	201	PO4	1	0
3	В	201	GOL	1	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	187/193 (96%)	-0.09	3 (1%) 72 68	29, 37, 56, 64	0
1	В	189/193 (97%)	-0.21	3 (1%) 72 68	28, 36, 45, 69	0
1	С	190/193 (98%)	-0.11	4 (2%) 63 59	28, 38, 47, 68	0
1	D	188/193 (97%)	-0.04	3 (1%) 72 68	30, 40, 52, 63	0
All	All	754/772 (97%)	-0.11	13 (1%) 70 66	28, 38, 52, 69	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	36	ASP	3.3
1	D	164	VAL	3.3
1	A	149	TYR	3.2
1	A	151	GLY	3.2
1	D	168	LEU	3.0
1	A	147	LYS	3.0
1	В	189	HIS	2.8
1	D	142	LYS	2.6
1	С	190	HIS	2.6
1	С	36	ASP	2.6
1	С	172	PRO	2.3
1	В	37	ASP	2.2
1	С	142	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	GOL	В	201	6/6	0.87	0.19	39,43,45,45	0
2	PO4	A	202	5/5	0.99	0.08	36,39,47,49	0
2	PO4	С	201	5/5	0.99	0.12	39,39,40,45	0
2	PO4	D	201	5/5	0.99	0.12	42,42,48,49	0
2	PO4	A	201	5/5	0.99	0.06	41,44,47,48	0

#### 6.5 Other polymers (i)

There are no such residues in this entry.

