



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2024 – 12:43 pm GMT

PDB ID : 8CLP
Title : Zearalenone lactonase from *Streptomyces coelicoflavus* mutant H286Y
Authors : Puehringer, D.; Mlynek, G.
Deposited on : 2023-02-17
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	 2% 89% 7%
1	B	315	 2% 86% 7% 7%
1	C	315	 2% 86% 10%
1	D	315	 2% 88% 8%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 18289 atoms, of which 8626 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	294	4442	1453	2151	401	426	11	0	0	0
1	B	292	4458	1446	2178	398	424	12	0	2	0
1	C	284	4341	1415	2117	388	410	11	0	0	0
1	D	291	4459	1448	2180	398	422	11	0	2	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	286	TYR	HIS	engineered mutation	UNP H1Q8U7
A	310	HIS	-	expression tag	UNP H1Q8U7
A	311	HIS	-	expression tag	UNP H1Q8U7
A	312	HIS	-	expression tag	UNP H1Q8U7
A	313	HIS	-	expression tag	UNP H1Q8U7
A	314	HIS	-	expression tag	UNP H1Q8U7
A	315	HIS	-	expression tag	UNP H1Q8U7
B	286	TYR	HIS	engineered mutation	UNP H1Q8U7
B	310	HIS	-	expression tag	UNP H1Q8U7
B	311	HIS	-	expression tag	UNP H1Q8U7
B	312	HIS	-	expression tag	UNP H1Q8U7
B	313	HIS	-	expression tag	UNP H1Q8U7
B	314	HIS	-	expression tag	UNP H1Q8U7
B	315	HIS	-	expression tag	UNP H1Q8U7
C	286	TYR	HIS	engineered mutation	UNP H1Q8U7
C	310	HIS	-	expression tag	UNP H1Q8U7
C	311	HIS	-	expression tag	UNP H1Q8U7
C	312	HIS	-	expression tag	UNP H1Q8U7
C	313	HIS	-	expression tag	UNP H1Q8U7
C	314	HIS	-	expression tag	UNP H1Q8U7
C	315	HIS	-	expression tag	UNP H1Q8U7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	286	TYR	HIS	engineered mutation	UNP H1Q8U7
D	310	HIS	-	expression tag	UNP H1Q8U7
D	311	HIS	-	expression tag	UNP H1Q8U7
D	312	HIS	-	expression tag	UNP H1Q8U7
D	313	HIS	-	expression tag	UNP H1Q8U7
D	314	HIS	-	expression tag	UNP H1Q8U7
D	315	HIS	-	expression tag	UNP H1Q8U7


- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	147	Total O 147 147	0	0
2	B	158	Total O 158 158	0	0
2	C	114	Total O 114 114	0	0
2	D	170	Total O 170 170	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydrolase

Chain A: 




- Molecule 1: Hydrolase

Chain B: 



- Molecule 1: Hydrolase

Chain C: 



- Molecule 1: Hydrolase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.57Å 126.13Å 128.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.57 – 1.92 45.04 – 1.92	Depositor EDS
% Data completeness (in resolution range)	95.9 (42.57-1.92) 96.0 (45.04-1.92)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.92Å)	Xtrriage
Refinement program	PHENIX 1.20.1-4489	Depositor
R, R_{free}	0.197 , 0.250 0.197 , 0.250	Depositor DCC
R_{free} test set	1994 reflections (2.61%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtrriage
Anisotropy	0.625	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.012 for -h,l,k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18289	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2358	0.49	0/3215
1	B	0.24	0/2350	0.49	0/3202
1	C	0.24	0/2290	0.48	0/3122
1	D	0.24	0/2357	0.50	0/3213
All	All	0.24	0/9355	0.49	0/12752

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	193	ARG	Sidechain
1	C	104	ARG	Sidechain
1	C	38	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2291	2151	2181	9	0
1	B	2280	2178	2172	9	0
1	C	2224	2117	2121	5	0
1	D	2279	2180	2169	6	0
2	A	147	0	0	0	0
2	B	158	0	0	0	0
2	C	114	0	0	0	0
2	D	170	0	0	1	0
All	All	9663	8626	8643	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:ALA:HB2	1:A:300:PHE:HE2	1.60	0.66
1:A:177:TRP:CZ3	1:A:204:LEU:HD23	2.38	0.57
1:B:102:VAL:HG11	1:B:106:VAL:CG1	2.35	0.56
1:B:184:ALA:O	1:B:187:SER:OG	2.20	0.50
1:B:134:LEU:HD12	1:B:238:LEU:HD23	1.96	0.48
1:B:160:VAL:HG22	1:B:192:ALA:HB2	1.95	0.47
1:A:109:ALA:HB2	1:A:300:PHE:CE2	2.45	0.47
1:C:130:LEU:HD23	1:C:133:VAL:CG1	2.45	0.46
1:D:102:VAL:HG11	1:D:106:VAL:CG1	2.44	0.46
1:C:32:GLU:OE2	1:C:66:HIS:NE2	2.48	0.46
1:D:182:ARG:HG2	1:D:182:ARG:NH1	2.31	0.45
1:C:53:SER:O	1:C:290:GLN:HG3	2.17	0.44
1:A:177:TRP:CD2	1:A:200:ILE:HG12	2.53	0.44
1:C:281:VAL:O	1:C:281:VAL:HG12	2.19	0.43
1:D:290:GLN:NE2	2:D:417:HOH:O	2.52	0.43
1:A:177:TRP:HZ3	1:A:204:LEU:CD2	2.32	0.43
1:D:237:VAL:HG11	1:D:269:MET:SD	2.59	0.42
1:A:102:VAL:HG11	1:A:106:VAL:CG1	2.50	0.42
1:D:130:LEU:HD23	1:D:133:VAL:CG1	2.49	0.41
1:B:112:SER:HA	1:B:139:PRO:HD3	2.02	0.41
1:A:146:VAL:HG12	1:A:146:VAL:O	2.21	0.41
1:B:286:TYR:CE1	1:B:287:MET:HG3	2.56	0.41
1:B:130:LEU:HD23	1:B:133:VAL:CG1	2.51	0.41
1:C:140:PHE:O	1:C:141:PHE:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:182:ARG:HG2	1:D:182:ARG:HH11	1.86	0.41
1:A:48:THR:HA	1:A:207:TYR:CD1	2.56	0.41
1:B:286:TYR:CZ	1:B:287:MET:HG3	2.56	0.40
1:A:241:HIS:NE2	1:A:257:LEU:O	2.53	0.40
1:B:54:TYR:O	1:B:58:MET:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/315 (92%)	278 (96%)	11 (4%)	1 (0%)	41	31
1	B	290/315 (92%)	282 (97%)	8 (3%)	0	100	100
1	C	280/315 (89%)	274 (98%)	6 (2%)	0	100	100
1	D	289/315 (92%)	283 (98%)	6 (2%)	0	100	100
All	All	1149/1260 (91%)	1117 (97%)	31 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/255 (93%)	237 (100%)	1 (0%)	91	91
1	B	236/255 (92%)	231 (98%)	5 (2%)	53	46
1	C	231/255 (91%)	228 (99%)	3 (1%)	69	66
1	D	239/255 (94%)	234 (98%)	5 (2%)	53	46
All	All	944/1020 (92%)	930 (98%)	14 (2%)	65	61

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	244	ARG
1	B	194	SER
1	B	243	MET
1	B	244	ARG
1	B	285	SER
1	B	290	GLN
1	C	106	VAL
1	C	243	MET
1	C	244	ARG
1	D	38	ARG
1	D	104	ARG
1	D	243	MET
1	D	244	ARG
1	D	277	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	GLN
1	D	290	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/315 (93%)	0.16	6 (2%) 65 68	21, 28, 44, 70	0
1	B	292/315 (92%)	0.11	7 (2%) 59 62	21, 29, 50, 87	0
1	C	284/315 (90%)	0.37	7 (2%) 57 60	23, 32, 52, 69	0
1	D	291/315 (92%)	0.14	5 (1%) 70 72	22, 28, 45, 64	0
All	All	1161/1260 (92%)	0.20	25 (2%) 62 65	21, 29, 49, 87	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	191	MET	6.6
1	B	192	ALA	5.0
1	A	200	ILE	4.5
1	C	10	VAL	4.0
1	A	196	VAL	4.0
1	D	286	TYR	3.8
1	A	199	GLY	3.7
1	B	286	TYR	3.4
1	A	198	ASP	3.3
1	B	194	SER	3.2
1	B	12	VAL	2.9
1	D	3	THR	2.9
1	C	13	PRO	2.8
1	D	305	ALA	2.7
1	C	199	GLY	2.6
1	A	186	ALA	2.5
1	C	302	ARG	2.5
1	B	305	ALA	2.4
1	A	177	TRP	2.3
1	D	243	MET	2.3
1	C	35	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	64	HIS	2.2
1	D	307	LEU	2.2
1	B	308	ALA	2.2
1	C	224	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.