



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2023 – 01:33 PM EDT

PDB ID : 8G0F  
Title : Crystal structure of diphtheria toxin H223Q/H257Q double mutant (pH 5.5)  
Authors : Lovell, S.; Kashipathy, M.M.; Battaile, K.P.; Ladokhin, A.S.  
Deposited on : 2023-01-31  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.33  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.33


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	 3% 85% 10% 5%
1	B	538	 4% 84% 10% 5%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7673 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diphtheria toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	511	3759	2384	635	727	13	0	0	0
1	B	511	3747	2379	628	727	13	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	GLU	LYS	engineered mutation	UNP Q5PY51
A	148	LYS	GLU	engineered mutation	UNP Q5PY51
A	223	GLN	HIS	engineered mutation	UNP Q5PY51
A	257	GLN	HIS	engineered mutation	UNP Q5PY51
A	536	MET	-	expression tag	UNP Q5PY51
A	537	ALA	-	expression tag	UNP Q5PY51
B	51	GLU	LYS	engineered mutation	UNP Q5PY51
B	148	LYS	GLU	engineered mutation	UNP Q5PY51
B	223	GLN	HIS	engineered mutation	UNP Q5PY51
B	257	GLN	HIS	engineered mutation	UNP Q5PY51
B	536	MET	-	expression tag	UNP Q5PY51
B	537	ALA	-	expression tag	UNP Q5PY51


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	85	Total	O	0	0
			85	85		
2	B	82	Total	O	0	0
			82	82		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Diphtheria toxin



- Molecule 1: Diphtheria toxin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.55Å 69.47Å 72.88Å 117.99° 93.87° 99.34°	Depositor
Resolution (Å)	36.31 – 2.25 46.78 – 2.25	Depositor EDS
% Data completeness (in resolution range)	95.8 (36.31-2.25) 95.8 (46.78-2.25)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 2.24Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.190 , 0.257 0.194 , 0.256	Depositor DCC
$R_{free}$ test set	2611 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.5	Xtrriage
Anisotropy	0.491	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 50.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7673	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/3833	0.62	0/5217
1	B	0.48	0/3821	0.60	0/5202
All	All	0.49	0/7654	0.61	0/10419

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3759	0	3581	27	0
1	B	3747	0	3563	27	0
2	A	85	0	0	3	0
2	B	82	0	0	0	0
All	All	7673	0	7144	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:458:ARG:NH2	2:A:602:HOH:O	2.26	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:440:LYS:NZ	1:B:498:LYS:O	2.29	0.61
1:A:251:HIS:CE1	1:A:255:LEU:HD22	2.36	0.60
1:B:5:VAL:HG23	1:B:96:VAL:HG22	1.85	0.57
1:A:114:LEU:O	1:A:118:VAL:HG13	2.06	0.56
1:A:341:ALA:O	1:A:353:ILE:HD11	2.07	0.55
1:B:519:ASP:O	1:B:521:THR:HG22	2.06	0.54
1:A:355:PHE:CE2	1:A:360:PHE:HB2	2.42	0.54
1:A:437:ILE:O	1:A:439:GLY:N	2.42	0.53
1:B:5:VAL:HG22	1:B:5:VAL:O	2.08	0.52
1:A:260:LEU:HA	1:A:331:GLN:OE1	2.10	0.52
1:B:406:ILE:HB	1:B:533:ILE:HD13	1.91	0.52
1:A:536:MET:O	1:A:537:ALA:HB3	2.10	0.51
1:A:315:GLY:HA3	1:A:323:HIS:CD2	2.48	0.49
1:A:24:LYS:NZ	1:A:65:TYR:O	2.42	0.49
1:A:353:ILE:O	1:A:353:ILE:HG23	2.11	0.49
1:A:377:ARG:HD3	2:A:606:HOH:O	2.13	0.49
1:A:166:ASN:CG	1:A:169:THR:HG23	2.34	0.48
1:B:104:LYS:O	1:B:107:GLY:N	2.38	0.48
1:A:398:TRP:HH2	1:A:509:ILE:HD12	1.78	0.48
1:B:392:ASP:OD2	1:B:516:LYS:NZ	2.44	0.48
1:A:180:GLU:OE1	2:A:601:HOH:O	2.20	0.47
1:A:73:LEU:HD13	1:A:301:THR:HG22	1.95	0.47
1:B:51:GLU:OE1	1:B:153:TRP:NE1	2.47	0.47
1:B:115:MET:SD	1:B:136:LEU:HD12	2.54	0.47
1:B:413:GLU:OE2	1:B:492:HIS:ND1	2.39	0.46
1:A:396:VAL:HA	1:A:419:LYS:O	2.15	0.46
1:B:301:THR:HA	1:B:363:SER:OG	2.17	0.45
1:B:355:PHE:CE1	1:B:360:PHE:HB2	2.52	0.44
1:B:240:GLU:O	1:B:244:LYS:HB2	2.18	0.44
1:B:518:VAL:O	1:B:521:THR:HG23	2.17	0.44
1:A:118:VAL:HA	1:A:123:PHE:CG	2.53	0.43
1:B:426:PRO:HA	1:B:479:VAL:O	2.19	0.43
1:B:234:PRO:O	1:B:236:LYS:N	2.52	0.43
1:A:250:PHE:CE2	1:A:342:GLN:HG3	2.54	0.43
1:A:315:GLY:O	1:A:322:HIS:HB2	2.19	0.43
1:A:504:ILE:HG23	1:A:504:ILE:O	2.18	0.43
1:B:28:VAL:HG11	1:B:166:ASN:ND2	2.34	0.43
1:A:342:GLN:O	1:A:345:PRO:HD2	2.19	0.42
1:B:407:ARG:HA	1:B:534:LYS:O	2.19	0.42
1:A:451:SER:O	1:A:485:ALA:HA	2.20	0.42
1:A:531:PHE:CE1	1:A:533:ILE:HD11	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:VAL:HG11	1:A:166:ASN:ND2	2.34	0.42
1:B:518:VAL:O	1:B:519:ASP:CB	2.68	0.42
1:B:200:SER:O	1:B:201:CYS:O	2.38	0.41
1:B:349:GLU:O	1:B:350:LEU:HG	2.20	0.41
1:A:306:ILE:HD12	1:B:515:GLN:HB2	2.02	0.41
1:A:291:SER:OG	1:A:350:LEU:HD23	2.20	0.41
1:B:244:LYS:O	1:B:248:GLU:HG3	2.21	0.41
1:B:52:GLY:HA2	1:B:153:TRP:CD1	2.56	0.41
1:B:349:GLU:C	1:B:350:LEU:HD12	2.40	0.41
1:B:451:SER:O	1:B:485:ALA:HA	2.20	0.41
1:B:315:GLY:HA3	1:B:323:HIS:CD2	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	503/538 (94%)	475 (94%)	25 (5%)	3 (1%)	25	25
1	B	503/538 (94%)	475 (94%)	25 (5%)	3 (1%)	25	25
All	All	1006/1076 (94%)	950 (94%)	50 (5%)	6 (1%)	25	25

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	201	CYS
1	A	438	PRO
1	A	68	ASP
1	B	519	ASP
1	B	347	VAL
1	A	347	VAL



### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/454 (85%)	374 (97%)	11 (3%)	42	51
1	B	384/454 (85%)	371 (97%)	13 (3%)	37	45
All	All	769/908 (85%)	745 (97%)	24 (3%)	40	49

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	SER
1	A	109	SER
1	A	133	ARG
1	A	170	ARG
1	A	291	SER
1	A	336	SER
1	A	475	SER
1	A	496	SER
1	A	501	SER
1	A	521	THR
1	A	536	MET
1	B	14	MET
1	B	18	SER
1	B	48	ASP
1	B	49	ASP
1	B	133	ARG
1	B	219	SER
1	B	364	ILE
1	B	467	ASP
1	B	494	SER
1	B	495	SER
1	B	521	THR
1	B	535	SER
1	B	536	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	511/538 (94%)	0.19	17 (3%) 46 48	31, 52, 82, 96	0
1	B	511/538 (94%)	0.22	22 (4%) 35 37	31, 51, 79, 108	0
All	All	1022/1076 (94%)	0.21	39 (3%) 40 43	31, 52, 81, 108	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	123	PHE	5.3
1	A	124	ILE	4.1
1	B	200	SER	3.5
1	B	111	THR	3.4
1	B	242	LYS	3.2
1	A	123	PHE	3.2
1	A	395	ALA	3.1
1	A	420	ILE	3.1
1	A	396	VAL	3.0
1	B	347	VAL	2.9
1	B	238	VAL	2.8
1	A	126	ARG	2.8
1	A	346	LEU	2.8
1	B	126	ARG	2.7
1	B	124	ILE	2.7
1	A	422	ALA	2.7
1	A	134	VAL	2.7
1	A	421	THR	2.7
1	B	107	GLY	2.5
1	A	238	VAL	2.5
1	A	479	VAL	2.5
1	B	127	PHE	2.5
1	A	246	TYR	2.5
1	A	111	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	109	SER	2.4
1	B	395	ALA	2.3
1	A	231	SER	2.2
1	B	4	ASP	2.2
1	B	102	ILE	2.2
1	B	346	LEU	2.2
1	B	348	GLY	2.2
1	B	485	ALA	2.1
1	B	120	THR	2.1
1	A	485	ALA	2.1
1	B	388	PRO	2.1
1	B	345	PRO	2.1
1	B	108	LEU	2.0
1	A	398	TRP	2.0
1	B	420	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.